

This is the post print version of the article, which has been published in Journal of the international neuropsychological society . 2018 24 (6), 617-628. <http://dx.doi.org/10.1017/S1355617718000176>.

The Occurrence and Recovery of Different Neglect-Related Symptoms in Right Hemisphere Infarct Patients during a One-Year Follow-Up

TamPub This document has been downloaded from TamPub.uta.fi
The Institutional Repository of University of Tampere

Laura Nurmi¹, Eija-Inkeri Ruuskanen¹, Mari Nurmi¹, Anna-Maija Koivisto², Anna-Kaisa Parkkila³,
Heikki Numminen³, Prasun Dastidar⁴ and Mervi Jehkonen¹

¹Faculty of Social Sciences, Psychology, University of Tampere, Tampere, Finland

²Faculty of Social Sciences, Health Sciences, University of Tampere, Tampere, Finland

³Department of Neurology and Rehabilitation, Tampere University Hospital, Tampere, Finland

⁴Department of Radiology, Tampere University Hospital, Tampere, Finland

Corresponding author: Laura Nurmi, University of Tampere, Faculty of Social Sciences, Psychology,
FIN-33014 University of Tampere, Finland. E-mail: laura.j.nurmi@uta.fi

Word count for the abstract: 250

Word count for the manuscript text: 5030

ABSTRACT

Objective: To examine the occurrence of and recovery from visual neglect-related symptoms with the focus on neglect laterality, ipsilateral orienting bias and slowed processing speed in right hemisphere (RH) infarct patients during a one-year follow-up. Furthermore, to propose guidelines for assessing processing speed alongside the Behavioural Inattention Test (BIT).

Method: We studied three RH patient groups: neglect (N+), mild left inattention (MLI+) and non-neglect (N-) patients, and healthy controls. The BIT with some additional analyses was conducted at the acute phase and at six and 12 months.

Results: The N+ group's BIT score increased and originally lateralized omissions became more evenly distributed during the follow-up. The N+ and MLI+ groups' starting points were more rightward located than the healthy group's at the acute phase and at six, and partly at 12 months. Patient groups were slower than the controls in performing cancellation tests at the acute phase. The N+ and MLI+ groups remained slower than the controls throughout the follow-up.

Conclusions: During the first year after RH infarct originally left-sided manifestation of neglect shifted towards milder non-lateralized attentional deficit. Ipsilateral orienting bias and slowed processing speed appeared to be rather persistent neglect-related symptoms both in neglect patients and patients with initially milder inattention. We propose some effortless, tentative ways of examining processing speed and ipsilateral orienting bias alongside the BIT in order to better recognise these neglect-related symptoms, and highlight the need to assess and treat patients with initially milder inattention, who have been under-recognized and under-treated in clinical work.

Keywords: inattention, infarct, processing speed, ipsilateral orienting bias, stroke, visual neglect

INTRODUCTION

Neglect indicates a failure in reporting, responding or orienting to contralesional stimuli that cannot be explained by primary sensory or motor deficits (Heilman, Watson, & Valenstein, 1993; Robertson & Halligan, 1999). Visual neglect commonly occurs in consequence of right hemisphere (RH) lesions. Two major symptoms are a tendency to initially orient attention to the ipsilesional side (Butler, Lawrence, Eskes, & Klein, 2009; Gainotti, De Luca, Figliozzi, & Doricchi, 2009; Gainotti, D’Erme, & Bartolomeo, 1991; Karnath & Rorden, 2012; Nurmi et al., 2010) and a deficit in orienting attention towards the contralesional side (e.g. Corbetta & Shulmann, 2011; Danckert & Ferber, 2005; Karnath & Rorden, 2012). Patients with left visual neglect tend to orient first to the right hemispace, and then fail to reorient towards the left (Robertson & Eglin, 1993). Besides lateralized symptoms, neglect is known to involve non-lateralized attentional deficits, which are also recognized as essential to neglect (Husain & Rorden, 2003; van Kessel, van Nes, Brouwer, Geurts, & Fasotti, 2010; Manly, 2002; Robertson, 1993). According to Karnath’s (1988) model, there are three components that contribute to neglect: a) a tendency to initially orient attention to the ipsilesional side, b) a deficit in shifting attention from the ipsilesional side towards the contralesional side, and c) a generalised (non-lateralized) decline in information processing and attention capacity. It has been suggested that the lateralized and the non-lateralized components interact with each other to intensify neglect symptoms (van Kessel et al., 2010; Manly, 2002; Ting et al., 2011) and to impair recovery from neglect (Husain & Rorden, 2003; Robertson, 1993, 2001; Samuelsson Hjelmquist, Jensen, Ekholm, & Blomstrand, 1998).

Adequate processing speed is essential in many cognitive functions, and slowed processing speed is often thought to underlie attentional deficits (Lezak, 2012). Slow processing speed has been associated with RH lesions (Farne et al., 2004; Heilman, Watson, & Valenstein, 1993) and to an even greater extent with neglect (Erez, Katz, & Ring, 2009; Bonato, 2012; Gerritsen, Berg, Deelman, Visser-

Keizer, & Jong, 2003). Several studies have demonstrated neglect-related slowness of perceptual processing using measures of reaction time (Bartolomeo, 1997; Bartolomeo & Chokron, 1999, 2002; Behrmann & Meegan, 1998; Erez et al., 2009; van Kessel et al., 2010; Samuelsson et al., 1998; Smania et al., 1998), attentional blink (Danckert & Ferber, 2005; Husain & Rorden, 2003) and cancellation rate (Manly et al., 2009; Robertson, 1993; Robertson & Eglin, 1993). Some studies have reported evidence of slow contralateral visual processing, which has been thought to reflect lateralized deficits, i.e. ipsilateral orienting and resultant slow attending to the left (e.g. Behrmann & Meegan, 1998; Smania et al., 1998). Other studies have described this slowness as ipsilateral or bilateral, and interpreted it as a limitation of non-lateralized attentional processing capacity (Bartolomeo & Chokron, 2002; van Kessel et al., 2010; Robertson, 1993).

Conventional neglect tests typically measure the deficit in orienting attention to the contralesional side, which may manifest as left-sided omissions in cancellation tasks (Heilman, Watson & Valenstein, 1993; Husain & Rorden, 2003; Karnath & Rorden, 2012; van Kessel et al. 2010). The Behavioural Inattention Test (BIT) that includes six conventional paper-pencil subtests (BITC) and nine behavioral subtests (Jehkonen, 2002; Wilson, Cockburn, & Halligan, 1987) has been shown to be a valid and reliable method for examining visual neglect (Wilson et al., 1987). In determining the laterality of attentional deficits the BITC can be further enhanced by measuring the Center of Cancellation (CoC) to define the mean horizontal location of the cancelled targets (Binder, Marshall, Lazar, Benjamin, & Mohr, 1992; Rorden & Karnath, 2010). However, neither the BIT nor other traditional methods usually acknowledge the other components of neglect described above, i.e. a tendency to initially orient attention to the ipsilesional side (here referred to as initial ipsilateral orienting bias) and slowed processing speed. The only exception that we are aware of is the Bell's test in a standardized French battery (*Batterie d'Evaluation de la Négligence*) that measures ipsilateral orienting bias (Azouvi et al., 2006). Therefore most conventional methods may be insensitive in

assessing less severe forms of neglect. Indeed, several studies have shown that at the acute phase of stroke, some RH patients who do not exhibit neglect in conventional measures do nonetheless demonstrate initial ipsilateral orienting bias (Erez et al., 2009; Gainotti et al., 1991; Jalas, Lindell, Brunila, Tenovuo, & Hämäläinen, 2002; Nurmi et al., 2010) and/or slowed processing speed (van Kessel et al., 2010). Similarly, many patients who according to traditional tests have recovered from neglect still show initial ipsilateral orienting bias (Bartolomeo, 1997; Bartolomeo & Chockron, 2002; Bonato, 2012; Kettunen, Nurmi, Dastidar, & Jehkonen, 2012; Mattingley, Bradshaw, Bradshaw, & Nettleton, 1994) and/or slowed processing speed (Bonato, 2012; Friedrich & Margolin, 1993; Robertson & Eglin, 1993; Samuelsson et al., 1998; Taylor, 2003; Viken, 2013) as residual neglect symptoms long after stroke.

It appears then that more sensitive methods are needed to assess mild or residual neglect. To this end it has been suggested that the starting point (SP) in cancellation tasks is highly sensitive in detecting initial ipsilateral orienting bias (Bonato, 2012; Nurmi et al. 2010; Samuelsson, Hjelmquist, Naver, & Blomstrand, 1996; Ting et al., 2011), According to Azouvi et al. (2002, 2006) it is in fact the most sensitive paper-and-pencil measure of neglect. In contrast to healthy controls, who typically start cancellation from left side (Nurmi et al., 2010; Robertson & Eglin, 1993; Warren, Moore, & Vogtle, 2008), neglect patients mainly start it from the right (Adair & Barret, 2008; Heilmann et al., 1993). Furthermore, it has been suggested that the inclusion of processing speed measurement might increase the diagnostic sensitivity of the assessment for mild neglect symptoms at the acute phase (Erez et al., 2009; van Kessel et al., 2010; Ting et al., 2011) as well as for residual neglect symptoms during the recovery (Bartolomeo, 1997; Bonato, 2012; Friedrich & Margolin, 1993; Taylor, 2003).

As mentioned above, it appears that different neglect-related symptoms do not recover at the same rate. Karnath (1988) has proposed that out of the three components of visual neglect, inability to orient attention towards the contralesional side recovers faster than ipsilateral orienting bias or general

attentional deficits. Indeed, when assessed with conventional methods that mainly focus on measuring the ability to orient attention contralaterally, it seems that in most patients neglect recovers within the first few months after RH stroke (e.g. Bonato, 2012; Cassidy, Lewis, & Grey, 1998; Harvey & Gilchrist, 2002), although recovery may fluctuate during the first year (Jehkonen, Laihosalo, Koivisto, Dastidar, & Ahonen, 2007). Initial ipsilateral orienting bias, on the other hand, is often present for six to seven (Bonato, 2012; Kettunen et al., 2012; Samuelsson et al., 1996), even 12 months (Mattingley et al., 1994) after stroke, despite the recovered ability to reorient attention towards the contralesional side. Similarly, slow processing speed has been seen as a residual symptom in clinically recovered neglect patients for three to seven months post-stroke (Harvey & Gilchrist, 2002; Samuelsson et al., 1998; Viken, 2013). It has also been shown that even if patients no longer show neglect symptoms in simpler tasks, attentionally more demanding situations, as often encountered in daily activities, may still provoke residual symptoms such as ipsilateral orienting bias (Friedrich & Margolin, 1993; Taylor, 2003).

The presence of neglect following stroke is known to predict poor functional recovery and rehabilitation outcomes as well as difficulties in activities of daily living (ADL; Buxbaum et al., 2004; Chen, Hreha, Kong, & Barret, 2015; Cherney, Halper, Kwasnica, Harvey, & Zhang, 2001; Di Monaco et al., 2011; Gillen, Tennen, & McKee, 2005; Jehkonen, 2002b; Katz, Hartman-Maeir, Ring, & Soroker, 1999). Viken, Jood, Jern, Blomstrand and Samuelsson (2014) found that in RH patients, among all the neglect-related symptoms present at the acute phase of stroke, ipsilateral orienting bias was the most important predictor of functional dependency at three months and slow visual processing speed at two years post-stroke. Additionally, even when full-blown neglect symptoms are absent, milder deficits such as ipsilateral orienting bias and slow processing speed *per se* cause difficulties in more complex ADL such as wheelchair manoeuvring and participating in traffic (e.g. driving, crossing roads), thus increasing the risk of accidents and impairing and endangering functioning in real-life

situations (Bonato, 2012; Suzuki, Chen, & Kondo, 1997; Webster et al., 1995).

This study set out to explore the occurrence and recovery of different neglect-related symptoms, focusing particularly on neglect laterality, initial ipsilateral orienting bias and slowed processing speed in visual search. These symptoms were examined during a one-year follow-up in RH infarct patients, divided into three groups based on the acute phase symptoms: a neglect (N+) group, a mild left inattention (MLI+) group, and a non-neglect (N-) group. These patient groups were also compared with healthy controls. The following hypotheses were set: 1) in the N+ group, originally left-sided manifestation of neglect shifts towards milder non-lateralized attentional deficit during the one-year follow-up; 2) both the N+ and MLI+ groups show initial ipsilateral orienting bias throughout the follow-up period; and 3) all RH patient groups show slowed processing speed in visual search at the acute phase, and the N+ and MLI+ groups continue to show this during the follow-up period. Our particular focus was to examine the occurrence of symptoms and their recovery in patients with MLI+, who cannot be detected with conventional neglect tests in clinical work and who have received minor attention in neglect research. Neglect assessment was based on the BIT with additional analyses of CoC and SP. In addition, we aimed to further enhance this conventional method by creating guidelines for assessing slowed processing speed using the BIT cancellation tasks.

METHODS

Subjects

The patient group included 65 first-ever RH infarct patients consecutively admitted to Tampere University Hospital during two different periods, from January 2007 to January 2008 and from March 2010 to December 2012. The infarct diagnosis was based on the neurological and neuroradiological

examinations. Patients who met the criteria for thrombolysis received the treatment within 4.5 hours of infarct onset, others were treated according to a conservative treatment protocol. Patients with a neurological diagnosis other than RH infarct, previous neurological or psychiatric diagnosis, significant cerebral atrophy, significant loss of consciousness, significant loss of primary vision or hearing, age < 30 or > 85 years, substance abuse, native language other than Finnish and inability to live independently prior to infarct, were excluded from the study. Informed consent was obtained before inclusion in the study. The patients underwent three neurological examinations: on arrival at hospital (baseline), at the acute phase (within a few days from onset) and at six months. Two neuroradiological examinations were carried out: a CT on arrival at hospital and a MRI at the acute phase or at six months. In addition, three neuropsychological examinations were conducted: at the acute phase, at six and at 12 months. No recurrent strokes emerged during the follow-up period. Rehabilitation status (received/not received during the one-year follow-up) regarding neuropsychological rehabilitation and occupational therapy was obtained from all patients.

The control group consisted of 40 healthy volunteers aged 30 to 80 years. They were gathered from among researchers' acquaintances and elderly social clubs. Subjects with a history of neurological or psychiatric disorders were not included. One neuropsychological examination was conducted. All subjects were blind to the hypotheses of the study. The study protocol was approved by the Ethical Committee of Tampere University Hospital and the human data included in this study was obtained in compliance with the Helsinki Declaration.

Methods

Neuropsychological examination

In the neuropsychological examination visual neglect was assessed with the BITC, laterality of neglect

with the CoC, initial ipsilateral orienting bias with the SP and processing speed with cancellation time. A comprehensive description of the neuropsychological methods is given in Appendix 1.

Neurological and neuroradiological examinations

The neurological and neuroradiological methods are described in Appendix 2.

Data Analysis

Nonparametric methods were used for data analysis due to the skewed distribution of the variables. For continuous variables, comparisons between several independent groups were performed with the Kruskal-Wallis test. Multiple pairwise comparisons were made with the Mann-Whitney U test. Bonferroni corrections were made by multiplying the p -value by six (comparisons between four groups) or by three (comparisons between three patient groups). Categorical variables were analyzed with the chi-square test. Effect size was calculated for Kruskal-Wallis analyses by computing η^2 and for Mann-Whitney U analyses by computing r (Tomczak & Tomczak, 2014). Cohen's guidelines for r were used (Coolican, 2009; a large effect: .5, a medium effect: .3, and a small effect: .1). The guideline cutoff for slowed processing speed in visual search was defined separately for the letter and star cancellation tasks by receiver operating characteristic (ROC) curve analyses. Adequate versus slowed processing speed was differentiated by comparing the healthy subjects' processing speed with that of the N+ patients, and by calculating sensitivity, specificity and areas under the curve (AUC). Guideline cutoffs were based on the best combination of sensitivity and specificity. Statistical analyses were conducted with the Statistical Package for the Social Sciences version 23 for Windows. The level of statistical significance was set at .05.

RESULTS

Subject characteristics

Eighteen patients (28%) had visual neglect (N+; failure in at least two BITC subtests) and 23 patients (35%) met the criteria for ipsilateral orienting bias (MLI+; atypical SP in at least two cancellation tasks). Twenty-four patients (37%) did not show neglect symptoms (N-). Characteristics for the patient groups and the healthy group are presented in Table 1. The groups did not differ significantly on any subject characteristics at any point of measurement. Forty percent of N+, 10% of MLI+ and 0% of N- patients received rehabilitation during the one-year follow-up.

Neurological characteristics and the infarct locations for the patient groups are presented in Table 2. Stroke severity (NIHSS) differed significantly between the groups at baseline and at the acute phase. At both times the N+ group had more severe stroke than the other groups. There were significant differences between the patient groups in basic ADL (BI) at the acute phase and at six months. At the acute phase the N+ group had greater dependence in basic ADL than the other groups. At six months the N+ group was still more dependent in basic ADL than the N- group. In all groups the most typical infarct location was in the region of MCA.

[INSERT TABLE 1 HERE]

[INSERT TABLE 2 HERE]

Visual neglect and laterality

Median and quartiles of the BIT and CoC sum scores for the four groups and between-group comparisons at the three points of measurement are presented in Tables 3 and 4. Significant group

differences were found in the BIT sum scores at the acute phase and at 12 months and in the CoC sum scores at the acute phase. At the acute phase, the N+ group had a significantly lower BIT sum score than the other groups (large effects) and a significantly more rightward CoC sum score than the H group (medium effects). At 12 months, the N+ group had a significantly lower BIT sum score than the N- and MLI+ groups (medium effects).

[INSERT TABLE 3 HERE]

[INSERT TABLE 4 HERE]

Individual BIT and CoC values for each patient are presented in Appendix 3. Based on failure in at least two BITC subtests, visual neglect was still present in 21% of N+ patients at six months and in 36% of N+ patients at 12 months.

Ipsilateral orienting bias

Median and quartiles of SPs for line and star cancellation tasks in the four groups and between-group comparisons at three points of measurement are presented in Table 5. There were statistically significant group differences in the SPs for both cancellation tasks at each point of measurement. At the acute phase, the SPs for line and star cancellation tasks were significantly more rightward located in the N+ and MLI+ groups than in the N- and healthy groups (large effects). At six months, the SPs for the same tasks were more rightward located in the N+ and MLI+ groups than in the healthy group, and in the MLI+ group than in the N- group (medium to large effects). At 12 months, the SPs for the line and star cancellation tasks were located significantly more to the right in the MLI+ group than in the healthy group (medium effect). The SPs of the N+ and MLI+ groups did not differ significantly from each other at any point of measurement.

[INSERT TABLE 5 HERE]

Individual SP values for each patient are presented in Appendix 3. Based on atypical SP in at least two cancellation tasks, ipsilateral orienting bias still occurred in 43% of N+ and 50% of MLI+ patients at six months, and in 29% of N+ and 32% of MLI+ patients at 12 months.

Processing speed in visual search

Median and quartiles of cancellation time for the letter and star cancellation tasks in the four groups and between-group comparisons at the three points of measurement are presented in Table 6. There were statistically significant cancellation time differences between the four groups in both cancellation tasks at each point of measurement. At the acute phase, cancellation time for the letter cancellation task was significantly slower in the MLI+ and N+ groups than in the healthy group (medium effects). Cancellation time for the star cancellation task was significantly slower in all three patient groups than in the healthy group (medium to large effects). At six months, cancellation time for both cancellation tasks was slower in the MLI+ and N+ groups than in the healthy group (medium to large effects). At 12 months, cancellation time for the letter cancellation task was slower in the N+ group than in the N- and healthy groups (medium effects). Cancellation time for the star cancellation task was slower in the MLI+ and N+ groups than in the healthy group (medium effects). The cancellation times of the N+ and MLI+ groups did not differ significantly from each other at any point of measurement.

[INSERT TABLE 6 HERE]

The ROC analyses of processing speed for healthy subjects versus N+ patients in the letter and star cancellation tasks are shown in Table 7. For the letter cancellation task, the ideal pairing of sensitivity (83%) and specificity (75%) was found with a cutoff of > 82.5 seconds for slowed processing speed (Figure 1a). For the star cancellation task, the ideal pairing of sensitivity (83%) and specificity (83%) was found with a cutoff of > 59.5 seconds for slowed processing speed (Figure 1b).

Because of its higher specificity, we chose to use the star cancellation task for assessing slowed processing speed in individual patients. Thus a cancellation time of 60 seconds or more in the star cancellation task was defined as an indicator of slowed processing speed in visual search.

Individual cancellation times for each patient are presented in Appendix 3. Based on the guideline cutoff, slowed processing speed was found in 83% of N+, 83% of MLI+ and 58% of N- patients at the acute phase, in 57% of N+, 55% of MLI+ and 35% of N- patients at six months, and in 71% of N+, 48% of MLI+ and 16% of N- patients at 12 months.

[INSERT TABLE 7 HERE]

[INSERT FIGURES 1a AND 1b HERE]

DISCUSSION

The aim of our study was to investigate the occurrence of and recovery from neglect-related symptoms, focusing on neglect laterality, ipsilateral orienting bias and slowed processing speed in visual search during a one-year follow-up. The RH patients were divided into N+, MLI+ and N- groups according to their acute phase symptoms and compared with healthy controls. Our particular interest was to examine the symptoms of patients with MLI+, who often remain undetected in clinical work and who have received less attention in neglect research. In addition, we aimed to further enhance the BIT by defining a guideline for assessing slowed processing speed using the cancellation tasks.

At the acute phase of RH infarct, 18 patients (28%) met the criterion for visual neglect (N+). During the one-year follow-up overall neglect decreased according to both the BIT and CoC sum scores. However, we also saw some fluctuation in recovery based on the group comparisons with BIT sum scores and the proportions of individual N+ patients showing neglect, which is in line with the previous findings of Jehkonen et al. (2007). On the other hand, the CoC sum scores indicated that at

follow-ups, omissions in the N+ group were no longer located more leftward than in the healthy group. As hypothesized, then, neglect manifested as left-sided only at the acute phase, after which the remaining omissions became more evenly distributed, indicating a shift towards milder non-lateralized deficit in attention. These results support Karnath's (1988) suggestion that the inability to orient attention towards the contralesional side recovers faster than the general attentional deficits of neglect.

At the acute phase we found initial ipsilateral orienting bias not only in the N+ group, but also in 23 RH patients (35%) who did not show neglect with conventional BIT measures (MLI+). Indeed, the N+ and MLI+ groups started the line and star cancellation tasks from the right side at the acute phase, in contrast to the healthy and N- groups, which started the tasks from the left margin. This is in line with previous studies showing that ipsilateral orienting bias is a common neglect symptom (Butler et al., 2009; Corbetta & Shulmann, 2011; Gainotti et al. 2009; Gainotti et al., 1991; Karnath & Rorden, 2012; Manly et al., 2009; Nurmi et al., 2010) and may also be seen in RH patients without full-blown neglect, indicating less severe neglect symptoms (Erez et al., 2009; Gainotti et al., 1991; Jalas et al., 2002; Nurmi et al., 2010). During the first six months the N+ and MLI+ groups' SPs shifted somewhat leftwards, but still located more to the right than in the healthy group, as hypothesized. At 12 months, however, only the MLI+ group's SPs were significantly more rightward than in the healthy group, while just a borderline difference occurred between the N+ and healthy groups in the star cancellation task. Nevertheless almost one-third of both the N+ and MLI+ patients still showed ipsilateral orienting bias at 12 months. These results support Karnath's (1988) suggestion that persisting ipsilateral orienting bias is one component of residual deficits in neglect patients who have recovered abilities of contralesional orienting. They also support previous findings that neglect patients may show ipsilateral orienting bias six to 12 months after stroke (Bonato, 2012; Kettunen et al., 2012; Mattingley et al., 1994; Samuelsson et al., 1996). However, only limited scientific attention has previously been devoted to the recovery of patients with initially milder inattention symptoms (i.e. MLI+), but no actual neglect

after stroke. Our results highlight the clinical relevance of these patients as well, since ipsilateral orienting bias proved to be particularly persistent in the MLI+ group.

For purposes of assessing the occurrence of slow processing speed in visual search in individual patients, we created guideline cutoffs in the letter and star cancellation tasks. Because of its better combination of sensitivity and specificity, we chose to focus on the star cancellation alone, in which a cancellation time of 60 seconds or more indicated slowed processing speed. At the acute phase, just over half of the N- patients and a clear majority of the MLI+ and N+ patients showed slow processing speed in visual search. In general, all patient groups were slower than the healthy group at the acute phase, but showed differing courses of recovery during the follow-up, as was hypothesized. The N- group's performance was significantly slower than the healthy group's only in the star cancellation task and only at the acute phase. Indeed, during the follow-up the proportion of slow N- patients fell markedly, and at 12 months reached the same level as in the healthy subjects. In contrast, the MLI+ and N+ groups were slower than the healthy group in both cancellation tasks up to six months, and in the star cancellation task still at 12 months. A little over half of the patients in both groups showed slow cancellation performance at six months, after which the proportion slightly decreased in the MLI+ patients but increased in the N+ patients towards the end of the year.

Altogether, these results are in line with previous studies that have associated slow processing speed with RH lesions in general (Farne et al., 2004; Heilman et al., 1993) and even more so with visual neglect (Erez et al., 2009; Bonato, 2012; Gerritsen et al., 2003). In addition, our results support the previous findings that slow processing speed is a rather persistent residual symptom of neglect and may be seen in patients with clinically recovered neglect for several months (Harvey & Gilchrist, 2002; Samuelsson et al., 1998; Viken, 2014), or as in our study, even a year after stroke. There is hardly any previous research regarding processing speed and its recovery specifically in a subgroup of RH patients with initially milder inattention symptoms after stroke but no actual neglect (i.e. MLI+). Our results

suggest that these patients may also show slowed processing speed in visual search as a persistent residual symptom even for a year after stroke, unlike the RH non-neglect patients (N-) who instead seem to recover quite soon after stroke.

We also obtained information about RH patients' rehabilitation, including neuropsychological rehabilitation and occupational therapy during the follow-up. A clear majority of patients who received rehabilitation were N+ patients, of whom 40% were in rehabilitation during the year after stroke. Only 10% of the MLI+ patients and none of the N- patients received rehabilitation. This may have influenced the recovery of the N+ group. For example, recovery from ipsilateral orienting bias appeared to be slightly better in the N+ than the MLI+ group. Moreover, this finding suggests that RH patients with milder inattention symptoms, but without full-blown neglect, are under-treated in clinical work.

A major strength of our study is a broad perspective on recovery from visual neglect after RH infarct by examining various neglect-related symptoms during a lengthy follow-up period of 12 months. We focused on a homogeneous group of consecutive patients with a first-ever RH infarct. As well as including neglect patients, we showed a particular interest in patients with initially milder inattention symptoms, who have received less attention both in clinical work and in neglect research. Overall, the results of our study were well in line with previous findings, which adds to their reliability. In addition, we propose a novel, potential method for measuring processing speed in visual search by suggesting a guideline cutoff of 60 seconds for slowed processing speed in the BIT star cancellation task. When combined with our previously suggested analysis of SP in the BIT cancellation tasks (Nurmi et al. 2010), this offers an effortless, tentative way of enhancing neglect assessment by adding an examination of processing speed and ipsilateral orienting bias alongside the standard administration of the well-know conventional neglect assessment.

However, it should be noted that our relatively small sample sizes together with the loss of

participants during the follow-up might have had some influence on the results. This applies particularly to the N+ group, which initially was the smallest group and also lost the biggest number of participants. This might go some way towards explaining why some of our hypotheses concerning the N+ group did not receive as strong support as we expected. Furthermore, this is why we were only able to suggest a guideline cutoff rather than establish a firm cutoff point for slowed processing speed in visual search. This will require further research. Paper-pencil measures of processing speed can be susceptible to factors other than processing speed (e.g. perseveration, distraction, lack of motivation), which may also contribute to prolonged test performance. Cancellation time is no exception, and as it has not been traditionally used as a measure of processing speed, the findings must be interpreted with caution. As stated, our results are altogether well in line with previous findings concerning neglect and slow processing speed, but more work is still needed to explore milder inattention symptoms and their association with processing speed and its recovery. Furthermore, previous studies concerning neglect and processing speed have lent support to two contrasting hypotheses; one suggesting that slow processing speed emerges contralaterally (e.g. Behrmann & Meegan, 1998; Smania et al., 1998) reflecting lateralized deficits, and the other that slowness is ipsi- or bilateral and reflects non-lateralized difficulties instead (Bartolomeo & Chokron, 2002; van Kessel et al., 2010; Robertson, 1993). The methods used in this study to measure processing speed do not allow us to determine which type of attentional deficit, lateralized or generalized, lies behind the slowed processing speed associated with neglect. This is another question that calls for further research. The considerable heterogeneity seen in N+ patients' acute phase BIT scores underlines the need to explore the possible association of neglect severity with the occurrence and recovery of different neglect-related symptoms. Finally, rehabilitation was here assessed on a yes/no basis only, but as the content and intensity of rehabilitation may substantially influence the recovery of different neglect-related symptoms, it needs to be considered in more detail in future studies.

In conclusion, our results suggest that during the first year after RH infarct there is a shift from left-sided manifestation of neglect towards milder non-lateralized deficit of attention. In addition, both ipsilateral orienting bias and slowed processing speed appear to be rather persistent neglect-related symptoms that may occur for up to a year after RH infarct in patients with originally full-blown neglect, but also in patients with initially milder inattention symptoms. Both these symptoms should be assessed in the clinical examination of neglect, since they are known to cause difficulties in more complex ADL such as participating in traffic, and are important predictors of functional dependency after stroke. We propose here some effortless, tentative ways of examining processing speed and ipsilateral orienting bias alongside the standard administration of the well-know conventional neglect assessment. The findings of our study highlight the importance of both a sensitive enough neglect assessment and a long enough follow-up for patients after RH infarct in order to better recognise and treat different neglect-related symptoms in neglect patients and in patients with initially milder inattention symptoms who both may suffer from significant residual symptoms long after stroke.

Acknowledgements

This study was supported by grants from the Finnish Cultural Foundation (00130649) and the Competitive State Research Financing of the Expert Responsibility Area of Tampere University Hospital (9M023).

Conflicts of interest

The authors have no conflicts of interest.

REFERENCES

- Adair, J.C., & Barrett, A.M. (2008). Spatial neglect clinical and neuroscience review: a wealth of information on the poverty of attention. *Annals of the New York Academy of Sciences*, 1142, 21–43.
- Azouvi, P., Bartolomeo, P., Beis, J.M., Perennou, D., Pradat-Diehl, P., & Rousseaux, M. (2006). A battery of tests for the quantitative assessment of unilateral neglect. *Restorative Neurology and Neuroscience*, 24, 273–285.
- Azouvi, P., Samuel, C., Louis-Dreyfus, A., Bernati, T., Bartolomeo, P., Beis, J.M., ... Rousseaux, M. (2002). Sensitivity of clinical and behavioural tests of spatial neglect after right hemisphere stroke. *Journal of Neurology, Neurosurgery, and Psychiatry*, 73, 160–166.
- Bartolomeo, P. (1997). The novelty effect in recovered hemineglect. *Cortex*, 33, 323–332.
- Bartolomeo, P., & Chokron, S. (1999). Left unilateral neglect or right hyperattention? *Neurology*, 53, 2023–2027.
- Bartolomeo, P., & Chokron, S. (2002). Orienting of attention in left unilateral neglect. *Neuroscience and Biobehavioral Reviews*, 262, 217–234.
- Behrmann, M., & Meegan, D.V. (1998). Visuomotor Processing in Unilateral Neglect. *Consciousness and Cognition*, 7, 381–409.

- Benton, A. (1986). Reaction time in brain disease: some reflections. *Cortex*, 22, 129–140.
- Binder, J., Marshall, R., Lazar, R., Benjamin, J., & Mohr, J.P. (1992) Distinct syndromes of hemineglect. *Archives of Neurology*, 49, 1187–1194.
- Bonato, M., Priftis, K., Marenzi, R., Umiltà, C., & Zorzi, M. (2010). Increased attentional demands impair contralesional space awareness following stroke. *Neuropsychologia* 48, 3934–3940.
- Bonato, M. (2012). Neglect and extinction depend greatly on task demands: a review. *Frontiers in Human Neuroscience*, 6: 195.
- Butler, B.C., Lawrence, M., Eskes, G.A., & Klein, R. (2009). Visual search patterns in neglect: Comparisons of peripersonal and extrapersonal space. *Neuropsychologia*, 47, 869–878.
- Buxbaum, L.J., Ferraro, M.K., Veramonti, T., Farne, A., Whyte, J., Lavadas, E., Frassinetti, F., & Coslett, H.B. (2004). Hemispatial neglect subtypes, neuroanatomy, and disability. *Neurology*, 62, 749–756.
- Cassidy, T.P., Lewis, S., & Grey, C.S. (1998). Recovery from visuospatial neglect in stroke patients. *Journal of Neurology, Neurosurgery, and Psychiatry*, 64, 555–557.
- Chen, P., Hreha, K., Kong, Y., & Barrett, A.M. (2015). Impact of Spatial Neglect on Stroke Rehabilitation: Evidence from the Setting of an Inpatient Rehabilitation Facility. *Archives of Physical Medicine and Rehabilitation*, 96, 1458-1466.

Cherney, L.R., Halper, A.S., Kwasnica, C.M., Harvey, R.L., & Zhang, M. (2001). Recovery of functional status after right hemisphere stroke: relationship with unilateral neglect. *Archives of Physical Medicine and Rehabilitation*, 82, 322-328.

Coolican, H. (2009). *Research methods and statistics in psychology*. London, United Kingdom: Hodder.

Corbetta, M., & Shulman, L. (2011). Spatial neglect and attention networks. *Annual Review of Neuroscience*, 34, 569–599.

Dankert, J., & Ferber, S. (2006). Perspective Revisiting unilateral neglect. *Neuropsychologia*, 44, 987–1006.

Erez, A.B.-H., Katz, N., Ring, H., & Soroker, N. (2009). Assessment of spatial neglect using computerised feature and conjunction visual search tasks. *Neuropsychological Rehabilitation*, 19, 677–695.

Farne, A., Buxbaum, L., Ferraro, M., Frassinetti, F., Whyte, J., Veramonti, T., ... Làdavas, E. (2004). Patterns of spontaneous recovery of neglect and associated disorders in acute right brain-damaged patients. *Journal of Neurology, Neurosurgery, and Psychiatry*, 75, 1401–1410.

Folstein, M.F., Folstein, S.E., & McHugh, P.R. (1975). Mini-mental state. A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatry Research*, 12, 189–198.

Friedrich, F.J., & Margolin, D.I. (1993) Responses time measures of hemi-inattention: a longitudinal case report. *Neuropsychiatry, Neuropsychology and Behavioral Neurology*, 6, 54-59.

Gainotti, G., De Luca, L., Figliozzi, F., & Doricchi, F. (2009). The influence of distracters, stimulus duration and hemianopia on first saccade in patients with unilateral neglect. *Cortex*, 45, 506–516.

Gainotti, G., D’Erme, P., & Bartolomeo, P. (1991). Early orientation of attention toward the half space ipsilateral to the lesion in patients with unilateral brain damage. *Journal of Neurology, Neurosurgery and Psychiatry*, 54, 1082–1089.

Gerritsen, M.J.J, Berg, I.J., Deelman, B.G, Visser-Keizer, A.C., & Meyboom-de Jong, B. (2003). Speed of Information Processing After Unilateral Stroke. *Journal of Clinical and Experimental Neuropsychology*, 25, 1-13.

Goldstein, L.B., Bertels, C., & Davis, J.N. (1989). Interrater reliability of the NIH stroke scale. *Archives of Neurology*, 46, 660– 662.

Harvey, B.O.M., & Gilchrist, I.D. (2002). First Saccades Reveal Biases in Recovered Neglect. *Neurocase*, 8, 306–313.

Heilman, K.M., Watson, R.T., & Valenstein, E. (1993). Neglect and related disorders. In K.M. Heilman, & E. Valenstein (Eds.), *Clinical Neuropsychology* (pp. 279–336). New York: Oxford University Press.

Husain, M., & Rorden, C. (2003). Non-spatially lateralized mechanisms in hemispatial neglect. *Neuroscience*, 4, 26–36.

Jalas, M.J., Lindell, A.B., Brunila, T., Tenovuo, O., & Hämäläinen, H. (2002). Initial rightward orienting bias in clinical tasks: Normal subjects and right hemispheric stroke patients with and without neglect. *Journal of Clinical and Experimental Neuropsychology*, 24, 479–490.

Jehkonen, M. (2002a). *Behavioural Inattention Test. Manual*. Helsinki: Psykologien Kustannus Oy. [In Finnish].

Jehkonen, M. (2002b). *The role of visual neglect and anosognosias in functional recovery after right hemisphere stroke*. (Doctoral dissertation, University of Tampere, Finland). Retrieved April 28, 2012 from <http://acta.uta.fi/english/teos.php?id=6242>

Jehkonen, M., Laihosalo, M., Koivisto, A.-M., Dastidar, P., & Ahonen, J. P. (2007). Fluctuation in spontaneous recovery of left visual neglect: a 1-year follow-up. *European Neurology*, 58, 210–214.

Karnath, H.O. (1988). Deficits of attention in acute and recovered visual hemi-neglect. *Neuropsychologia*, 26, 27–43.

Karnath, H.-O., & Rorden, C. (2012). The anatomy of spatial neglect. *Neuropsychologia*, 50, 1010–1017.

Katz, N., Hartman-Maeir, A., Ring, H., Soroker, N. (1999). Functional disability and rehabilitation outcome in right hemisphere damaged patients with and without unilateral spatial neglect. *Archives of Physical Medicine and Rehabilitation*, 80, 379–384.

Kessel, M.E., van Nes, I.J.W., Brouwer, W.H., Alexander C.H. Geurts, A.C.H., & Fasotti, L. (2010). Visuospatial asymmetry and non-spatial attention in subacute stroke patients with and without neglect. *Cortex*, 46, 602–612.

Kettunen, J.E., Nurmi, M., Dastidar, P., & Jehkonen, M. (2012). Recovery From Visual Neglect After Right Hemisphere Stroke: Does Starting Point in Cancellation Tasks Change After 6 Months? *The Clinical Neuropsychologist*, 26, 305–320.

Lezak, M.D., Howieson, D.B., Bigler, E.D., & Tranel, D. (2012). *Neuropsychological Assessment* (5th ed.). New York: Oxford University Press.

Mahoney, F.I., & Barthel, D.W. (1965). Functional evaluation: The Barthel index. *Maryland State Medical Journal*, 14, 61–65.

Manly, T. (2002). Cognitive rehabilitation for unilateral neglect: Review. *Neuropsychological Rehabilitation*, 12, 289–310.

Manly, T., Dove, A., Blows, S., George, M., Noonan, M.P., Dodds, C.M., ... Warburton, E. (2009). Assessment of unilateral spatial neglect: scoring star cancellation performance from video recordings – method, reliability, benefits, and normative data. *Neuropsychology*. 23, 519–528.

Mattingley, J.B., Bradshaw, J.L., Bradshaw, J.A., & Nettleton, N.C. (1994). Residual rightward attentional bias after apparent recovery from right hemisphere damage: Implications for a multicomponent model of neglect. *Journal of Neurology, Neurosurgery and Psychiatry*, 57, 597–604.

Mesulam, M. M. (2000). *Principles of behavioral and cognitive neurology (second edition)*. New York, NY: Oxford University Press.

Nurmi, L.J., Kettunen, J., Laihosalo, M., Ruuskanen, E.-I., Koivisto, A.-M., & Jehkonen, M. (2010). Right hemisphere infarct patients and healthy controls: Evaluation of starting points in cancellation tasks. *Journal of the International Neuropsychological Society*, 16, 902–909.

Robertson, I.H. (1993). The relationship between lateralised and non-lateralised attentional deficits in unilateral neglect. In I.H. Robertson, and J.C. Marshall (Eds.), *Unilateral Neglect: Clinical and Experimental Studies* (pp. 257-275). Hove (UK): Lawrence Erlbaum Associates.

Robertson, I.H. (2001). Do we need the “lateral” in unilateral neglect? Spatially nonselective attention deficits in unilateral neglect and their implications for rehabilitation. *NeuroImage*, 14, S85–S90.

Robertson, L.C., & Eglin, M. (1993). Attentional search in unilateral visual neglect. In I.H. Robertson & J.C. Marshall (Eds.), *Unilateral neglect: Clinical and Experimental Studies* (pp. 169–191). Hillsdale, NJ: Erlbaum.

Robertson, I. H., & Halligan, P. W., (1999). *A clinical handbook for diagnosis and treatment*. Hove,

UK: Psychology Press.

Samuelsson, H., Hjelmquist, E., Naver, H., & Blomstrand, C. (1996). Visuospatial neglect and an ipsilesional bias during the start of performance in conventional tests of neglect. *The Clinical Neuropsychologist*, 10, 15-24.

Samuelsson, H., Hjelmquist, E.K.E., Jensen, C., Ekholm, S., & Blomstrand, C. (1998). Nonlateralized attentional deficits: An important component behind persisting visuospatial neglect? *Journal of Clinical and Experimental Neuropsychology*. 20, 73–88.

Smania, N., Martini, M.C., Gambina, G., Tomelleri, G., Palamara, A., Natale, E., & Marzi, C.A. (1998). The spatial distribution of visual attention in hemineglect and extinction patients. *Brain*, 121, 1759–1770.

Suzuki E, Chen W, Kondo T. (1997). Measuring unilateral spatial neglect during stepping. *Archives of Physical Medicine and Rehabilitation*, 78, 173–178.

Taylor, D. (2003). Measuring mild visual neglect: Do complex visual tests activate rightward attentional bias? *New Zealand Journal of Physiotherapy*, 31, 67–72.

Ting, D.S.J., Pollock, A., Dutton, G.N., Doubal, F.N., Ting, D.S.W. Thompson, M., & Dhillon, B. (2011). Visual Neglect Following Stroke: Current Concepts and Future Focus. *Survey of Ophthalmology*, 56, 114–133.

Tomczak, M. & Tomczak, E. (2014). The need to report effect size estimates revisited. An overview of some recommended measures of effect size. *Trends in Sport Science*, 1, 19–25.

Viken, J. (2013). *Visuospatial neglect and processing speed: importance of lateralized and nonlateralized symptoms as predictors of functional outcome after stroke*. (Doctoral dissertation, University of Gothenburg, Sweden). Retrieved June 18, 2017 from <https://gupea.ub.gu.se/handle/2077/33882?locale=sv>

Viken, J.I., Jood, K., Jern, C., Blomstrand, C., & Samuelsson, H. (2014). Ipsilesional Bias and Processing Speed are Important Predictors of Functional Dependency in the Neglect Phenomenon after a Right Hemisphere Stroke. *The Clinical Neuropsychologist*, 28, 974–993.

Warren, M., Moore, J.M., & Vogtle, L.K. (2008). Search performance of healthy adults on cancellation tests. *American Journal of Occupational Therapy*, 62, 588–594.

Webster, J.S., Rodes, L.A., Morrill, B., Rapport, L.J., Abadee, P.S., Sowa, M.V., ... Godlewski, M. C. (1995). Rightward orienting bias, wheelchair maneuvering, and fall risk. *Archives of Physical Medicine and Rehabilitation*, 76, 924–928.

Wilson, B., Cockburn, J., & Halligan, P. (1987). *Behavioural Inattention Test*. Titchfield, UK: Thames Valley Test Company.

Table 1. Subject characteristics and comparisons of healthy control (H), neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups

Descriptive variable	H n = 40	N+ n = 18	MLI+ n = 23	N- n = 24	χ^2	Kruskal- Wallis	df	<i>p</i> -value
Male/female	16/24	9/9	13/10	16/8	4.576		3	<i>p</i> = .206
Age: Md (Q ₁ ; Q ₃)	66 (53; 74)	72 (61; 78)	69 (59; 73)	65 (57; 70)		4.484	3	<i>p</i> = .214
Education in years: Md (Q ₁ ; Q ₃)	11 (9; 14)	9 (8; 12) ²	10 (8; 12) ¹	9 (8; 12) ²		4.143	3	<i>p</i> = .246
MMSE: Md (Q ₁ ; Q ₃)	29 (28; 29)							
Thrombolysis: yes/no		12/6	16/7	18/6	.370		2	<i>p</i> = .831
Days to APE: Md (Q ₁ ; Q ₃)		5 (2; 6)	3 (2; 4)	3 (2; 5)		2.847	2	<i>p</i> = .241
Days to 6ME: Md (Q ₁ ; Q ₃)		193 (185; 197) ³	184 (179; 190) ²	186 (180; 193) ¹		5.330	2	<i>p</i> = .070
Days to 12ME: Md (Q ₁ ; Q ₃)		372 (348; 384) ⁴	376 (363; 401) ¹	369 (361; 374) ³		3.323	2	<i>p</i> = .190
Rehabilitation: yes/no		6/9 ³	2/19 ²	0/23 ¹				

Note. H = healthy control group; N- = non-neglect group; MLI+ = mild left inattention group; N+ = neglect group; Md = median; Q₁ = lower quartile; Q₃ = upper quartile; MMSE = Mini Mental State Examination; APE = acute phase examination; ME = months' examination; Rehabilitation = rehabilitation received during one-year follow-up including neuropsychological rehabilitation and occupational therapy

¹Data for 1 patient missing

²Data for 2 patients missing

³Data for 3 patients missing

⁴Data for 4 patients missing

Table 2. Neurological characteristics and comparisons of neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups

Descriptive variable	N+ n = 18	MLI+ n = 23	N- n = 24	Kruskal-Wallis	Compared pairs	Mann-Whitney U	<i>p</i> -value ¹
NIHSS at BL: Md (Q ₁ ; Q ₃)	9 (6; 13) ²	4 (3; 7)	5 (4; 6)	11.503; df=2; <i>p</i> = .003	N- vs. MLI+ N- vs. N+ MLI+ vs. N+	235.5 100.0 83.0	<i>p</i> = 1.000 <i>p</i> = .018 <i>p</i> = .006
NIHSS at AP: Md (Q ₁ ; Q ₃)	7 (3; 10) ²	1 (0; 3)	0 (0; 1)	24.441; df=2; <i>p</i> < .001	N- vs. MLI+ N- vs. N+ MLI+ vs. N+	211.5 45.5 56.5	<i>p</i> = .378 <i>p</i> < .001 <i>p</i> < .001
NIHSS at 6M: Md (Q ₁ ; Q ₃)	1 (0; 5) ⁴	0 (0; 1) ³	0 (0; 1) ²	4.659; df=2; <i>p</i> = .097			
Hemianopia present at AP	3 ²	3	0	4.093; df=2; <i>p</i> = .129			
Hemianopia present at 6M	3 ⁴	1 ³	0 ²	15.690; df=2; <i>p</i> = .058			
BI at AP: Md (Q ₁ ; Q ₃)	50 (25; 95) ²	100 (98; 100) ³	100 (99; 100) ³	22.230; df=2; <i>p</i> < .001	N- vs. MLI+ N- vs. N+ MLI+ vs. N+	226.0 50.5 61.0	<i>p</i> = 1.000 <i>p</i> < .001 <i>p</i> < .001
BI at 6M: Md (Q ₁ ; Q ₃)	100 (90; 100) ⁴	100 (100; 100) ³	100 (100; 100) ²	7.731; df=2; <i>p</i> = .021	N- vs. MLI+ N- vs. N+ MLI+ vs. N+	230.0 121.0 115.0	<i>p</i> = 1.000 <i>p</i> = .045 <i>p</i> = .114
ACA	0 (0%)	2 (9%)	0 (0%)				
MCA	13 (72%)	11 (50%)	10 (42%)				
PCA	2 (11%)	3 (14%)	2 (8%)				
ACA+MCA	1 (6%)	0 (0%)	0 (0%)				
MCA+PCA	1 (6%)	0 (0%)	1 (4%)				
ACA+MCA+PCA	1 (6%)	3 (14%)	3 (13%)				
No MRI finding ⁵	0 (0%)	3 (14%)	8 (33%)				

Note. N- = non-neglect group; MLI+ = mild left inattention group; N+ = neglect group; Md = median; Q₁ =

lower quartile; Q₃ = upper quartile; BL = baseline; NIHSS = National Institution of Health Stroke Scale;

AP = acute phase; M = months; BI = Barthel Index, ACA = infarct in the region of anterior cerebral artery;

MCA = infarct in the region of medial cerebral artery; PCA = infarct in the region of posterior cerebral

artery; MRI = magnetic resonance imaging

¹*p*-values Bonferroni corrected multiplying by three

²Data for 1 patient missing

³Data for 2 patients missing

⁴Data for 3 patients missing

⁵All the patients without a finding in the MRI have received thrombolytic treatment at the baseline prior to the MRI examination

Table 3. Median (Md) and quartiles (Q₁; Q₃) of the Behavioural Inattention Test (BIT) sum scores in healthy control (H), neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups and between-group comparisons

Variable	Group	n (missing)	Md (Q ₁ ; Q ₃)	Kruskal- Wallis	Compared pairs	Mann- Whitney U	<i>p</i> -value ¹	Effect size
BIT sum score at AP	H	40	142 (139; 144)	36.189; df=3; <i>p</i> < .001 $\eta^2 = .34$	H vs. N-	406.0	1.000	<i>r</i> = -.13
	N-	24	143 (141; 145)		H vs. MLI+	344.5	.948	<i>r</i> = -.18
	MLI+	22 (1)	144 (141; 145)		H vs. N+	39.0	< .001	<i>r</i> = -.70
	N+	17 (1)	128 (113; 136)		N- vs. MLI+	249.5	1.000	<i>r</i> = -.05
					N- vs. N+	18.0	< .001	<i>r</i> = -.77
					MLI+ vs. N+	15.5	< .001	<i>r</i> = -.78
BIT sum score at 6M	H	40	142 (139; 144)	5.822; df=3; <i>p</i> = .121 $\eta^2 = .03$				
	N-	23 (1)	144 (142; 145)					
	MLI+	20 (3)	143 (140; 145)					
	N+	14 (4)	140 (137; 143)					
BIT sum score at 12M	H	40	142 (139; 144)	10.798; df=3; <i>p</i> = .013 $\eta^2 = .09$	H vs. N-	285.0	1.000	<i>r</i> = -.13
	N-	17 (7)	143 (141; 145)		H vs. MLI+	346.5	1.000	<i>r</i> = -.14
	MLI+	21 (2)	143 (140; 145)		H vs. N+	154.0	.072	<i>r</i> = -.34
	N+	14 (4)	139 (135; 142)		N- vs. MLI+	178.0	1.000	<i>r</i> = -.00
					N- vs. N+	50.5	.036	<i>r</i> = -.49
			MLI+ vs. N+	63.0	.024	<i>r</i> = -.48		

Note. BIT = the Behavioural Inattention Test; H = healthy control group; N- = non-neglect group; MLI+ = mild left inattention group; N+ = neglect group; Md = median; Q₁ = lower quartile; Q₃ = upper quartile;

AP = acute phase; M = months

¹*p*-values Bonferroni corrected multiplying by six

Table 4. Median (Md) and quartiles (Q₁; Q₃) of the Center of Cancellation (CoC) sum scores in healthy control (H), neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups and between-group comparisons

Variable	Group	n (missing)	Md (Q ₁ ; Q ₃)	Kruskal- Wallis	Compared pairs	Mann- Whitney U	p-value ¹	Effect size
CoC sum score at AP	H	40	0,0 (-1,9; 0,1)	9.695; df=3; p = .022 $\eta^2 = .06$	H vs. N-	456.5	1.000	<i>r</i> = -.04
	N-	24	0,0 (-0,2; 0,1)		H vs. MLI+	438.0	1.000	<i>r</i> = -.04
	MLI+	23	0,0 (-0,2; 0,1)		H vs. N+	191.0	.024	<i>r</i> = -.37
	N+	18	0,8 (-0,1; 1,6)		N- vs. MLI+	272.5	1.000	<i>r</i> = -.01
					N- vs. N+	116.0	.066	<i>r</i> = -.39
					MLI+ vs. N+	113.0	.084	<i>r</i> = -.39
CoC sum score at 6M	H	40	0,0 (-1,9; 0,1)	3.980; df=3; <i>p</i> = .264 $\eta^2 = .01$				
	N-	23 (1)	0,0 (-0,2; 0,1)					
	MLI+	20 (3)	-0,1 (-0,2; 0,0)					
	N+	14 (4)	0,0 (-0,2; 0,2)					
CoC sum score at 12M	H	40	0,0 (-1,9; 0,1)	2.582; df=3; <i>p</i> = .461 $\eta^2 = .00$				
	N-	19 (5)	0,0 (-0,4; 0,0)					
	MLI+	22 (1)	0,0 (-0,2; 0,1)					
	N+	14 (4)	-0,2 (-0,5; 0,1)					

Note. CoC = the Center of Cancellation; H = healthy control group; N- = non-neglect group; MLI+ = mild left inattention group; N+ = neglect group; Md = median; Q₁ = lower quartile; Q₃ = upper quartile; AP = acute phase; M = months

¹*p*-values Bonferroni corrected multiplying by six

Table 5. Median (Md) and quartiles (Q₁; Q₃) of starting points (SPs) for line and star cancellation tasks in healthy control (H), neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups and between-group comparisons

Variable	Group	n (missing)	Md (Q ₁ ; Q ₃)	Kruskal- Wallis	Compared pairs	Mann- Whitney U	<i>p</i> -value ¹	Effect size
SP for line cancellation at AP	H	40	-11,7 (-11,7; -11,7)	71.329; df=3; <i>p</i> < .001 η ² = .68	H vs. N-	416.0	.750	<i>r</i> = -.19
	N-	24	-11,7 (-11,7; -11,7)		H vs. MLI+	26.0	< .001	<i>r</i> = -.84
	MLI+	23	10,8 (3,5; 12,0)		H vs. N+	93.0	< .001	<i>r</i> = -.68
	N+	18	3,4 (-4,6; 11,7)		N- vs. MLI+	4.0	< .001	<i>r</i> = -.90
					N- vs. N+	40.5	< .001	<i>r</i> = -.79
					MLI+ vs. N+	160.0	1.000	<i>r</i> = -.19
SP for star cancellation at AP	H	40	-11,9 (-11,9; -11,9)	60.236; df=3; <i>p</i> < .001 η ² = .57	H vs. N-	387.0	.768	<i>r</i> = -.19
	N-	23 (1)	-11,9 (-11,9; -7,8)		H vs. MLI+	34.0	< .001	<i>r</i> = -.84
	MLI+	23	1,8 (0,8; 12,0)		H vs. N+	107.0	< .001	<i>r</i> = -.67
	N+	18	4,5 (-11,2; 12,3)		N- vs. MLI+	17.0	< .001	<i>r</i> = -.82
					N- vs. N+	71.0	< .001	<i>r</i> = -.59
					MLI+ vs. N+	192.5	1.000	<i>r</i> = -.06
SP for line cancellation at 6M	H	40	-11,7 (-11,7; -11,7)	26.020; df=3; <i>p</i> < .001 η ² = .24	H vs. N-	456.5	1.000	<i>r</i> = -.01
	N-	23 (1)	-11,7 (-11,7; -11,7)		H vs. MLI+	175.5	< .001	<i>r</i> = -.56
	MLI+	21 (2)	3,5 (-11,7; 11,6)		H vs. N+	154.5	.012	<i>r</i> = -.43
	N+	14 (4)	-4,4 (-11,7; 5,4)		N- vs. MLI+	109.5	< .001	<i>r</i> = -.53
					N- vs. N+	93.5	.066	<i>r</i> = -.42
					MLI+ vs. N+	119.0	1.000	<i>r</i> = -.16
SP for star cancellation at 6M	H	40	-11,9 (-11,9; -11,9)	21.423; df=3; <i>p</i> < .001 η ² = .20	H vs. N-	418.0	1.000	<i>r</i> = -.12
	N-	23 (1)	-11,9 (-11,9; -11,9)		H vs. MLI+	187.5	< .001	<i>r</i> = -.53
	MLI+	20 (3)	-5,2 (-11,9; 2,2)		H vs. N+	156.0	.006	<i>r</i> = -.44
	N+	14 (4)	-9,7 (-11,9; 3,4)		N- vs. MLI+	109.5	.030	<i>r</i> = -.42
					N- vs. N+	102.5	.192	<i>r</i> = -.35
					MLI+ vs. N+	127.5	1.000	<i>r</i> = -.08
SP for line cancellation at 12M	H	40	-11,7 (-11,7; -11,7)	9.716; df=3; <i>p</i> = .021 η ² = .07	H vs. N-	357.5	1.000	<i>r</i> = -.07
	N-	19 (5)	-11,7 (-11,7; -11,7)		H vs. MLI+	291.5	.030	<i>r</i> = -.35
	MLI+	22 (1)	-11,7 (-11,7; 8,5)		H vs. N+	196.0	.162	<i>r</i> = -.30
	N+	14 (4)	-11,7 (-11,7; 8,3)		N- vs. MLI+	153.5	.516	<i>r</i> = -.27
					N- vs. N+	102.5	1.000	<i>r</i> = -.24
					MLI+ vs. N+	150.5	1.000	<i>r</i> = -.02
SP for star cancellation at 12M	H	40	-11,9 (-11,9; -11,9)	11.422; df=3; <i>p</i> = .010 η ² = .09	H vs. N-	362.0	1.000	<i>r</i> = -.06
	N-	19 (5)	-11,9 (-11,9; -11,9)		H vs. MLI+	289.0	.018	<i>r</i> = -.38
	MLI+	22 (1)	-11,9 (-11,9; 6,8)		H vs. N+	188.0	.060	<i>r</i> = -.35
	N+	14 (4)	-11,9 (-11,9; 7,6)		N- vs. MLI+	153.0	.456	<i>r</i> = -.28
					N- vs. N+	99.0	.684	<i>r</i> = -.27
					MLI+ vs. N+	152.0	1.000	<i>r</i> = -.01

Note. SP = starting point; H = healthy control group; N- = non-neglect group; MLI+ = mild left

inattention group; N+ = neglect group; Md = median; Q₁ = lower quartile; Q₃ = upper quartile; AP =

acute phase; M = months

¹*p*-values Bonferroni corrected multiplying by six

Table 6. Median (Md) and quartiles (Q₁; Q₃) of cancellation time for letter and star cancellation tasks in healthy control (H), neglect (N+), mild left inattention (MLI+) and non-neglect (N-) groups and between-group comparisons

Variable	Group	n (missing)	Md (Q ₁ ; Q ₃)	Kruskal-Wallis	Compared pairs	Mann-Whitney U	p-value ¹	Effect size
Cancellation time for letter cancellation at AP	H	40	65 (57; 96)	22.063; df=3; p < .001 $\eta^2 = .19$	H vs. N-	306.5	.168	<i>r</i> = -.28
	N-	23 (1)	81 (69; 105)		H vs. MLI+	218.0	.006	<i>r</i> = -.44
	MLI+	23	94 (77; 127)		H vs. N+	141.0	< .001	<i>r</i> = -.48
	N+	18	113 (83; 181)		N- vs. MLI+	178.5	.354	<i>r</i> = -.28
					N- vs. N+	114.0	.084	<i>r</i> = -.38
Cancellation time for star cancellation at AP	H	40	43 (33; 57)	42.030; df=3; p < .001 $\eta^2 = .39$	MLI+ vs. N+	159.5	1.000	<i>r</i> = -.19
	N-	24	68 (47; 105)		H vs. N-	206.0	< .001	<i>r</i> = -.48
	MLI+	23	86 (68; 105)		H vs. MLI+	78.5	< .001	<i>r</i> = -.69
	N+	18	90 (60; 125)		H vs. N+	76.5	< .001	<i>r</i> = -.63
					N- vs. MLI+	192.5	.456	<i>r</i> = -.26
N- vs. N+	147.0	.474	<i>r</i> = -.27					
Cancellation time for letter cancellation at 6M	H	40	65 (57; 96)	14.286; df=3; p = .003 $\eta^2 = .12$	MLI+ vs. N+	189.0	1.000	<i>r</i> = -.07
	N-	21 (3)	81 (61; 107)		H vs. N-	336.0	1.000	<i>r</i> = -.16
	MLI+	21 (2)	97 (84; 121)		H vs. MLI+	222.5	.018	<i>r</i> = -.38
	N+	14 (4)	90 (77; 144)		H vs. N+	135.0	.024	<i>r</i> = -.39
					N- vs. MLI+	139.5	.246	<i>r</i> = -.31
N- vs. N+	96.5	.534	<i>r</i> = -.29					
Cancellation time for star cancellation at 6M	H	40	43 (33; 57)	20.028; df=3; p < .001 $\eta^2 = .18$	MLI+ vs. N+	145.5	1.000	<i>r</i> = -.01
	N-	23 (1)	48 (43; 63)		H vs. N-	295.5	.114	<i>r</i> = -.30
	MLI+	20 (3)	68 (53; 102)		H vs. MLI+	153.5	< .001	<i>r</i> = -.50
	N+	14 (4)	61 (48; 83)		H vs. N+	139.0	.030	<i>r</i> = -.38
					N- vs. MLI+	137.0	.138	<i>r</i> = -.35
N- vs. N+	124.0	1.000	<i>r</i> = -.19					
Cancellation time for letter cancellation at 12M	H	40	65 (57; 96)	10.371; df=3; p = .016 $\eta^2 = .08$	MLI+ vs. N+	124.5	1.000	<i>r</i> = -.09
	N-	19 (5)	65 (58; 88)		H vs. N-	374.5	1.000	<i>r</i> = -.01
	MLI+	22 (1)	79 (69; 91)		H vs. MLI+	348.0	1.000	<i>r</i> = -.17
	N+	14 (4)	106 (72; 147)		H vs. N+	146.0	.048	<i>r</i> = -.36
					N- vs. MLI+	151.5	.798	<i>r</i> = -.23
N- vs. N+	58.5	.042	<i>r</i> = -.47					
Cancellation time for star cancellation at 12M	H	40	43 (33; 57)	17.440; df=3; p = .001 $\eta^2 = .16$	MLI+ vs. N+	89.0	.210	<i>r</i> = -.35
	N-	19 (5)	43 (38; 58)		H vs. N-	321.5	1.000	<i>r</i> = -.12
	MLI+	21 (2)	58 (47; 68)		H vs. MLI+	221.5	.018	<i>r</i> = -.39
	N+	14 (4)	78 (41; 120)		H vs. N+	117.0	.006	<i>r</i> = -.44
					N- vs. MLI+	120.0	.186	<i>r</i> = -.34
N- vs. N+	66.0	.090	<i>r</i> = -.43					
				MLI+ vs. N+	93.0	.414	<i>r</i> = -.31	

Note. H = healthy control group; N- = non-neglect group; MLI+ = mild left inattention group; N+ =

neglect group; Md = median; Q₁ = lower quartile; Q₃ = upper quartile; AP = acute phase; M = months

¹p-values Bonferroni corrected multiplying by six

Table 7. ROC analyses of processing speed for healthy subjects vs. N+ patients in letter and star cancellation tasks

Task	Sensitivity (%)	Specificity (%)	AUC	95% CI for AUC	Cutoff (sec)
Letter cancellation	83	75	0.804	0.682–0.927	82.5
Star cancellation	83	83	0.894	0.795–0.992	59.5

Note. ROC = receiver operating characteristic; H = healthy; N+ = neglect; AUC = area under the curve; CI = confidence interval; Cutoff = guideline cutoff for slowed processing speed (cancellation time slower than cutoff indicates slowed processing speed)




Figure 1a. Receiver operating characteristic curve: processing speed for healthy subjects versus visual neglect (N+) patients in letter cancellation task.




Figure 1b. Receiver operating characteristic curve: processing speed for healthy subjects versus visual neglect (N+) patients in star cancellation task.

Appendix 1. Methods used in neuropsychological examination

Assessed domain	Method	Description
Visual neglect	Behavioural Inattention Test (BIT)	<p>Visual neglect was assessed with the six subtests of the BITC (Jehkonen, 2002a; Wilson et al., 1987). For individual patients, the criterion for visual neglect and inclusion in the N+ group was failure in at least two subtests (possible range 0-6), with cutoff points based on Wilson et al. (1987). The BIT sum score was used as a variable for statistical analyses in group comparisons (possible range 0-146). During assessment, each task sheet was placed in front of the subject in line with the midsagittal plane, an arrow indicating the location of the midline on the A4-sized sheet. Moving the sheet was not allowed.</p> <p><i>The line cancellation task</i> includes 40 lines randomly spaced on a landscape sheet. Eighteen target lines locate on both sides of the sheet. Four central lines are used as examples by the examiner and are not scored (maximum score 36, cutoff 34).</p> <p><i>The letter cancellation task</i> includes 40 target letters (E, R) and 130 non-target letters in five rows on a landscape sheet (maximum score 40, cutoff 32). Two example letters (E & R) beneath the rows are not scored.</p> <p><i>The star cancellation task</i> includes 56 targets (small stars) and 75 non-targets (big stars, letters and words) randomly spaced on a landscape sheet. Twenty-seven targets locate on both sides of the sheet. Two central stars are examples and not scored (maximum score 54, cutoff 51). In these tasks, the subjects were instructed to search all the targets.</p> <p><i>The figure and shape copying</i> contains two tasks. Figure copying includes three figures – a star, a cube, and a daisy – located on the left side of a portrait sheet, and the subjects were instructed to copy the figures to the boxes on the right side. Shape copying contains three geometric shapes on a landscape sheet, which the subjects copy to a separate blank sheet. The scoring was based on the completeness of the drawings (maximum score 4, cutoff 3).</p> <p><i>The line bisection</i> includes three horizontal 20.4 cm lines in a staircase manner across a landscape sheet. The subjects were instructed to estimate the center for each line. The scoring was based on the level of the deviation between the estimated and the true center of the lines (maximum score 9, cutoff 7).</p> <p><i>The representational drawing</i> contains three blank landscape sheets, to which the subjects were instructed to draw a clock face with numbers, a human being, and a butterfly. The scoring was based on the completeness of the drawings (maximum score 3, cutoff 2).</p>
Laterality of neglect	Center of Cancellation (CoC)	<p>Laterality of visual neglect was evaluated based on the CoC (Binder, Marshall, Lazar, Benjamin, & Mohr, 1992) in the BIT cancellation tasks. CoC refers to the mean location of the targets found, i.e. the mean of the targets' distances from the midline, with left-sided distances expressed as negative values and right-sided as positive values. CoC value 0 indicates a symmetrical performance, < 0 refers to a performance lateral to the left side (more targets found on the left side), and > 0</p>

indicates a performance lateral towards the right side (more targets found on the right side). The CoC sum variable was formed by adding up the CoC values of the three BIT cancellation tasks (possible range: -37.6 - +39.4).

Initial ipsilateral orienting bias	Starting point (SP) in cancellation tasks	Initial ipsilateral orienting bias was assessed based on the SPs of the three BIT cancellation tasks. SP indicates a location, i.e. a specific stimulus (target or non-target) from which the subject starts the cancellation task. In each task, SP was expressed as distance from the sheet's midline in centimeters, with negative values for left-sided SPs and positive values for right-sided SPs (possible ranges: line cancellation -13.4 – +13.2 cm; letter cancellation -12.4 – +12.2 cm; star cancellation -11.9 – +14.0 cm). For individual patients, the criterion for ipsilateral orienting bias and inclusion in the MLI+ group (when actual neglect was not present) was an atypical SP in at least two cancellation tasks, meaning that these SPs were outside the guideline value for a normal SP. The guideline values for a normal SP were based on Nurmi et al. (2010; line cancellation: normal SP < -5.8 cm ≤ atypical SP, letter cancellation: normal SP < -10.0 cm ≤ atypical SP, star cancellation: normal SP < -11 cm ≤ atypical SP). In addition, because the SPs particularly in the line and star cancellation tasks have been found to be sensitive in detecting ipsilateral orienting bias (Nurmi et al., 2010; Jalas et al., 2002), they were chosen as SP variables for statistical analyses in group comparisons.
Processing speed	Cancellation time	Processing speed in visual search was assessed by measuring cancellation time in seconds to determine the duration of performance in cancellation tasks. Since complex tasks are more sensitive in detecting deficits in processing speed than simpler ones (Benton, 1986; Lezak, Howieson, & Tranel, 2012), processing speed was measured with the letter and star cancellation tasks, which place greater demands on attention than line cancellation (Mesulam, 2000). Cancellation time was measured for both tasks separately. Subjects were instructed to start the cancellation from the examiner's signal, operate fast and accurately, and to tell the examiner when they had completed the task. The examiner measured the duration from the start signal to completion.
Cognitive status of healthy controls	Mini Mental State Examination (MMSE)	The healthy controls' cognitive status was evaluated with the MMSE (Folstein, Folstein, & McHugh, 1975; range 0–30; 0 = severe cognitive impairment, 30 = normal cognitive functioning) using the cutoff score of ≤ 24 for exclusion from the study.

Appendix 2. Methods used in neurological and neuroradiological examinations

Assessed domain	Method	Description
Stroke severity and hemianopia	National Institute of Health Stroke Scale (NIHSS)	Stroke severity and hemianopia were assessed using the NIHSS (Goldstein, Bertels, & Davis, 1989). Stroke severity was based on the NIHSS sum score (range 0–34; 0 = no defect, 34 = severe stroke). Visual field deficit was evaluated on a scale from 0 to 2 (0 = no vision loss, 1 = partial hemianopia, 2 = complete hemianopia).
Functioning in basic activities of daily living (ADL)	Barthel Index (BI)	The BI was used for evaluating functioning in basic ADL (range 0–100; 0 = dependent, 100 = independent; Mahoney & Barthel, 1965).
Infarct diagnosis and exclusion of hemorrhage	Computerized tomography (CT)	CT was conducted as a standard method of treatment on arrival at hospital in order to corroborate the infarct diagnosis and to exclude any hemorrhage.
Infarct location and exclusion of significant cerebral atrophy	Magnetic Resonance Imaging (MRI)	MRI examination was used to rule out a presence of significant cerebral atrophy and to identify the location of infarct in anterior cerebral artery (ACA), medial cerebral artery (MCA) and posterior cerebral artery (PCA).

Appendix 3. Individual values of neuropsychological variables for each patient in N+, MLI+ and N- groups

Patient	Acute phase								6 months								12 months								Rehabilitation yes/no	
	BIT sum score	Failed sub- tests	CoC sum score (cm)	SP in LiC (cm)	SP in StC (cm)	Aty- pical SP:s	LeC time (sec)	StC time (sec)	BIT sum score	Failed sub- tests	CoC sum score (cm)	SP in LiC (cm)	SP in StC (cm)	Aty- pical SP:s	LeC time (sec)	StC time (sec)	BIT sum score	Failed sub- tests	CoC sum score (cm)	SP in LiC (cm)	SP in StC (cm)	Aty- pical SP:s	LeC time (sec)	StC time (sec)		
N+																										
1	140	2	-0.8	-11.7	2.3	1	163	60	143	0	-0.3	11.1	12.8	2	129	52	-	-	-	-	-	-	-	-	-	no
2	107	4	1.2	-7.4	6.7	1	62	57	140	1	-0.4	-11.7	-11.9	0	78	41	142	1	-0.1	-11.7	-11.9	0	112	40	no	
3	133	2	-0.7	3.5	12.4	3	83	116	146	0	0.0	3.5	-11.0	2	128	77	141	0	-0.4	11.5	-11.9	1	75	60	no	
4	123	3	1.8	3.5	6.7	3	209	77	138	2	-0.6	3.5	-11.9	1	81	53	134	3	-1.0	11.7	9.7	2	103	82	no	
5	120 ¹	3 ¹	0.7	12.2	1.7	3	179	83	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	yes	
6	138	2	-0.3	-3.6	-0.1	2	92	129	137	1	0.8	-8.7	1.7	2	84	50	142	0	-0.1	-11.7	-11.0	1	53	41	no	
7	132	3	-0.4	-3.6	-1.9	2	128	107	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8	136	2	0.0	0.0	-11.9	1	54	31	142	0	0.2	-11.7	-11.9	0	57	31	135	2	-0.2	-11.7	-11.9	0	63	35	no	
9	106	6	2.6	11.5	-11.9	1	115	96	131	2	0.5	12.2	-11.9	1	80	62	136	1	-0.8	-11.7	-11.9	0	105	70	yes	
10	139	2	0.3	-11.7	-11.0	1	98	58	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
11	128	5	1.2	12.2	-11.9	2	83	70	144	1	0.0	-11.7	-6.6	1	187	85	141	1	-0.4	-11.7	-11.9	0	148	103	yes	
12	136	2	0.0	0.0	13.9	3	96	60	136	2	0.0	0.0	13.9	3	96	60	137	2	-0.6	3.4	12.5	2	79	74	no	
13	127	3	1.0	12.3	12.9	2	148	124	146	0	0.0	-11.7	-11.9	1	114	82	143	0	0.0	-11.7	-11.9	0	137	117	no	
14	116	3	1.5	12.1	14.0	3	302	163	136	1	0.0	12.4	-6.6	3	205	164	144	0	-0.2	-7.3	1.7	1	347	211	yes	
15	109	4	6.8	7.2	8.1	3	245	139	-	-	-	-	-	-	-	-	129	5	0.4	7.2	6.9	3	151	149	-	
16	38	6	28.9	10.9	12.3	3	111	98	137	1	1.7	-11.7	-8.5	1	72	78	139	0	-0.2	-11.7	-11.9	0	106	116	yes	
17	136	2	0.2	7.0	9.6	3	188	210	140	0	-0.1	3.3	8.6	2	222	162	135	2	0.4	12.2	12.5	3	147	128	no	
18	127	3	1.1	-11.7	-11.9	0	70	73	143	0	0.2	-11.7	-11.9	0	62	38	138	1	0.7	-11.7	-11.9	0	58	34	yes	
MLI+																										
1	145	0	-0.1	3.5	12.4	2	72	94	144	0	-0.2	11.5	-11.9	1	51	50	144	0	-0.1	3.5	-11.9	1	71	57	no	
2	145	0	-0.1	3.5	6.9	2	127	110	143	0	0.0	3.5	1.7	2	129	108	144	0	0.0	3.5	6.8	2	110	100	no	
3	145	0	0.0	-3.6	-1.9	2	215	103	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
4	143	0	-0.1	12.2	12.7	2	114	105	141	1	-0.2	11.5	-1.9	2	128	88	143	0	-0.1	11.5	-11.9	1	105	79	yes	
5	142	0	-0.4	3.5	1.7	2	92	86	146	0	0.0	12.2	-11.9	1	98	77	144	0	-0.3	-11.7	-11.9	0	75	57	no	
6	141	0	0.4	12.2	13.2	3	162	139	142	0	-0.1	3.5	8.8	2	198	107	-	-	-0.3	-11.7	-6.7	1	77	69	yes	
7	146	0	0.0	-3.9	2.3	2	143	135	-	0	-	-11.7	-	-	95	-	145	0	0.0	-11.7	-11.1	0	90	73	no	
8	146	0	0.0	0.0	1.7	2	91	67	141	0	-0.3	-11.7	-11.9	0	70	44	146	0	0.0	-11.7	-11.9	0	81	58	no	
9	139 ¹	1	0.1	12.2	12.4	2	143	89	139	0	0.0	12.2	12.4	3	90	52	143	1	0.1	12.2	8.2	2	87	66	no	
10	143	0	0.2	12.2	6.8	3	80	82	140	1	-0.2	12.3	-11.2	1	90	80	140	0	-0.1	11.4	12.4	2	82	65	no	
11	136	1	0.7	11.5	-3.7	2	94	112	137	1	-0.2	-11.7	-11.9	0	78	55	141	0	-0.1	-11.7	-11.9	0	51	-	no	
12	145	0	-0.1	3.5	1.7	2	49	49	145	0	0.2	-11.7	-11.1	0	46	33	143	0	0.2	3.3	-11.9	1	43	31	no	
13	141	0	-0.2	12.0	1.2	2	141	79	143	0	-0.3	4.0	2.3	2	144	70	140	0	0.3	12.0	6.8	3	107	62	no	

14	146	0	0.0	10.8	12.0	2	77	51	146	0	0.0	0.0	-11.9	1	114	54	146	0	0.0	-11.7	-11.9	0	96	37	no
15	145	0	-0.2	11.2	11.6	2	117	76	-	-	-	-	-	-	-	-	144	0	-0.4	-11.7	-6.7	1	87	44	-
16	144	0	0.2	12.0	1.5	2	74	83	140	0	0.2	12.0	13.5	2	110	71	146	0	0.0	-11.7	-11.9	0	85	63	no
17	132	1	-1.2	11.3	13.0	3	90	86	142	0	-0.3	11.7	-2.8	2	110	108	140	0	-0.2	0.0	12.5	2	95	94	no
18	140	0	0.0	3.5	-2.0	2	68	59	143	0	-0.1	-11.7	-11.9	0	95	55	143	0	0.2	-11.7	-11.9	0	70	58	no
19	137	0	-0.8	-4.0	0.7	2	91	78	145	0	0.3	4.1	-3.7	2	97	66	137	3	-0.2	7.5	-1.8	2	70	66	no
20	142	0	0.1	-3.5	-6.4	2	106	68	139	1	-0.4	-11.7	-6.6	1	156	139	134	2	-0.4	-11.7	-11.9	0	74	46	no
21	142	0	0.1	3.5	6.2	3	114	95	141	0	-0.4	0.0	12.6	2	114	167	135	1	1.0	-11.7	-11.9	0	66	49	no
22	145	0	-0.2	11.8	0.8	2	64	53	145	0	-0.2	-11.7	-11.9	0	66	45	145	0	-0.1	-11.7	-11.9	0	57	32	no
23	144	0	-0.1	11.5	1.8	2	118	115	144	0	-0.1	4.0	0.8	2	95	54	145	0	0.2	11.5	12.5	3	64	48	no
N-																									
1	144	0	-0.2	-11.7	-7.8	1	85	48	136	1	-0.4	-11.7	1.7	1	93	40	140	0	-0.5	-11.7	13.6	1	84	35	no
2	143	0	0.2	-11.7	-7.8	1	105	116	142	1	-0.5	-11.7	-11.9	0	122	85	144	0	0.0	-11.7	-11.9	0	95	59	no
3	146	0	0.0	-11.7	-11.9	0	156	136	145	0	0.2	-11.7	-11.9	0	88	54	142	0	-0.4	-11.7	-11.9	0	57	45	no
4	143	1	-0.1	-11.7	-11.9	0	107	66	145	0	0.0	-11.7	-11.9	0	81	57	-	-	-	-	-	-	-	-	no
5	142	0	-0.4	-0.1	-11.9	1	100	75	145	0	0.2	12.2	-11.9	1	128	85	146	0	0.0	-11.7	-11.9	0	109	96	no
6	144	0	0.2	-11.7	-11.9	0	70	46	143	0	-0.4	-11.7	-11.9	0	69	37	141	0	-0.4	-11.7	-11.9	0	65	37	no
7	139	0	0.3	-11.7	-7.8	1	-	121	132	2	-0.6	-7.4	0.7	1	103	63	-	-	-	-	-	-	-	-	no
8	146	0	0.0	-11.7	1.5	1	69	80	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9	143	0	-0.2	-11.7	-	²	66	58	143	1	0.0	-11.7	-11.9	0	61	43	-	-	-	-	-	-	-	-	no
10	138	1	0.1	-11.7	-11.9	0	73	44	145	0	-0.1	-11.7	-11.9	0	-	43	143	0	-0.1	-11.7	-11.9	0	68	44	no
11	132	1	-1.6	-11.7	-11.9	0	105	187	134	1	0.9	-11.7	-11.9	0	54	46	133 ¹	1 ¹	0.1	-3.6	-11.9	1	67	40	no
12	146	0	0.0	-11.7	-11.9	0	119	89	146	0	0.0	-11.7	-11.9	0	67	48	144	0	0.1	-11.7	-11.9	0	78	42	no
13	145	0	0.0	-11.7	-11.9	0	139	110	146	0	0.0	-11.7	-11.9	0	118	70	-	-	-	-	-	-	-	-	no
14	142	1	-0.5	-11.7	-11.9	0	62	44	145	1	0.0	-11.7	-11.9	0	59	42	146	0	0.0	-11.7	-11.9	0	60	40	no
15	143	0	0.5	-11.7	-7.3	1	121	118	141	0	0.1	-11.7	-11.9	0	110	80	132 ¹	1 ¹	0.3	-11.7	-11.9	0	88	69	no
16	145	0	-0.1	-11.7	-11.9	0	61	28	144	0	-0.2	-11.7	-11.9	0	63	31	144	0	0.0	-11.7	-11.9	0	58	36	no
17	142	0	-0.1	-11.7	-11.9	0	81	49	142	0	-0.2	3.5	1.5	2	84	48	146	0	0.0	3.5	-11.9	1	91	48	no
18	140	1	0.1	-11.7	-11.9	0	74	69	145	0	0.2	-11.7	-6.8	1	68	63	138	1	0.4	3.5	2.5	2	61	58	no
19	144	0	0.1	-11.7	-11.9	0	62	48	144	0	0.1	-11.7	-11.9	0	61	45	139	0	-0.3	-11.7	-11.9	0	50	38	no
20	146	0	0.0	-11.7	-2.3	1	76	61	146	0	0.0	-11.7	-11.9	0	82	62	144	0	-0.2	-11.7	-11.9	0	61	52	no
21	137	1	-0.5	-11.7	1.5	1	69	40	145	0	0.0	-11.7	-11.9	0	52	39	141	0	-0.9	-11.7	-11.9	0	55	35	no
22	144	0	-0.1	-11.7	-11.9	0	82	72	142	0	-0.1	-7.1	-11.9	0	-	44	142	1	-0.2	-11.7	-11.9	0	65	43	no
23	143	0	0.3	-11.7	-11.9	0	87	91	144	0	0.1	-11.7	-11.9	0	116	106	146	0	0.0	-11.7	-11.9	0	102	67	no
24	137	1	-0.4	-11.7	-11.9	0	62	42	140	0	0.8	-11.7	1.6	1	60	58	135	2	-0.6	12.2	8.9	2	51	39	no

Note. BIT = the Behavioural Inattention Test; CoC = the Center of Cancellation; SP = starting point; LiC = the line cancellation task; StC = the star cancellation task; LeC = the letter cancellation task; sec = seconds; N- = non-neglect group; MLI+ = mild left inattention group; N+ = neglect group

¹ Exact value is unknown because of missing data, the given value is a minimum value

² Exact value is unknown because of missing data, the value may vary from 0 to 1