

RESEARCH

Open Access



# QTc prolongation is associated with severe desaturations in stroke patients with sleep apnea

Saara Sillanmäki<sup>1\*</sup>, Jukka A. Lipponen<sup>2</sup>, Henri Korkalainen<sup>1,2</sup>, Antti Kulkas<sup>3</sup>, Timo Leppänen<sup>1,2,5</sup>, Sami Nikkonen<sup>1,2</sup>, Juha Töyräs<sup>2,4,5</sup>, Brett Duce<sup>6,7</sup>, Aaron Suni<sup>8</sup> and Samu Kainulainen<sup>1,2</sup>

## Abstract

**Background:** Obstructive sleep apnea (OSA) is associated with vascular diseases from which stroke and sudden cardiac death are the most significant ones. It is known that disturbances of the autonomic nervous system and electrocardiographic changes are seen in patients with a previous cerebrovascular event. However, the pathophysiological cascade between breathing cessations, autonomic regulation, and cardiovascular events is not fully understood.

**Methods:** We aimed to investigate the acute effect of desaturation on repolarisation in OSA patients with a previous stroke. We retrospectively analysed heart-rate corrected QT (QTc) intervals before, within, and after 975 desaturations in OSA patients with a stroke history and at least moderate sleep apnea (apnea–hypopnea index  $\geq 15$  events/h,  $n = 18$ ). For the control population ( $n = 18$ ), QTc intervals related to 1070 desaturation were analysed. Desaturations were assigned to groups according to their length and duration. Groupwise comparisons and regression analyses were further executed to investigate the influence of desaturation features on repolarization.

**Results:** In the stroke population the QTc prolonged at least 11 ms during 27.1% of desaturations, and over 20 ms during 12.2% of desaturations. QTc was significantly prolonged during longer ( $> 30$  s,  $p < 0.04$ ) and deeper ( $> 7\%$ ,  $p < 0.03$ ) desaturations. Less severe desaturations didn't influence QTc. In median, QTc prolonged 7.5 ms during  $> 45$  s desaturations and 7.4 ms during  $> 9\%$  deep desaturations. In the control population, QTc prolongation was observed but to a significantly lesser extent than in stroke patients. In addition, desaturation duration was found to be an independent predictor of QTc prolongation ( $\beta = 0.08$ ,  $p < 0.001$ ) among all study patients.

**Conclusions:** We demonstrated that longer ( $> 30$  s) and deeper ( $> 7\%$ ) desaturations prolong QTc in patients with stroke history. A significant proportion of desaturations produced clinically relevant QTc prolongation. As it is known that a long QTc interval is associated with lethal arrhythmias, this finding might in part explain the pathophysiological sequelae of cardiovascular mortality in OSA patients with a history of stroke.

**Keywords:** QTc, Stroke, Obstructive sleep apnea, Desaturation, Repolarisation

## Introduction

Cardiovascular diseases are one of the leading causes of morbidity and mortality. Obstructive sleep apnea (OSA), a sleep disorder affecting approximately one billion adults worldwide [1], shares several common risk factors and comorbid conditions with cardiovascular disease. Furthermore, OSA is associated with an increased

\*Correspondence: saara.sillanmaki@kuh.fi

<sup>1</sup> Diagnostic Imaging Center, Kuopio University Hospital (KYS), P.O. Box 100, 70029 Kuopio, Finland

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

risk of cerebrovascular stroke [2]. It has been shown that 90% of patients with a previous stroke have at least mild OSA (apnea–hypopnea index (AHI)  $\geq 5$  events/h) [3]. This connection between stroke, OSA, and cardiac arrhythmogenicity is clinically relevant since OSA is a known risk factor for sudden cardiac death (SCD) [4].

Cerebrovascular stroke can cause changes in autonomic tone, electrocardiographic abnormalities, and increase the incidence of lethal cardiac arrhythmias [5]. Furthermore, it has been shown that stroke survivors with long QT intervals are at a higher risk of cardiac death [6]. In most cases, sudden, unexpected death is caused by fatal cardiac arrhythmias [7] while QT prolongation is a risk factor for the development of ventricular tachycardia and even SCD [8]. Although extensive research has been carried out on electrocardiogram (ECG) changes and arrhythmia in OSA patients, there are no studies adequately investigating QT interval prolongation relative to desaturation features in patients with known stroke history.

Even though the AHI is the most used parameter for OSA severity estimation, it fails as a benchmark for detailing the OSA severity. Thus, parameters describing physiological cardiopulmonary consequences are needed [5–7]. Moreover, a greater understanding of the connection between respiratory events (breathing cessations and obstructions) and cardiovascular electrophysiological alterations may provide useful clinical insights into the causes and prevention of SCD. Therefore, we investigated this association in a population comprising OSA patients having suffered a stroke. Our main hypothesis was that deeper and longer desaturation cause greater prolongation of the QTc interval in OSA patients with a previous stroke. We aimed to study whether OSA-related desaturations are associated with repolarization prolongation and to investigate which desaturation features (e.g., length or depth) causes the highest risk for QTc prolongation predisposing to a possible risk for ventricular arrhythmias.

## Methods

### Study population

The initial dataset used in this study comprised ( $n=916$ ) suspected OSA patients, who had undergone type I full diagnostic polysomnography (PSG) at the Princess Alexandra Hospital (Brisbane, Australia) during 2015–2017. The PSGs were performed by using the Compumedics Grael acquisition system (Compumedics, Abbotsford, Australia). From the initial dataset, patients with a previous stroke and controls were matched based on OSA severity.

Approval for retrospective data collection was obtained from the Institutional Human Research

Ethics Committee of the Princess Alexandra Hospital (HREC/16/QPAH/021 and LNR/2019/QMS/54313). All procedures were performed following the ethical standards of the institutional and national research committee, and in accordance with the Helsinki declaration. The need for informed consent was waived by the Metro South Human Research Ethics Committee due to the retrospective nature of the study.

Out of the initial data set, 21 subjects fulfilled the inclusion criteria for this study: previous cerebrovascular stroke and at least moderate sleep apnea (AHI  $\geq 15$  events/h) with a total sleep time of  $\geq 4$  h in the PSG. Within these patients, the quality of ECG was insufficient in two cases and one patient was excluded due to ventricular pacing. No other exclusion criteria were used. Finally, eighteen patients were included in the further analyses. The reference population ( $n=18$ ) was matched based on OSA severity from the same dataset with the same inclusion criteria but excluding patients with a history of stroke. Demographic information for the stroke and control population are presented in Tables 1 and 2, respectively.

### PSG analysis

The PSG recordings were manually scored according to the American Academy of Sleep Medicine 2012 guidelines [9] by experienced sleep technicians in the Princess Alexandra Hospital using Compumedics ProFusion PSG 4 software (Compumedics, Abbotsford, Australia). All  $\geq 3\%$  desaturations were scored from the onset of the desaturation to the recovery of oxygenation and all desaturations during wake were excluded. The scoring process is described in a previous paper [10]. The detailed information of each desaturation was exported from ProFusion to MATLAB (R2019b; Mathworks, Natick, MA, USA) for data processing.

### ECG analysis related to desaturation events

ECG signal samples were extracted based on the start and end times of each desaturation. The nocturnal ECGs (lead II) were recorded during the PSG study with a sampling frequency of 256 Hz. The ECG signal was examined in three parts: (1) 10-s pre-desaturation ECG sample before the onset of the desaturation, (2) ECG sample within-desaturation, and (3) 15-s post-desaturation ECG sample after the end of desaturation (Fig. 1). If the post- and pre-desaturation ECG samples of two consecutive desaturations overlapped, all samples of the latter desaturation were excluded from further analyses. In addition, desaturations with a duration of less than 10 s were excluded. After these exclusions, a total of 975 (out of 2310) ECG samples were included in further analyses for the stroke population. For the reference population

**Table 1** Demographic, PSG, and ECG characteristics in the stroke population

Demographics	n/Median (%/IQR)	ECG characteristics	Median (IQR)
Patients (male)	18 (66.7)	RR interval (ms)	923.9 (834.1–1060.2)
Age (years)	67.9 (55.9–71.9)	PQ interval (ms)	188.9 (174.0–212.6)
BMI (kg/m <sup>2</sup> )	35.8 (33.7–41.8)	QRS interval (ms)	94.0 (85.3–102.0)
Atrial fibrillation	1 (5.6)	QT interval (ms)	417.9 (384.0–452.8)
COPD	1 (5.6)	QTc interval (ms)	424.1 (409.7–455.3)
Dyslipidemia	8 (44.4)	<b>Nocturnal HVR</b>	<b>Median (IQR)</b>
Hypertension	13 (72.2)	SDNN (ms)	33.7 (16.0–41.2)
Smoker	2 (11.1)	RMS-SD (ms)	21.7 (14.1–47.4)
<b>PSG characteristics</b>	<b>Median (IQR)</b>	LF (m <sup>2</sup> )	404.4 (134.9–710.6)
AHI (events/h)	35.3 (23.5–45.5)	HF (m <sup>2</sup> )	175.6 (43.3–519.5)
ODI (events/h)	23.1 (14.9–37.1)	LF/HF	1.4 (0.6–3.4)
Desaturation duration (s)	28.0 (19.6–41.0)	VLF (m <sup>2</sup> )	188.8 (56.8–334.9)
Desaturation depth (%)	4.3 (3.1–6.5)	SD1 (ms)	15.4 (10.0–33.5)
TST (min)	319.0 (285.3–351.9)	SD2 (ms)	40.8 (20.7–49.4)
Arousal index (events/h)	32.3 (24.1–44.1)	SD1/SD2	1.7 (1.0–2.9)

PSG polysomnography, ECG electrocardiogram, n number of patients, BMI body mass index, COPD chronic obstructive pulmonary disease, IQR interquartile range, AHI apnea–hypopnea index, ODI oxygen desaturation index, TST total sleep time, HRV heart rate variability, SDNN standard deviation of NN intervals, RMS-SD root mean square of successive RR interval differences, LF absolute power of the low-frequency band, HF absolute power of the high-frequency band, LF/RF ratio of LF-to-HF power, VLF absolute power of the very-low-frequency band, SD1 Poincaré plot standard deviation perpendicular the line of identity, SD2 Poincaré plot standard deviation along the line of identity, SD1/SD2 ratio of SD1-to-SD2

**Table 2** Demographic, PSG, and ECG characteristics in the reference population


Demographics	n/Median (%/IQR)	ECG characteristics	Median (IQR)
Patients (male)	18 (72.2)	RR interval (ms)	887.1 (808.1–998.3)
Age (years)	48.3 (46.2–49.6)	PQ interval (ms)	179.7 (164.1–191.6)
BMI (kg/m <sup>2</sup> )	35.1 (31.0–37.9)	QRS interval (ms)	93.8 (85.0–102.1)
Atrial fibrillation	1 (5.6)	QT interval (ms)	390.6 (359.4–410.2)
COPD	0 (0)	QTc interval (ms)	405.0 (392.7–420.8)
Dyslipidemia	6 (33.3)	<b>Nocturnal HVR</b>	<b>Median (IQR)</b>
Hypertension	7 (38.9)	SDNN (ms)	38.2 (17.7–54.2)
Smoker	3 (16.7)	RMS-SD (ma)	29.7 (14.7–52.5)
<b>PSG characteristics</b>	<b>Median (IQR)</b>	LF (m <sup>2</sup> )	563.2 (180.6–1456.1)
AHI (events/h)	29.8 (24.3–37.8)	HF (m <sup>2</sup> )	320.5 (79.1–788.4)
ODI (events/h)	25.1 (13.2–40.7)	LF/HF	2.1 (1.3–3.6)
Desaturation duration (s)	30.0 (21.1–40.0)	VLF (m <sup>2</sup> )	186.7 (91.9–481.5)
Desaturation depth (%)	4.7 (3.4–7.9)	SD1 (ms)	21.9 (10.4–37.2)
TST (min)	339.8 (321.0–368.5)	SD2 (ms)	44.5 (23.7–67.7)
Arousal index (events/h)	28.5 (23.9–41.2)	SD1/SD2	2.1 (1.8–2.7)

PSG polysomnography, ECG electrocardiogram, n number of patients, BMI body mass index, COPD chronic obstructive pulmonary disease, IQR interquartile range, AHI apnea–hypopnea index, ODI oxygen desaturation index, TST total sleep time, HRV heart rate variability, SDNN standard deviation of NN intervals, RMS-SD root mean square of successive RR interval differences, LF absolute power of the low-frequency band, HF absolute power of the high-frequency band, LF/RF ratio of LF-to-HF power, VLF absolute power of the very-low-frequency band, SD1 Poincaré plot standard deviation perpendicular the line of identity, SD2 Poincaré plot standard deviation along the line of identity, SD1/SD2 ratio of SD1-to-SD2

( $n = 18$ ), a total of 1070 (out of 3277) ECG samples were included in the analyses.

After extracting the ECG samples, the R-wave fiducial points were detected using Kubios HRV Premium software (Kubios Oy, Kuopio, Finland) [11] and detections

were verified by visual inspection of all extracted ECG samples (Fig. 1). Ventricular extrasystoles were excluded from the analysis. The average waveform for electrical activity was computed for each ECG sample in all three categories (i.e. before, within, and after the related



desaturation) to produce averaged P-QRS-T segments. The onset and offset times of the averaged P-, QRS-, and T-waves were manually set for each sample using Kubios HRV software [11] by trained ECG readers (Fig. 1). Heart rate corrected QT (QTc) intervals were calculated using Bazett's formula [12].

The nocturnal heart rate variability (HRV) parameters were analysed with Kubios Premium software. The ECG samples 30-min after the onset of PSG recording and 30-min before the end of PSG recordings were excluded from the nocturnal HRV analysis. The average of the analysed HRV recording was 6 h.

To investigate the effect of desaturation characteristics on QTc intervals, the desaturations were first grouped based on their duration ( $10 \leq s < 20$ ,  $20 \leq s < 30$ ,  $30 \leq s < 45$ , and  $\geq 45$  s) and depth, i.e. the magnitude of blood oxygen saturation (SpO<sub>2</sub>) drop ( $3\% \leq \Delta\text{SpO}_2 < 5\%$ ,  $5\% \leq \Delta\text{SpO}_2 < 7\%$ ,  $7\% \leq \Delta\text{SpO}_2 < 9\%$ , and  $\Delta\text{SpO}_2 \geq 9\%$ ).

### Statistical analysis

The statistical significance of the differences in QTc intervals between different duration and depth groups was evaluated using Wilcoxon's rank-sum (W) test. Additionally, the statistical significance of differences between

pre-desaturation QTc intervals and within-desaturation QTc intervals was evaluated using Wilcoxon's signed-rank sum (T) test as the within-desaturation QTc was assumed to be dependent on the pre-desaturation QTc time. The limit for statistical significance was set to be  $p < 0.05$ . The statistical significance of differences between the reference population and the stroke population were evaluated with Wilcoxon's W test (baseline characteristics) and the two-sided Kolmogorov–Smirnov test (distributions of  $\Delta\text{QTc}$ ). Moreover, generalized linear regression and stepwise linear regression were conducted to analyze the effect of demographics, conventional OSA severity, and event-by-event desaturation characteristics on the  $\Delta\text{QTc}$  between pre- and within-desaturation segments. In stepwise regression, deviance with  $p$ -value's F-statistics was used as entering and removing criteria.

### Results

QTc was prolonged (at least 1 ms) during 52.6% of desaturations in the stroke population. Moreover, QTc was prolonged for at least 11 ms during 27.1% of desaturations and over 20 ms during 12.2% of desaturation (Table 3). In the reference population, most of the prolongations were minor (1–11 ms, 38.2%), and only 4.0%

**Table 3** The effect of desaturation characteristics on QTc intervals in the stroke and control population

$\Delta$ QTc	Number of desaturations	Desaturation duration (s)	Desaturation depth (%)
<i>Stroke population</i>			
1–11 ms	264 (27.1%)	28.2 (19.8–40.0)	3.9 (3.0–7.4)
11–16 ms	82 (8.4%)	40.0 (26.0–49.0)	5.0 (3.4–11.2)
16–20 ms	48 (4.9%)	30.5 (22.5–44.0)	4.1 (3.2–7.2)
> 20 ms	119 (12.2%)	30.0 (18.1–43.8)	4.7 (3.4–7.3)
<i>Control population</i>			
1–11 ms	409 (38.2%)	30.0 (22.0–40.0)	4.6 (3.3–7.2)
11–16 ms	57 (5.3%)	28.0 (18.0–40.5)	4.3 (3.2–7.3)
16–20 ms	37 (3.5%)	34.0 (22.9–42.8)	4.7 (3.9–6.2)
> 20 ms	43 (4.0%)	34.0 (18.3–43.8)	5.2 (3.7–7.0)

Desaturations are grouped according to the absolute change in the heart rate corrected QT interval ( $\Delta$ QTc) within desaturations. The median and (interquartile range) of desaturation duration (s) and depth (%) are presented related to each  $\Delta$ QTc group

of the desaturations caused over 20 ms prolongation (Table 3).


To further study the influence of desaturation characteristics on QTc among stroke patients, desaturations were also categorized into sub-groups according to the desaturation duration (Table 4; Fig. 2). The pre-desaturation QTc was longer before  $30 \leq s < 45$  and  $\geq 45$  s desaturations compared to short ( $10 \leq s < 20$ )

desaturations ( $p < 0.01$  for both) (Table 4). There was no statistical significance in pre-desaturation QTc durations between the other desaturation duration groups. Furthermore, the QTc interval within-desaturations was significantly ( $p < 0.01$ ) longer compared to the corresponding pre-desaturation QTc intervals in the  $30 \leq s < 45$  and  $\geq 45$  s desaturation duration groups (Table 4; Fig. 2A). In shorter desaturation

**Table 4** Median (interquartile range) changes in QTc in the stroke and control population

	Pre-desaturation QTc (ms)	$\Delta$ QTc within-desaturation (ms)	$\Delta$ QTc post-desaturation (ms)
<i>Stroke population</i>			
Desaturation duration			
$10 \leq s < 20$	418.9 (405.4–439.4)	0.8 (–8.9–9.7)	<b>2.3 (–7.4–11.0)</b>
$20 \leq s < 30$	422.3 (409.5–453.5)	–1.1 (–8.0–8.7)	–1.0 (–11.3–7.8)
$30 \leq s < 45$	427.4 (412.7–457.2)*	<b>3.4 (–6.9–12.3)</b>	<b>2.4 (–6.3–10.5)</b>
$\geq 45$ s	430.6 (412.6–460.9)*	<b>7.5 (–3.0–14.8)*</b>	0.5 (–10.3–10.8)
Desaturation depth			
$3\% \leq \Delta$ SpO <sub>2</sub> < 5%	424.8 (409.1–454.9)	0.6 (–8.9–11.1)	0.8 (–11.5–9.5)
$5\% \leq \Delta$ SpO <sub>2</sub> < 7%	424.8 (414.1–458.3)	–1.4 (–9.7–10.5)	–1.4 (–9.8–9.4)
$7\% \leq \Delta$ SpO <sub>2</sub> < 9%	424.5 (408.9–444.1)	<b>4.6 (–3.2–12.3)*</b>	2.0 (–4.8–7.8)*
$\Delta$ SpO <sub>2</sub> $\geq 9\%$	424.3 (413.7–437.4)	<b>7.4 (–1.1–14.9)*</b>	<b>3.7 (–5.6–10.0)†</b>
<i>Control population</i>			
Desaturation duration			
$10 \leq s < 20$	416.2 (390.9–436.7)	1.4 (–5.2–8.1)	–1.7 (–7.9–5.9)
$20 \leq s < 30$	405.8 (392.1–424.9)	0.8 (–5.1–6.4)	0.6 (–6.2–7.6)
$30 \leq s < 45$	401.5 (391.2–413.4)*	0.3 (–4.5–5.6)	2.2 (–3.6–10.1)*
$\geq 45$ s	403.2 (391.3–414.5)*	<b>3.7 (–2.6–8.9)†</b>	–0.4 (–4.9–6.2)
Desaturation depth			
$3\% \leq \Delta$ SpO <sub>2</sub> < 5%	402.6 (387.0–423.1)	<b>1.8 (–3.7–6.8)</b>	0.9 (–4.7–7.9)
$5\% \leq \Delta$ SpO <sub>2</sub> < 7%	405.0 (390.8–422.0)	<b>1.8 (–3.5–8.2)</b>	0.8 (–5.8–9.5)
$7\% \leq \Delta$ SpO <sub>2</sub> < 9%	405.5 (397.3–421.4)	1.3 (–6.5–6.1)	1.2 (–7.8–6.7)
SpO <sub>2</sub> $\geq 9\%$	404.2 (396.3–414.9)	–0.7 (–7.3–5.1)*	2.6 (–4.9–9.6)

Change in the within-desaturation segment ( $\Delta$ QTc within-desaturation) and the post-desaturation segment ( $\Delta$ QTc post-desaturation) were calculated based to the pre-desaturation QTc segment values. Bolded values indicate a statistically significant ( $p < 0.05$ ) change from the pre-desaturation QTc value. \*The value differs significantly from the 10–20 s or 3–5% group. †The value differs significantly from all duration or depth groups. Abbreviations:  $\Delta$ SpO<sub>2</sub> = drop in the blood oxygen saturation during desaturation



duration groups, QTc intervals did not change significantly ( $p > 0.11$ ) within-desaturations compared to pre-desaturation QTc intervals. The median QTc was slightly longer after  $10 \leq s < 20$  and  $30 \leq s < 45$  desaturations compared to corresponding pre-desaturation QTc intervals (2.3 and 2.4 ms,  $p < 0.05$  for both). There were no statistically significant differences in pre-desaturation and post-desaturation QTc intervals in  $20 \leq s < 30$  and  $\geq 45$  s desaturation duration groups (Table 4). No statistically significant difference between pre-desaturation and post-desaturation QTc in the stroke population (Table 4). In the reference population, only mild prolongation was seen post-desaturation after  $10 \leq s < 20$  (median 2.3 ms) and  $30 \leq s < 45$  (median 2.4 ms) (Table 4). In the reference population

pre-desaturation QTc was longer before  $10 \leq s < 20$  compared to  $\geq 30$  s desaturations ( $p < 0.05$ ). Only the  $\geq 45$  s desaturations caused significant ( $p < 0.01$ ) prolongation of the QTc among the reference population. Furthermore, in  $30 \leq s < 45$  and  $\geq 45$  s desaturation duration groups, stroke patients had a significantly ( $p < 0.04$ ) greater change between pre-desaturation and within-desaturation QTc intervals compared to the reference population (Fig. 2A, C).

We also categorized desaturations into sub-groups according to the depth of desaturation. There were no statistically significant differences in pre-desaturation QTc intervals in the stroke group (Table 4). However, the QTc prolonged significantly during  $7\% \leq \Delta SpO_2 < 9\%$  and  $\geq 9\%$  desaturations (Table 4; Fig. 2B). QTc did not

prolong significantly during <7% desaturations. Compared to pre-desaturation QTc, the post-desaturation QTc was slightly elevated (median 3.7 ms) after  $\geq 9\%$  desaturations in stroke patients. No statistically significant difference between pre-desaturation and post-desaturation QTc intervals was observed in other desaturation depth groups or in the reference population (Table 4). In the reference population, the QTc prolongations were minor, and the deeper desaturations did not similarly affect the QTc than in stroke patients (Table 4). Moreover, the deep ( $\Delta\text{SpO}_2 \geq 7\%$ ) desaturations caused significantly ( $p < 0.01$ ) greater QTc prolongation in the stroke patients compared to the reference population.

In regression analysis, both in the general linear (GL) model as well as in the stepwise (SW) model, only the total sleep time in PSG ( $\beta_{\text{GL}} = .366$ ,  $\beta_{\text{SW}} = -1.549$ ,  $p < 0.01$ ) and desaturation duration ( $\beta_{\text{GL}} = 0.065$ ,  $\beta_{\text{SW}} = 0.080$ ,  $p < 0.002$ ) were significant predictors of the increase in  $\Delta\text{QTc}$  (Table 5). Conversely, age, sex, or AHI were not predictors of acute QTc prolongation.

## Discussion

In the present study, we evaluated the acute influence of desaturation length and depth in repolarization duration in OSA patients with a previously diagnosed stroke,

and OSA severity matched counterparts without a previous stroke. We found that longer and deeper desaturations can prolong QTc interval during a desaturation, especially in patients with previous stroke. Moreover, the desaturation length was an independent predictor for QTc prolongation in both stroke and control populations.

Intermittent hypoxemia is assumed to promote cardiac arrhythmias by producing oxidative stress, systemic inflammation, and increased sympathetic activity leading to anatomical and electrophysiological remodeling [13, 14]. Interestingly, it has been shown that hypoxia induces QTc prolongation also in healthy subjects [15]. Still, the pathophysiological cascade between breathing cessations, autonomic dysregulation, and fatal cardiovascular events is partly unclear. Based on previous studies, we hypothesized that desaturation leads to acute intermittent prolongation of the QTc. Since the autonomic nervous system (ANS) plays an important role in the regulation of repolarization [16] we focused on OSA patients with a previous stroke as stroke can lead to impaired autonomic control [19]. As hypothesized, we found that as the heart rate increases within-desaturations the shortening of the QTc interval was inadequate among OSA patients with a previous stroke. We further evaluated ANS activation in our patient populations with nocturnal HRV. The stroke patient had slightly lower median HRV parameters compared to the control population, which can suggest that they tend to have lower ANS function.

OSA causes acute and long-time sympathetic autonomic nervous system activation by increasing pulmonary artery pressure and chemoreceptor activation caused by acidosis, hypoxia, and hypercapnia [17]. Previously, Tavares et al. have shown that when blood  $\text{SaO}_2$  levels decline, the sympathetic activation increases progressively [17]. Moreover, a cohort study showed that the lowest nocturnal  $\text{O}_2$  saturation value and the mean  $\text{O}_2$  saturation at night, are independent predictors of SCD [4]. These findings are in line with our results since we found that more substantial desaturations are associated with QTc prolongation which is a risk factor for lethal arrhythmias [8]. The ANS activation alters not only heart rate, conduction, and hemodynamics, but also the cellular and subcellular properties of individual myocytes [18]. A previous study has shown that intrathoracic pressure swings cause acute left-sided heart morphology and function changes [19] and this may lead to electrophysiological remodeling and further to the manifestation of arrhythmias.

The QTc prolongation seen in the present study might explain the increased risk for nocturnal SCD of OSA patients with long desaturations for the following reasons. First, it has been shown that the frequency and duration of respiratory and desaturation events increase

**Table 5** Generalized linear regression model and a stepwise regression model for absolute  $\Delta\text{QTc}$  (ms) between within-desaturation period and pre-desaturation in stroke and control population

Generalized model	$\beta$	SE	T	p
Sex	0.578	0.774	0.747	0.455
Age (y)	0.049	0.037	1.310	0.190
TST (h)	-1.366	0.524	-2.591	0.010
AHI (1/h)	-0.031	0.018	-1.725	0.09
Desaturation duration (s)	0.065	0.022	2.978	0.002
$\Delta\text{SpO}_2$ (%)	0.069	0.077	0.895	0.371
Stepwise model	$\beta$	SE	T	p
Sex	-	-	-	-
Age (y)	-	-	-	-
TST (h)	-1.549	0.457	-3.391	<0.001
AHI (1/h)	-	-	-	-
Desaturation duration (s)	0.080	0.021	3.818	<0.001
$\Delta\text{SpO}_2$ (%)	-	-	-	-

All other parameters except Desaturation duration and  $\Delta\text{SpO}_2$  are treated patient-wise; the two are computed for each segment separately. The criteria for excluding parameters from the stepwise model was Deviance and  $p$ -value for F-statistics. Abbreviations: TST total sleep time, AHI apnea-hypopnea index,  $\Delta\text{SpO}_2$  = depth of desaturation

towards the morning [20]. Second, it is known that both prolonged QTc and dynamic changes in QT interval increase the risk for arrhythmias [21]. Third, it has been shown that the risk for sudden death from cardiac causes is increased in OSA patients from midnight to 6 a.m. [22]. These previous findings reinforce the association between pro-arrhythmogenic QTc prolongation and long (>30 s) and deep (>7%) desaturations.

The retrospective nature of the study leads to certain limitations. First, we had no information available on the anatomical location of the stroke or the time between stroke and PSG/ECG recordings. Thus, there is the possibility that autonomic regulation may improve during the intervening period between the stroke event and the diagnostic PSG. However, a previous study has shown that the long-term negative effects on cardiovascular regulation can still be seen after stroke [23]. Secondly, the stroke patients were older compared to controls, the difference in acute electrophysiological response on desaturation may be also influenced by longer exposure to the pathological insults associated with sleep apnea and other comorbidities. However, the results of regression analyses showed that when sex, age, and conventional OSA severity are considered alongside desaturation metrics, desaturation duration was the only independent predictor of acute QTc prolongation together with total sleep time (Table 5). Therefore, even though the stroke population was older, and had higher baseline QTc, the acute effect of desaturation to repolarization is significant and modulated especially by the length of the desaturation event. Thus, the difference in QTc prolongation between the populations cannot be explained by age or AHI. Yet, we know that TST is related to age so the age might not be completely irrelevant and further analyses outside the scope of this study are needed.

Moreover, we had to assume an appropriate evaluation window for ECG samples before and after desaturation. For this reason, we excluded desaturations originating less than 25 s after the previous desaturation had ended. It is possible that the exclusion of a significant number of desaturations can cause some bias in the result and can partly explain why short ( $10 \leq s < 20$ ) desaturations seem to cause a more substantial influence on QTc than longer ( $20 \leq s < 30$ ) desaturations (Fig. 2A). Furthermore, it is common that patients with very severe sleep apnea suffer from frequent desaturations that are short but cause rapid drops in SpO<sub>2</sub> level and can still be very deep. Therefore, the desaturations of these patients would be mostly represented in the short ( $10 \leq s < 20$ ) desaturation group. This can also partly explain why short desaturations seem to have a greater effect on QTc interval. Another reason behind this unexpected finding can be due to desaturations being estimated from a level just

before desaturation onset and wake-time normal blood saturation levels are not considered. It is possible that in these measurements, the short events start during already lower SpO<sub>2</sub> levels, causing SpO<sub>2</sub> to drop faster and further induce QTc prolongation. Last, the QT measuring and correction can be challenging. In this study, the QTc was measured semiautomatically from the lead II which is the most often used in QT interval duration measurements because it is more likely to have the longest QT interval [24]. In addition, several approaches have been developed to correct the QT interval. It is known that the Bazett's correction formula is not optimal due to its tendency to underestimate and overestimate the duration of cardiac repolarization at extremely low and high heart rates. However, we chose to use the Bazett's formula as it is the most used and universal QT interval correction method to achieve results that are easily comparable with the literature. The median heart rate in this study was 64.9 beats/min in the stroke population and 67.6 beats/min in the reference population (Tables 1, 2). Thus, as there were no major differences in median heart rates, we consider that using the Bazett's formula does not have major effects on the obtained results.

### Clinical implications and conclusions

In the stroke population, 12.2% of desaturations induced  $\geq 20$  ms prolongation in QTc (Table 3). Generally, when drug safety is estimated, prolongation of QTc of  $\geq 20$  ms is associated with a substantial likelihood of the drug being proarrhythmic [25]. Therefore, we consider that OSA should be recognized as a risk factor for the acquired long-QT syndrome (LQTS) in stroke patients. Moreover, when describing medication associated with QT prolongation possible nocturnal QTc prolongation should be noticed. We believe that this is essential, especially in OSA patients diagnosed with LQTS or if they have borderline daytime QT levels.

In the current clinical practice, atrial fibrillation is routinely screened with 24-h Holter monitoring if no obvious cause of stroke can be found. However, routine screening for sleep-disordered breathing is uncommon in patients with stroke even though OSA is overrepresented in this patient group [3]. Combining home polysomnography with the 24-h Holter monitoring would be a feasible and possibly beneficial addition to stroke patients' clinical risk evaluation. This is because sleep apnea is an important target for public health interventions aiming at reducing the risk for arrhythmias and SCD. Moreover, continuous positive airway pressure treatment has been shown to significantly decrease QTc in moderate/severe OSA patients [26]. Our finding might partially explain why continuous positive airway pressure therapy, which decreases the frequency and duration of desaturations,



prevents cardiac arrhythmias [13]. This finding endorses the need to screen and treat OSA, especially in patients with a previous stroke.

#### Acknowledgements

Not applicable.

#### Author contributions

B.D. provided the patient data and J.L. the Kubios software. S.S., S.K., and A.S. analysed the patient data. All authors interpreted the patient data and contributed to writing the manuscript. All authors read and approved the final manuscript.

#### Funding

This work was supported by the Research Committee of the Kuopio University Hospital Catchment Area for the State Research Funding [5041798 to S.S., 5041767 to T.L., 5041794 to T.L., 5041803 to H.K.]; the European Union's Horizon 2020 research and innovation program [965417 to T.L.]; the Academy of Finland [323536 to T.L.], the Competitive State Research Financing of Expert Responsibility Area of Tampere University Hospital [VTR3242 to A.K., VTR3249 to A.K., VTR 3256 to A.K., EVO2089 to A.K.]; NordForsk [90458 to J.T. and T.L.] via Business Finland [5133/31/2018 to J.T. and T.L.]; the Research Foundation of the Pulmonary Diseases [to H.K. and to S.K.]; the Finnish Anti-Tuberculosis Foundation [to H.K.]; the Finnish Cultural Foundation—Pohjois-Savo regional fund [to S.K.]; and the Maud Kuistila Memorial Foundation [to S.K.].

#### Availability of data and materials

The datasets analysed during the current study are not publicly available due to privacy policy but are available from the corresponding author on reasonable request.

#### Declarations

##### Ethics approval and consent to participate

Approval for retrospective data collection was obtained from the Institutional Human Research Ethics Committee of the Princess Alexandra Hospital (HREC/16/QPAH/021 and LNR/2019/QMS/54313). The need for informed consent was waived by the Metro South Human Research Ethics Committee due to the retrospective nature of the study.

##### Consent to publish

Not applicable.

##### Competing interests

Lipponen JA is a shareholder of a company (Kubios) that designs ECG-analysis and heart rate variability analysis software. Other co-authors have nothing to declare.

##### Author details

<sup>1</sup>Diagnostic Imaging Center, Kuopio University Hospital (KYS), P.O. Box 100, 70029 Kuopio, Finland. <sup>2</sup>Department of Applied Physics, University of Eastern Finland, Kuopio, Finland. <sup>3</sup>Department of Clinical Neurophysiology, Seinäjoki Central Hospital, Seinäjoki, Finland. <sup>4</sup>Science Service Center, Kuopio University Hospital, Kuopio, Finland. <sup>5</sup>The School of Information Technology and Electrical Engineering, The University of Queensland, Brisbane, Australia. <sup>6</sup>Department of Respiratory and Sleep Medicine, Sleep Disorders Centre, Princess Alexandra Hospital, Ipswich Rd, Woolloongabba, QLD, Australia. <sup>7</sup>Institute for Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia. <sup>8</sup>The School of Medicine, University of Eastern Finland, Kuopio, Finland.

Received: 18 February 2022 Accepted: 10 May 2022

Published online: 24 May 2022

#### References

- Benjafeld AV, Ayas NT, Eastwood PR, Heinzer R, Ip MSM, Morrell MJ, Nunez CM, Patel SR, Penzel T, Pepin JL, Peppard PE, Sinha S, Tufik S,

- Valentine K, Malhotra A. Estimation of the global prevalence and burden of obstructive sleep apnoea: a literature-based analysis. *Lancet Respir Med.* 2019;7(8):687–98.
- Redline S, Yenokyan G, Gottlieb DJ, Shahar E, O'Connor GT, Resnick HE, Diener-West M, Sanders MH, Wolf PA, Geraghty EM, Ali T, Lebowitz M, Punjabi NM. Obstructive sleep apnea-hypopnea and incident stroke: the sleep heart health study. *Am J Respir Crit Care Med.* 2010;182(2):269–77.
- Leino A, Westeren-Punnonen S, Toyra J, Myllymaa S, Leppanen T, Yla-Herttua S, Muraja-Murro A, Kantanen AM, Autere J, Jakala P, Mervaala E, Myllymaa K. Acute stroke and TIA patients have specific polygraphic features of obstructive sleep apnea. *Sleep Breath.* 2020;24(4):1495–505.
- Gami AS, Olson EJ, Shen WK, Wright RS, Ballman KV, Hodge DO, Herges RM, Howard DE, Somers VK. Obstructive sleep apnea and the risk of sudden cardiac death: a longitudinal study of 10,701 adults. *J Am Coll Cardiol.* 2013;62(7):610–6.
- Soros P, Hachinski V. Cardiovascular and neurological causes of sudden death after ischaemic stroke. *Lancet Neurol.* 2012;11(2):179–88.
- Wong KY, Mac Walter RS, Douglas D, Fraser HW, Ogston SA, Struthers AD. Long QTc predicts future cardiac death in stroke survivors. *Heart.* 2003;89(4):377–81.
- Huikuri HV, MD, Castellanos A, MD, Myerburg RJ, MD: Sudden death due to cardiac arrhythmias. *N Engl J Med.* 2001;345(20):1473–82.
- Chugh SS, Reinier K, Singh T, Uy-Evanado A, Socoteanu C, Peters D, Mariani R, Gunson K, Jui J. Determinants of prolonged QT interval and their contribution to sudden death risk in coronary artery disease: the Oregon Sudden Unexpected Death Study. *Circulation.* 2009;119(5):663–70.
- Berry RB, Budhiraja R, Gottlieb DJ, Gozal D, Iber C, Kapur VK, Marcus CL, Mehra R, Parthasarathy S, Quan SF, Redline S, Strohl KP, Davidson Ward SL, Tangredi MM, American Academy of Sleep Medicine: Rules for scoring respiratory events in sleep: update of the 2007 AASM manual for the scoring of sleep and associated events. Deliberations of the Sleep Apnea Definitions Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2012, 8(5):597–619.
- Duce B, Kulkas A, Langton C, Töyräs J, Hukins C. The AASM 2012 recommended hypopnea criteria increase the incidence of obstructive sleep apnea but not the proportion of positional obstructive sleep apnea. *Sleep Med.* 2016;26:23–9.
- Tarvainen MP, Niskanen JP, Lipponen JA, Ranta-Aho PO, Karjalainen PA. Kubios HRV—heart rate variability analysis software. *Comput Methods Programs Biomed.* 2014;113(1):210–20.
- Rautaharju PM, Surawicz B, Gettes LS, Bailey JJ, Childers R, Deal BJ, Gorgels A, Hancock EW, Josephson M, Kligfield P, Kors JA, Macfarlane P, Mason JW, Mirvis DM, Okin P, Pahlm O, van Herpen G, Wagner GS, Wellens H. American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology, American College of Cardiology Foundation, Heart Rhythm Society: AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: part IV: the ST segment, T and U waves, and the QT interval: a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society: endorsed by the International Society for Computerized Electrocardiology. *Circulation.* 2009;119(10):241.
- Rossi VA, Stradling JR, Kohler M. Effects of obstructive sleep apnoea on heart rhythm. *Eur Respir J.* 2013;41(6):1439–51.
- Walker M, Blackwell JN, Stafford P, Patel P, Mazimba S, Mehta N, Cho Y, Mangrum M, Nazarian S, Bilchick K, Kwon Y. Daytime QT by routine 12-lead ECG is prolonged in patients with severe obstructive sleep apnea. *Sleep Disord.* 2020;2020:3029836.
- Roche Fr, Reynaud C, Pichot V, Duverney D, Costes Fr, Garet M, Gaspoz J, Barthélémy J. Effect of acute hypoxia on QT rate dependence and corrected QT interval in healthy subjects. *Am J Cardiol* 2003, 91(7):916–9.
- Shen MJ, Zipes DP. Role of the autonomic nervous system in modulating cardiac arrhythmias. *Circ Res.* 2014;114(6):1004–21.
- Tavares L, Rodriguez-Manero M, Kreidieh B, Ibarra-Cortez SH, Chen J, Wang S, Markovits J, Barrios R, Valderrabano M. Cardiac afferent denervation abolishes ganglionated plexi and sympathetic responses to apnea: implications for atrial fibrillation. *Circ Arrhythm Electrophysiol.* 2019;12(6):e006942.

18. Morand J, Arnaud C, Pepin JL, Godin-Ribuot D. Chronic intermittent hypoxia promotes myocardial ischemia-related ventricular arrhythmias and sudden cardiac death. *Sci Rep*. 2018;8(1):2997.
19. Orban M, Bruce CJ, Pressman GS, Leinveber P, Romero-Corral A, Korinek J, Konecny T, Villarraga HR, Kara T, Caples SM, Somers VK. Dynamic changes of left ventricular performance and left atrial volume induced by the mueller maneuver in healthy young adults and implications for obstructive sleep apnea, atrial fibrillation, and heart failure. *Am J Cardiol*. 2008;102(11):1557–61.
20. Nikkonen S, Toyras J, Mervaala E, Myllymaa S, Terrill P, Leppanen T. Intra-night variation in apnea-hypopnea index affects diagnostics and prognostics of obstructive sleep apnea. *Sleep Breath*. 2020;24(1):379–86.
21. Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A, Daniels S, Floras JS, Hunt CE, Olson LJ, Pickering TG, Russell R, Woo M, Young T, American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, American Heart Association Stroke Council, American Heart Association Council on Cardiovascular Nursing, American College of Cardiology Foundation: Sleep apnea and cardiovascular disease: an American Heart Association/American College Of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council On Cardiovascular Nursing. In collaboration with the National Heart, Lung, and Blood Institute National Center on Sleep Disorders Research (National Institutes of Health). *Circulation* 2008, 118(10):1080–111.
22. Gami AS, Howard DE, Olson EJ, Somers VK. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med*. 2005;352(12):1206–14.
23. McLaren A, Kerr S, Allan L, Steen IN, Ballard C, Allen J, Murray A, Kenny RA. Autonomic function is impaired in elderly stroke survivors. *Stroke*. 2005;36(5):1026–30.
24. Malik M, Camm AJ. Evaluation of drug-induced QT interval prolongation: implications for drug approval and labelling. *Drug Saf*. 2001;24(5):323–51.
25. QT interval and drug therapy (Clinical review from Drug and Therapeutics Bulletin). *BMJ* 2016, 354:i4331.
26. Cicek D, Balcioglu AS, Lakadamyali H, Müderrisoğlu H. Effects of three month nasal continuous positive airway pressure treatment on electrocardiographic, echocardiographic and overnight polysomnographic parameters in newly diagnosed moderate/severe obstructive sleep apnea patients. *Int Heart J*. 2015;56(1):94–9.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

