

**Thinness, overweight, and obesity among 6- to 17-year-old Malaysians: Secular trends and sociodemographic determinants from 2006-2015**

Maria S. Mohamad<sup>1</sup>, Balkish Mahadir Naidu<sup>2</sup>, Riittakerttu Kaltiala<sup>3</sup>, Suvi M. Virtanen<sup>1,4,5,6</sup>,  
Susanna Lehtinen-Jacks<sup>1,7</sup>

<sup>1</sup>Unit of Health Sciences, Faculty of Social Sciences, Tampere University, Finland

<sup>2</sup>Research and Methodology Unit, Department of Statistics Malaysia, Putrajaya, Malaysia

<sup>3</sup>Department of Adolescent Psychiatry, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

<sup>4</sup>Public Health Promotion Unit, Finnish Institute for Health and Welfare, Helsinki, Finland

<sup>5</sup>Center for Child Health Research, Tampere University and Tampere University Hospital, Tampere, Finland

<sup>6</sup>The Science Center of Pirkanmaa Hospital District, Tampere, Finland

<sup>7</sup>Division of Public Health Sciences, School of Health, Care and Social Welfare, Mälardalen University, Västerås, Sweden

**Corresponding author:** Maria S. Mohamad, Mailing address: Unit of Health Sciences, Faculty of Social Sciences, Tampere University, Arvo Ylpön Katu 34, 33520, Tampere, Finland, Email address: [maria.mohamad@tuni.fi](mailto:maria.mohamad@tuni.fi), Telephone number: +3584578749260

**Short title:** Trends of malnutrition among Malaysian children.



This is an Accepted Manuscript for Public Health Nutrition as part of the Cambridge Coronavirus Collection. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process.

The article is considered published and may be cited using its DOI

10.1017/S1368980021003190

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

**Acknowledgements:** We would like to thank the Director General of Health Malaysia for the permission to use the datasets and publish this article.

**Financial Support:** None.

**Conflict of Interest:** None.

**Authorship:** M.S.M, B.M.N, R.K, S.M.V and S.L-J formulated the research questions and designed the study. M.S.M, B.M.N and S.L-J conducted the study and performed statistical analysis. M.S.M and S.L-J wrote paper. RK and SMV contributed (to the writing) with critical comments. All authors read and approved the final manuscript.

**Ethical Standards Disclosure:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the Medical Research Ethics Committee under the Ministry of Health Malaysia. Written informed consent was obtained from all participants.

## Abstract

**Objective:** To examine secular trends and sociodemographic determinants of thinness, overweight, and obesity among Malaysian children and adolescents in 2006-2015.

**Design:** We used cross-sectional data from the National Health and Morbidity Surveys 2006, 2011, and 2015. Individuals were classified into pre- (6-9 years), early (10-13 years) and mid (14-17 years) adolescence. BMI status was determined according to the IOTF and WHO criteria, using measured height and weight. We analysed trends using log-binomial regression, by sex-age groups, stratified by sociodemographic factors (ethnicity, residential area, household size, household income), and accounting for the complex survey design. Associations between sociodemographic factors and prevalence of thinness and overweight (obesity included) in 2015 were assessed using log-Poisson regression.

**Setting:** Nationwide population-based surveys, Malaysia.

**Participants:** Eligible 6-17-year-olds from urban and rural residential areas (N=28,094).

**Results:** The prevalence of thinness decreased from 2006 to 2015 (IOTF: boys from 22% to 18%, girls from 23% to 19%; WHO: boys from 9% to 7%, girls from 8% to 6%), while the prevalence of overweight increased (IOTF: boys from 20% to 26%, girls from 19% to 24%; WHO: boys from 25% to 31%, girls from 22% to 27%). These changes were statistically significant in most sex-age groups. Thinness and overweight co-existed in all sociodemographic subgroups, with variation in the prevalence estimates, but similar secular changes in most subgroups.

**Conclusions:** Malaysia is facing a double burden of malnutrition at population level with a secular increase in overweight and obesity and a gradual decrease in thinness among 6-17-year-olds from varying sociodemographic backgrounds.

**Keywords:** Child, Malaysia, Overweight, Thinness, Trends

## Introduction

The World Health Organisation (WHO) has defined the double burden of malnutrition as the coexistence of undernutrition with overweight and obesity that may manifest in individuals, households, and populations across the life course<sup>(1)</sup>. Undernutrition includes conditions such as stunting, wasting, thinness, and micronutrient deficiency<sup>(2)</sup>. Worldwide, 88% of countries are facing more than one form of malnutrition<sup>(3)</sup>. In 2016, it was estimated that 75 million girls and 117 million boys aged 5 to 19 years had thinness<sup>(4)</sup>. Despite decreasing global trends in prevalence of thinness from 1975 to 2016, the absolute numbers of children and adolescents with thinness increased in Asia and Africa due to population growth<sup>(4)</sup>. At the same time, the number of children and adolescents with obesity worldwide increased from 5 million to 50 million in girls and from 6 million to 74 million in boys, with a rapid increase among boys in Southeast Asia<sup>(4)</sup>. Both forms of malnutrition are associated with short- and long-term adverse health consequences. Thinness in childhood and adolescence is associated with an increased risk of mortality and morbidity. This could delay maturity and reduced physical work capacity and productivity<sup>(5,6)</sup>. In girls, thinness could increase the risk of adverse pregnancy outcomes and postmenopausal osteoporosis in later life<sup>(5,6)</sup>. Likewise, overweight and obesity in childhood and adolescence tend to persist into adulthood and are associated with physically and psychologically adverse health outcomes, including many chronic diseases and mental health disorders<sup>(7,8)</sup>.

Since its independence in 1957, Malaysia, a country located in Southeast Asia has experienced decades of a thriving economy and rapid development. From a country dominated by a rural region and depending on the agricultural sector at the time of independence, Malaysia has transformed into a highly urbanised country with dependency on the service sector economy. The rapid transformation has led to changes in the lifestyles and behaviours of Malaysians. Westernised diets, low levels of physical activity, and sedentary behaviours have become the norms<sup>(9)</sup>. Evidence shows that Malaysia is currently in the midst of nutrition and epidemiological transitions with increasing prevalence of overweight, obesity, and diet-related non-communicable diseases, with a high burden of undernutrition<sup>(9–11)</sup>.

A standardised and reliable reference to measure malnutrition in children and adolescents is essential to guide policy efforts to tackle malnutrition. Regarding thinness, overweight and obesity, body mass index (BMI) is nowadays commonly used in children and adolescents for growth monitoring, clinical management, and research<sup>(12)</sup>. In addition to local references, several international references are available, including the widely used age- and sex-specific references

developed by the International Obesity Task Force (IOTF)<sup>(13)</sup>, the WHO<sup>(14)</sup>, and the Centres for Disease Control and Prevention (CDC)<sup>(15)</sup>. Previous studies among Malaysian children and adolescents have used international references in different ways<sup>(16–20)</sup>. For example, the CDC Growth Chart 2000 reference was applied in the National Health and Morbidity Survey (NHMS) conducted in 2011 and 2015<sup>(18,19)</sup>. The prevalence of thinness among children and adolescents aged under 18 years decreased from 12% to 8%, while the prevalence of obesity doubled from 6% to 12% between the survey's intervals. In the most recent NHMS survey in 2019, the WHO 2007 reference was used<sup>(20)</sup>. Using that reference, the prevalence of thinness was 10%, while the prevalence of obesity was 15% among children and adolescents aged 5 to 17 years old<sup>(20)</sup>.

Several factors such as age, sex, ethnicity, urbanisation level of the place of residence, and socioeconomic factors including household income and household size have been associated with the double burden of thinness and overweight and obesity in previous studies among Malaysian children and adolescents<sup>(21–27)</sup>. For example, male sex<sup>(21,22)</sup>, Indian ethnic origin<sup>(24)</sup>, and urban residence<sup>(24)</sup> have been associated with higher prevalence of both thinness and obesity, while overweight was found to be as common in all major ethnic groups in the few national studies<sup>(21,22)</sup>. Higher household income has been associated with lower prevalence of thinness<sup>(24)</sup> and with higher prevalence of overweight<sup>(26)</sup> or obesity<sup>(24)</sup>.

The double burden of malnutrition imposes a challenge in terms of the healthcare and wellbeing of the population. In addition to the high burden of undernutrition, it is predicted that 1.8 million Malaysians aged 5 to 19 years would be living with obesity in 2030<sup>(28)</sup>. To our knowledge, there were no previous studies using nationally representative data looking at secular trends and sociodemographic determinants of both thinness and overweight and obesity among Malaysian children and adolescents. In addition, the use of different BMI references in the previous studies complicates comparisons across studies locally and internationally. Therefore, in order to develop effective strategies targeting the most afflicted parts of the population, the present study aims to provide vital information about time trends and potential sociodemographic differences in the prevalence of thinness and overweight (obesity included) at the population level in 6- to 17-year-old children and adolescents in Malaysia. For comparability at the local and international level, the present study applied the IOTF and the WHO BMI criteria as references for child and adolescent BMI status.

## Methods

### *Study Design and Population*

We used three repeated cross-sectional data from the NHMS conducted in 2006, 2011, and 2015 in Malaysia. The NHMS is a nationwide population-based survey conducted once every decade since 1986, and at 4-year intervals since 2011. The main aim of the NHMS is to monitor the prevalence and distribution of common diseases, health needs, and health expenditures. The survey was designed to represent all non-institutionalised individuals residing in Malaysia for at least four (NHMS 2006) or two (NHMS 2011 and NHMS 2015) weeks before data collection<sup>(18,19,29)</sup>. A similar sampling design and methodology was used in each NHMS. The survey was designed as a two-stage stratified cluster sampling proportionate to the population size. Malaysia was first stratified into states that were further stratified into rural and urban areas (<10,000 vs. ≥10,000 inhabitants), based on geostatistical census areas defined by the Department of Statistics, Malaysia (DOSM)<sup>(30)</sup>. The first stage of the sampling involved the random selection of enumeration blocks from the clusters of rural and urban areas separately. In the second stage, living quarters (buildings where people live) were randomly sampled from each selected enumeration block. All households and their members within the selected living quarters were included in the study. A larger sample was selected in 2006 to represent data at the national and state level, while the surveys in 2011 and 2015 were designed to provide data at the national level only. The response rates in living quarters for NHMS 2006, 2011, and 2015 were 90.0%, 88.2%, and 89.2%, and the individual participation rates were 94.6%, 93.0%, and 86.4%, respectively. In this study, data from 6- to 17-year-old children and adolescents were used.

### *Procedures and Measures*

The NHMS consists of face-to-face interviews, self-administrated questionnaires, and clinical measurements including height and weight. All questionnaires were structured, pre-tested and available as bilingual (English and Malay). A questionnaire manual was prepared to guide the data collection, done by trained research assistants. Information about children younger than 13 years of age was collected from their parents or guardians (by proxy), and information concerning individuals older than 13 years directly from them. The variables of this study were measured as described below.

#### (a) Anthropometry

Body weight was measured on an electronic digital weighing scale (Tanita Personal Scale) according to a standard procedure for children and adolescents<sup>(5)</sup>. Everyone was weighed twice,

barefoot, with minimum clothing; weight was recorded to the nearest 0.1 kg. A standard weight was supplied for each team for standardisation of the weighing scales. Height was measured twice based on the SECA Body Meter to the nearest 0.1 cm. The tools were validated and calibrated before use. The average of the two weight and height measurements, respectively, were used to calculate BMI (weight in kilograms divided by the square of height in meters,  $\text{kg/m}^2$ ). Two age- and sex-specific BMI criteria were used to define thinness, overweight, and obesity: (1) The IOTF 2012 reference is appropriate for children and adolescents aged 2-18 years, and provides BMI cut-offs that correspond to adult cut-offs of 18.5, 25.0, and 30.0  $\text{kg/m}^2$  at age 18<sup>(13)</sup>. We used a published SPSS code for the calculations<sup>(31)</sup>. (2) In the WHO 2007 criteria, thinness is defined as a BMI-for-age z-score lower than -2SD, overweight is defined as a BMI-for-age z-score greater than +1SD, and obesity is defined as a BMI-for-age z-score greater than +2SD, from the mean of the WHO reference population<sup>(14)</sup>. We used the WHO Reference 2007 SPSS macro package to calculate the BMI z-scores<sup>(32)</sup>. We excluded BMI z-scores greater than +5SD or less than -5SD. Age at the time of the interview, recorded in full years, was used when applying the IOTF and the WHO BMI criteria.

#### (b) Sociodemography

Age at the time of the interview, recorded in full years, was used in the analysis. Ages 6 to 9 years were classified as preadolescence, 10 to 13 years as early adolescence, and 14 to 17 years as mid-adolescence, based on adolescent growth and development<sup>(33)</sup>. Residential area was categorised into urban or rural according to the Department of Statistics Malaysia definition<sup>(30)</sup>. Ethnicity was based on major ethnicities in Malaysia: Malay, Chinese, Indian, *Other Bumiputeras* (which comprises indigenous people living in Peninsular and East Malaysia), and *Other* (including Malaysian minorities such as Sikh, Baba, Chitty, Eurasian, and non-citizens residing in Malaysia). Household size was classified based on the number of individuals living in the household into small (maximum three), medium (between four and six), and large (seven or more) households. Household income was based on the self-reported income of all individuals living in the household per month. For the analyses, household income was divided into quintiles (Q1 to Q5; Q1 representing the lowest and Q5 the highest income), separately for each survey year.

#### *Statistical Analysis*

The crude prevalence estimates of thinness, overweight, and obesity were calculated for each survey year, according to the IOTF and WHO BMI criteria. Complex sample analyses were conducted to account for the complex survey design. As BMI varies substantially by sex and age

during childhood and adolescence, all analyses were stratified by sex and the three age groups. Individuals with missing BMI values (3.7%, 6.2%, and 5.2% in 2006, 2011, and 2015, respectively) were excluded from all analyses. In addition, study subjects in the ethnicity category of *Other* were excluded from the analyses involving sociodemographic factors, due to the small number of participants in that category (approximately 3% of the total sample each survey year). The statistical significance of the time trends in the prevalence of thinness, overweight, and obesity across the study period was assessed by log-binomial regression, with year as a continuous variable and adjusted for age. The results from these analyses are presented as prevalence ratios (PR) (with 95% confidence intervals, 95% CI) that describe the relative change in the prevalence of thinness, overweight, and obesity, respectively, per year. Then, the prevalence of thinness and overweight was calculated for each survey year (using the IOTF criteria) separately in each subgroup of ethnicity, residential area, household size, and household income. We only used the IOTF criteria here, to avoid small numbers in some subgroups when defining thinness according to the WHO criteria. The statistical significance of the time trends in the prevalence of thinness and overweight within each sociodemographic subgroup was assessed by log-binomial regression, with year as a continuous variable, and adjusted for age. The results are presented as PR (with *p*-value) that describe the relative change in the prevalence of thinness and overweight per year in each sociodemographic subgroup. Finally, associations of individual sociodemographic factors (age group, ethnicity, residential area, household size, household income) with the prevalence of thinness (vs. normal weight) and overweight (vs. normal weight) in 2015 were studied. The statistical significance of these associations was assessed by log-Poisson regression, adjusted for age. The results of these analyses are presented as PRs (with 95% CI) that describe the ratio of the prevalence of thinness and overweight, respectively, in each sociodemographic subgroup in comparison to the respective reference group (in 2015). We used  $\log(\mu)$  as the link function in modeling the binary outcomes in log-binomial and log-Poisson regressions. All statistical tests were attained by using a two-tailed test, and an alpha level  $<0.05$  was used for statistical significance. Analyses were performed using StataSE 14 and IBM SPSS 25.0.



## Results

### *Study Population*

Data were available on 28,094 individuals aged 6 to 17 years, with the biggest sample (n=14,838) in 2006 and the smallest sample (n=6,262) in 2015 (**Table 1**). The proportion of individuals with Malay ethnicity increased, and the proportion of large households decreased somewhat during the study period. However, the distributions of sociodemographic characteristics were relatively similar in the three surveys, as follows: Malay ethnicity made up two-thirds of the sample, slightly over 50% of the individuals lived in urban areas, and around 60% came from medium-sized households.

### *Thinness*

The prevalence of thinness fell significantly from 22.7% (95% CI 21.8, 23.5) in 2006 to 18.7% (95% CI 17.3, 20.1) in 2015 among all individuals, from 22.1% (95% CI 21.0, 23.1) to 18.2% (95% CI 16.2, 20.3) in boys, and from 23.3% (95% CI 22.2, 24.2) to 19.1% (95% CI 17.4, 21.0) in girls according to the **IOTF criteria**. **Figure 1** shows the distribution of nutritional status, including the prevalence of thinness, across the study period according to sex and age. The decrease in the prevalence of thinness (IOTF criteria) over time was significant in all sex-age groups except for preadolescent boys (**Figure 2**, Panel A and Panel B). According to the **WHO criteria**, the prevalence of thinness decreased from 8.3% (95% CI 7.8, 8.8) to 6.7% (95% CI 5.8, 7.7) among all individuals, from 8.7% (95% CI 8.0, 9.4) to 7.4% (95% CI 6.0, 9.1) in boys, and from 7.8% (95% CI 7.1, 8.5) to 5.9% (95% CI 5.0, 7.1) in girls. Overall, there were no significant differences in the secular changes in thinness between boys and girls, whether defined by the IOTF or the WHO criteria.

Prevalence of thinness varied in the sociodemographic subgroups from 2006 to 2015 as shown in **Tables 2a (boys) and 2b (girls)**. The prevalence ratios, PRs (describing the relative change in the prevalence of thinness per year, adjusted for age, in subgroups of ethnicity, residential area, household size, and household income), were mostly below one, indicating a decrease in the prevalence of thinness from 2006 to 2015 in those subgroups; however, relatively few of the PRs were statistically significant. The only statistically significant interaction between survey year and a sociodemographic variable was observed for household size in early adolescent boys ( $p=0.014$ ). While the prevalence of thinness decreased over time among boys living in medium-sized and large households, it increased among boys living in small households.

In 2015, thinness was more common in preadolescence than in mid-adolescence (**Supplemental Table 1(a) and 1(b)**). In addition, a statistically significant association was observed between ethnicity and thinness among preadolescent girls; those with Indian ethnic origin had a higher prevalence of thinness, and those with Chinese origin had a lower prevalence of thinness, as compared to Malay (the strata-specific  $p$ -values with borderline significance).

### *Overweight and Obesity*

The prevalence of overweight (including obesity) increased significantly from 19.6% (95% CI 18.8, 20.4) in 2006 to 25.2% (95% CI 23.6, 27.0) in 2015 among all individuals, from 20.2% (95% CI 19.2, 21.3) to 26.0% (95% CI 23.8, 28.5) in boys, and from 18.9% (95% CI 17.9, 19.9) to 24.4% (95% CI 22.4, 26.5) in girls according to the **IOTF criteria**. Respectively, the prevalence of obesity increased from 7.2% (95% CI 6.7, 7.7) to 10.1% (95% CI 9.1, 11.2) among all individuals, from 7.3% (95% CI 6.7, 8.0) to 10.6% (95% CI 9.2, 12.2) in boys, and from 7.0% (95% CI 6.4, 7.7) to 9.5% (95% CI 8.3, 10.9) in girls. **Figure 1** shows the distribution of nutritional status, including the prevalence of overweight and obesity across the study period according to sex and age. The increases in the prevalence of overweight and obesity over time were statistically significant in all sex-age groups, except for mid-adolescent boys and, regarding obesity, in early adolescent girls (**Figure 2**, Panels C to F). Based on the **WHO criteria**, the prevalence of overweight (including obesity) increased from 23.4% (95% CI 22.5, 24.2) in 2006 to 29.2% (95% CI 27.5, 30.9) in 2015 in all individuals, and, respectively, from 24.9% (95% CI 23.8, 26.1) to 31.0% (95% CI 28.7, 33.3) in boys and from 21.7% (95% CI 20.7, 22.8) to 27.4% (95% CI 25.3, 29.4) in girls. Regarding obesity, the respective increases were from 10.4% (95% CI 9.8, 11.0) to 14.2% (95% CI 12.9, 15.5) in all individuals, from 11.9% (95% CI 11.1, 12.8) to 16.1% (95% CI 14.4, 18.1) in boys, and from 8.8% (95% CI 8.1, 9.6) to 12.1% (95% CI 10.6, 13.7) in girls. Overall, there was no significant difference in the secular changes in overweight and obesity between boys and girls, whether defined by the IOTF or WHO criteria.

**Tables 3a (boys) and 3b (girls)** show the prevalence of overweight (including obesity) in different sociodemographic subgroups from 2006 to 2015. The trends in the prevalence of overweight over time were mainly increasing ( $PR > 1$ ), and statistically significant in several subgroups, particularly in preadolescent boys and early adolescent girls. Statistically significant interactions between survey year and a sociodemographic variable were observed in three analyses, as follows: 1) Among mid-adolescent boys, overweight became more common in rural but not urban areas (interaction term  $p = 0.014$ ); 2) Among mid-adolescent girls, overweight became more common

among *Other* Bumiputeras and Chinese, while changes in the other groups were smaller and insignificant ( $p=0.020$ ); 3) Among early adolescent girls, overweight became more common in the highest and lowest quintiles of household income ( $p=0.009$ ), while changes in the other groups were smaller and insignificant.

In 2015, the following associations between the sociodemographic factors and overweight (including obesity) were statistically significant (**Supplemental Table 1(c) and 1(d)**): Early adolescent boys of Chinese ethnicity had a higher prevalence of overweight than Malay. Preadolescent boys from small households had a higher prevalence of overweight, and early adolescent girls from large households had a lower prevalence of overweight, as compared to medium-sized household residents.

Translating the prevalence estimates to absolute numbers, there were 1.1 million children and adolescents with thinness, 1.5 million children and adolescents with overweight (including obesity), and almost 600,000 children and adolescents with obesity in Malaysia in 2015, based on the IOTF classification. Using the WHO classification, the respective population estimates were 390,000, 1.7 million, and 820,000.

## **Discussion**

### *Summary of findings*

This is the first study describing secular trends and sociodemographic determinants of two forms of malnutrition (thinness, and overweight/obesity) simultaneously, among children and adolescents in Malaysia. As evidenced by this study, Malaysia is facing a double burden of malnutrition with increasing trends of overweight and obesity and a gradual decline in thinness prevalence from 2006 to 2015. Apart from a few exceptions, these secular changes were, in general, similar in the different sociodemographic subgroups (sex, age, ethnicity, place of residence, household size, and household income), and thinness and overweight co-existed in all the sociodemographic subgroups. The present study also highlights that different BMI criteria result in very different estimates of absolute numbers of thinness, but also of overweight and obesity, in Malaysian children and adolescents.

### *Comparison to previous studies*

Four national school-based studies were conducted in Malaysia between 2008 and 2017, in which the nutritional status of children and adolescents was assessed using the WHO BMI

classification<sup>(21–23,34)</sup>. The overall prevalence of thinness, overweight, and obesity in these previous studies were comparable to our results based on the WHO BMI criteria. Comparing to the prevalence estimates of thinness and excess weight by the NCD Risk Factor Collaboration among 5- to 19-year-olds in 2015<sup>(4)</sup>, thinness seems to be much more common, and overweight and obesity much less common, in some of the neighbouring countries than in Malaysia. On the other hand, 13- to 16-year-old adolescents in Brunei had a higher prevalence of overweight and obesity as compared to adolescents in Malaysia<sup>(35)</sup>. Secular trends of thinness and excess weight similar to those in our study have been observed in other developing countries in Southeast Asia<sup>(4,36–38)</sup> and China<sup>(39,40)</sup>, and among children and adolescents from South Asian<sup>(41)</sup> and African countries<sup>(42)</sup>.

Regarding the sociodemographic factors, the secular changes in the prevalence of thinness and overweight between 2006 and 2015 were, in general, similar in the different sociodemographic subgroups (sex, age, ethnicity, place of residence, household size, and household income). Looking at the most recent study year (2015), we observed a significantly higher prevalence of thinness in preadolescence (6-9 years) than in mid-adolescence (14-17 years). Because of rapid growth in children between the age of 5 years and pubertal age<sup>(43)</sup>, any imbalance in nutrient intake and energy expenditure during this period will increase the likelihood of thinness among the younger children. The distribution of thinness according to the different ethnic groups in our study is similar to that in previous national studies<sup>(21–23)</sup>. Regarding overweight, previous national studies reported roughly similar findings across the major ethnic groups, which is in line with our findings<sup>(21,22)</sup>. Differences in genetic factors, cultural traditions, and religious beliefs, as well as in socioeconomics, may influence dietary patterns, lifestyle habits, and weight perceptions—and, thereby, the weight status of ethnic groups<sup>(44,45)</sup>. Finally, we observed some differences in the prevalence of thinness and overweight according to household size, varying by sex and age. Previously, larger household size has been associated with a higher prevalence of thinness and a lower prevalence of excess weight among children from an urban area in Malaysia<sup>(27)</sup>. However, we are not aware of any national-level studies to compare our results. This signifies the need for further research.

### *Strengths and limitations*

A strength of our study is that we were able to use three large, nationally representative population-based data with comparable study designs and high response rates, and complex sample analysis to account for the disproportionate sampling of the population. Further, trained staff measured height and weight using standardised and calibrated equipment. Using BMI as a measure of adiposity can

be seen as a limitation<sup>(46)</sup>, although BMI is generally accepted as a screening measure of weight status in population studies<sup>(47)</sup>. The BMI cut-offs used for overweight and obesity<sup>(13,14)</sup> may have been too high in the sense that Asian populations have been observed to have higher health risks at lower BMI values<sup>(48)</sup>. Information on age, ethnicity, and household income was self-reported, which may cause some error, though we have no specific reason to suspect that such error would be systematic. We did not adjust household income for household size, and we did not include information on dietary intake, physical activity, or parental factors that could confound or explain associations between sociodemographic factors and weight status or explain the observed secular trends. Finally, although the sample sizes of the three NHMS surveys were large, some of the stratified analyses may have had insufficient statistical power. On the other hand, we made multiple comparisons within strata of sociodemographic variables, which increases the probability of receiving a statistically significant  $p$ -value in at least one stratum, even if the null hypothesis about an association would be true<sup>(49)</sup>. Because our study was descriptive and we used a frequentist analytical approach, we followed the suggestion by Sjölander and Vansteelandt and did not formally adjust the statistical analyses for multiple testing<sup>(49)</sup>.

### *Public health implications*

In the present study, the WHO population estimate for thinness was around 60% lower, and that the estimate for obesity was around 40% higher, than the respective estimates based on the IOTF classification. The discrepancies between the estimates based on the two BMI criteria are well-known by researchers in the field<sup>(50)</sup>. However, as prevalence figures inform public health policy decision-making, the importance of being clear about which criteria are used in each case cannot be over-emphasized.

Regardless of which BMI criteria are used, the double burden of malnutrition in terms of thinness and overweight and obesity exists among Malaysian children and adolescents. Considering the serious health consequences associated with undernutrition and overweight and obesity in children and adolescents, effective strategies incorporating double-duty actions to prevent co-existence of undernutrition with overweight and obesity are crucial. Based on our results, we recommend close monitoring of children's nutritional status and that public health interventions target families with children, from an early age.

## **Conclusion**

The present findings provide evidence that Malaysia is facing a double burden of malnutrition at national level among 6- to 17-year-old adolescents, with increasing trends of overweight and obesity and a persistent despite gradually decreasing trend of thinness across one decade, and affecting adolescents of different ethnic origins, from rural and urban areas, and from households of varying sizes and income levels. Further understanding is needed about sociodemographic, lifestyle-related, and environmental determinants of thinness and overweight among Malaysian children and adolescents and their families. Research is also warranted to investigate whether a double burden exists at other than the country level, such as the household level and the individual level.

**References**

1. World Health Organisation (2017) - The Double Burden of Malnutrition. Policy Brief. <https://www.who.int/nutrition/publications/doubleburdenmalnutrition-policybrief/en/> (accessed October 2019).
2. World Health Organization (2018) - Malnutrition. <https://www.who.int/news-room/fact-sheets/detail/malnutrition> (accessed October 2019).
3. Global Nutrition Report (2018) - Chapter 2. The Burden of Malnutrition. <https://globalnutritionreport.org/reports/global-nutrition-report-2018/burden-malnutrition/> (accessed October 2019).
4. NCD Risk Factor Collaboration (NCD-RisC). (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: A pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet* **390**, 2627–2642.
5. World Health Organisation (1995) *Physical status: The Use and Interpretation of Anthropometry*. World Health Organisation Expert Committee Report. WHO Technical Report Series No. 854. Geneva: WHO.
6. FAO, IFAD, UNICEF, WFP and WHO (2018) *The State of Food Security and Nutrition in the World. Building Climate Resilience for Food Security and Nutrition*. Rome: FAO.
7. Pulgarón ER (2013) Childhood Obesity: A review of increased risk for physical and psychological comorbidities. *Clin Ther* **35**, A18–A32.
8. World Health Organisation (2016) - Consideration of the evidence on childhood obesity for the Commission on Ending Childhood Obesity Report of the Ad hoc Working Group on Science and Evidence for Ending Childhood Obesity. <https://apps.who.int/iris/handle/10665/206549> (accessed October 2019).
9. Goh E Von, Azam-Ali S, McCullough F et al. The nutrition transition in Malaysia; key drivers and recommendations for improved health outcomes. *BMC Nutr*. Published online: 29 June 2020. doi: 10.1186/s40795-020-00348-5.
10. Popkin BM (2006). Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *Am J Clin Nutr* **84**, 289–298.
11. Global Nutrition Report (2020) Country Nutrition Profiles - Malaysia. <https://globalnutritionreport.org/resources/nutrition-profiles/asia/south-eastern-asia/malaysia/> (accessed Feb 2021).
12. Must A & Anderson SE (2006) Body mass index in children and adolescents: Considerations for population-based applications. *Int J Obes* **30**, 590–594.

13. Cole TJ & Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* **7**, 284–294.
14. de Onis M, Onyango AW, Borghi E et al. (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660–667.
15. Kuczmarski R, Ogden C, Guo S et al. (2002) 2000 CDC growth charts for the United States: Methods and development. *Vital Health Stat* **11**, **246**, 1-190.
16. Noor MI. (1992) Malnutrition and food consumption patterns in Malaysia. *Int J Food Sci Nutr* **43**, 69–78.
17. Khambalia AZ, Lim SS, Gill T et al. (2012) Prevalence and sociodemographic factors of malnutrition among children in Malaysia. *Food Nutr Bull* **33**, 31–42.
18. Institute for Public Health Malaysia. (2011) National Health and Morbidity Survey 2011.
19. Institute for Public Health Malaysia. (2015) National Health and Morbidity Survey 2015.
20. Institute for Public Health Malaysia. (2020) National Health and Morbidity Survey 2019.
21. Institute for Public Health Malaysia. (2013) National Health and Morbidity Survey: Malaysia School Based Nutrition Survey 2012.
22. Institute for Public Health Malaysia (2017) National Health and Morbidity Survey: Adolescent Nutrition Health Survey 2017.
23. Zainuddin AA, Selamat R, Baharudin A et al. (2013) Nutritional status of Malaysian primary school children aged 8-10 years: Findings from the 2008 National IDD Survey. *Malays J Nutr* **19**, 149–161.
24. Baharudin A, Zainuddin AA, Selamat R et al. (2013) Malnutrition among Malaysian adolescents: Findings from National Health and Morbidity Survey (NHMS) 2011. *Int J Public Heal Res* **3**, 282–289.
25. Poh BK, Wong JE, Norimah AK et al. (2016) Differences in body build in children of different ethnic groups and their impact on the prevalence of stunting, thinness, overweight, and obesity. *Food Nutr Bull* **37**, 3–13.
26. Naidu BM, Mahmud SZ, Ambak R et al. (2013) Overweight among primary school-age children in Malaysia. *Asia Pac J Clin Nutr* **22**, 408–415.
27. Ahmad A, Zulaily N, Shahril MR et al. Association between socioeconomic status and obesity among 12-year-old Malaysian adolescents. *PLoS One*. Published online: 25 July 2018. doi: 10.1371/journal.pone.0200577.
28. World Obesity Federation (2019) Atlas of Childhood Obesity. <https://www.worldobesity.org/nlsegmentation/global-atlas-on-childhood-obesity> (accessed October 2019).




29. Institute for Public Health Malaysia. (2008) National Health and Morbidity Survey 2006.
30. Department of Statistics Malaysia. (2010) Population Distribution and Basic Demographic Characteristics 2010.
31. SPSS Code for the "IOTF body mass index cut-offs for thinness, overweight and obesity". (2017) University of Luxemborg. Open Repository and Bibliography. <https://orbilu.uni.lu/handle/10993/31448> (accessed August 2018).
32. World Health Organisation - WHO Reference 2007 SPSS macro package. [https://www.who.int/growthref/tools/readme\\_spss.pdf?ua=1](https://www.who.int/growthref/tools/readme_spss.pdf?ua=1). (accessed August 2018).
33. Strasburger VC, Brown RT, Braverman PK et al. (2006) Adolescent growth and development. In *Adolescent Medicine : A Handbook for Primary Care*. pp. 3-9. Philadelphia. Lippincott Williams & Wilkins.
34. Poh BK, Ng BK, Siti Haslinda MD et al. (2013) Nutritional status and dietary intakes of children aged 6 months to 12 years: Findings of the Nutrition Survey of Malaysian Children (SEANUTS Malaysia). *Br J Nutr* **110**, Suppl.3, S21-S35.
35. Pengpid S & Peltzer K (2018) Overweight or obesity and related lifestyle and psychosocial factors among adolescents in Brunei Darussalam. *Int J Adolesc Med Health*. Published online: 8 Mar 2018. doi: 10.1515/ijamh-2018-0019.
36. Novianti Rachmi C, Li M, Baur LA. (2018) The double burden of malnutrition in Association of South East Asian Nations (ASEAN) countries: A comprehensive review of the literature. *Asia Pac J Clin Nutr* **27**, 736–755.
37. Rachmi CN, Agho KE, Li M et al. (2016) Stunting, underweight and overweight in children aged 2.0-4.9 years in Indonesia: Prevalence trends and associated risk factors. *PLoS One*. Published online: 11 May 2016. doi: 10.1371/journal.pone.0154756.
38. Kien VD, Lee HY, Nam YS et al. (2016) Trends in socioeconomic inequalities in child malnutrition in Vietnam: Findings from the Multiple Indicator Cluster Surveys, 2000-2011. *Glob Health Action*. Published online: 1 Mar 2016. doi: 10.3402/gha.v9.29263.
39. Song Y, Agardh A, Ma J et al. (2018) National trends in stunting, thinness and overweight among Chinese school-aged children, 1985–2014. *Int J Obes*. Published online: 18 June 2018. doi: 10.1038/s41366-018-0129-7.
40. Zhai L, Dong Y, Bai Y et al. (2017) Trends in obesity, overweight, and malnutrition among children and adolescents in Shenyang, China in 2010 and 2014: a multiple cross-sectional study. *BMC Public Health* **17**, 1–7.
41. Mistry SK & Puthussery S. (2015) Risk factors of overweight and obesity in childhood and adolescence in South Asian countries: A systematic review of the evidence. Published

online: 6 Mar 2015. doi: 10.1016/j.puhe.2014.12.004.


42. Muthuri SK, Francis CE, Wachira LJM et al. (2014) Evidence of an overweight/obesity transition among school-aged children and youth in Sub-Saharan Africa: A systematic review. *PLoS One*. Published online: 27 Mar 2014. doi: 10.1371/journal.pone.009284.
43. Rogol A, Clark P, Roemmich J (2000) Growth and pubertal development in children and adolescents: effects of diet and physical activity. *Am J Clin Nutr* **72**, Suppl., S521- S528.
44. Kasmini K, Idris MN, Fatimah A et al. (1997) Prevalence of overweight and obese school children aged between 7 to 16 years amongst the major 3 ethnic groups in Kuala Lumpur, Malaysia. *Asia Pac J Clin Nutr* **6**, 172–174.
45. Rezali FW, Chin YS, Shariff ZM et al. (2015) Evaluation of diet quality and its associated factors among adolescents in Kuala Lumpur, Malaysia. *Nutr Res Pract* **9**, 511–516.
46. Doak CM, Hoffman DJ, Norris SA et al. (2013) Is body mass index an appropriate proxy for body fat in children? *Glob Food Sec* **2**, 65–71.
47. Department of Health and Human Services Centers for Disease Control and Prevention - Body Mass Index: Considerations for Practitioners.  
<https://www.cdc.gov/obesity/downloads/BMIforPactitioners.pdf>. (accessed November 2019)
48. Hills AP, Mokhtar N, Brownie S et al. (2014) Childhood obesity in Asia: The value of accurate body composition methodology. *Asia Pac J Clin Nutr* **23**, 339–343.
49. Sjölander A & Vansteelandt S (2019) Frequentist versus Bayesian approaches to multiple testing. *Eur J Epidemiol* **34**, 809–821.
50. Partap U, Young EH, Allotey P et al. (2017) The use of different international references to assess child anthropometric status in a Malaysian population. *J Pediatr*. Published online: Nov 2017. doi: 10.1016/j.jpeds.2017.07.049.

## Legends for Figures:



**Figure 1.** Distribution of nutritional status based on BMI and defined by the IOTF<sup>(13)</sup> and WHO<sup>(14)</sup> criteria according to sex and age (preadolescent 6-9 years old, early adolescent 10-13 years old and mid-adolescent 14-17 years old) in the National Health and Morbidity Survey (NHMS) 2006, 2011 and 2015, Malaysia.

Thin = Thinness, OW = Overweight, OB = Obesity



**Figure 2.** Panel A-F. Prevalence Ratio (describes the relative change per year across the study period 2006-2015) of thinness, overweight (including obesity) and obesity based on the IOTF<sup>(13)</sup> and WHO<sup>(14)</sup> classifications of sex and age (preadolescent 6-9 years old, early adolescent 10-13 years old, and mid-adolescent 14-17 years old). National Health and Morbidity Survey (NHMS), Malaysia.

**Table 1.** Sociodemographic characteristics of participating individuals aged 6 to 17 years old in the National Health and Morbidity Survey (NHMS) 2006, 2011, and 2015, Malaysia.

Sociodemographic characteristics	NHMS 2006		NHMS 2011		NHMS 2015	
	n=14,838		n=6,994		n=6,262	
<b>Sex (%)</b>						
Boys	7550	50.9	3572	51.1	3082	49.2
Girls	7288	49.1	3422	48.9	3180	50.8
<b>Age Group (%)</b>						
Preadolescent (6-9 years)	5422	36.5	2403	34.4	2138	34.1
Early adolescent (10-13 years)	5166	34.8	2357	33.7	2081	33.2
Mid-adolescent (14-17 years)	4250	28.6	2234	31.9	2043	32.6
<b>Ethnicity (%)</b>						
Malay	9103	61.3	4457	63.7	4235	67.6
Chinese	2211	14.9	1048	15.0	766	12.2
Indian	1066	7.2	477	6.8	421	6.7
Other Bumiputeras <sup>a</sup>	2015	13.6	787	11.3	634	10.1
Other <sup>b</sup>	443	3.0	225	3.2	206	3.3
<b>Residential Area<sup>c</sup> (%)</b>						
Urban	8066	54.4	3916	56.0	3517	56.2
Rural	6772	45.6	3078	44.0	2745	43.8
<b>Household Size<sup>d</sup> (%)</b>						
Small	1455	9.8	588	8.4	833	13.3
Medium	8564	58.0	4291	61.6	4012	64.1
Large	4754	32.2	2088	30.0	1417	22.6
Missing Data <sup>*</sup>	65	0.4	27	0.4	-	-
<b>Household Income Quintile<sup>e</sup> (%)</b>						
Quintile 1 (Lowest Quintile)	2894	20.2	1410	20.2	1216	20.3
Quintile 2	3601	25.1	1405	20.1	1209	20.2
Quintile 3	2198	15.3	1594	22.8	1171	19.6
Quintile 4	2783	19.4	1188	17.0	1194	20.0
Quintile 5 (Highest Quintile)	2867	20.0	1397	20.0	1193	19.9
Missing Data <sup>*</sup>	495	3.3	-	-	279	4.5

<sup>a</sup>Other Bumiputeras comprised of indigenous individuals living in Peninsular and East Malaysia.

<sup>b</sup>Other ethnicities comprised of other minorities such as Sikh, Baba, Chitty, Eurasian, and non-citizens.

<sup>c</sup>Urban: Areas with population sizes  $\geq 10,000$ , Rural: Areas with population sizes  $< 10,000$ .

<sup>d</sup>Household Size was classified based on the number of individuals living in the household as small (maximum three), medium (between four and six) and large (seven or more).

<sup>e</sup>Household Income is the total average earnings of all individuals living in the household per month and was classified into Quintiles. Quintile 1 = the lowest income quintile, Quintile 5 = the highest income quintile.

\*Missing values are expressed separately from the valid percentage distribution.

NHMS 2006: Q1 ( $\leq$  RM 550.00), Q2 (RM 550.01-RM 1,000.00), Q3 (RM 1,000.01-RM 1,500.00), Q4 (RM 1,500.01-RM 2,510.00), Q5 ( $\geq$  RM 2,510.01); 1 USD = RM 3.78 in January 2006.

NHMS 2011: Q1 ( $\leq$  RM 900.00), Q2 (RM 901.00-RM 1,750.00), Q3 (RM 1,751.00-RM 3,000.00), Q4 (RM 3,001.00-RM 4,840.00), Q5 ( $\geq$  4,841.00); 1 USD = RM 3.06 in January 2011.

NHMS 2015: Q1 ( $\leq$  RM 1,300.00), Q2 (RM 1,301.00-RM 2,300.00), Q3 (RM 2,301.00-RM 3,500.00), Q4 (RM 3,501.00-RM 5,900.00), Q5 ( $\geq$  RM 5,901.00); 1 USD = RM 3.52 in January 2015.

**Table 2a.** Prevalence of thinness (IOTF criteria) in boys according to individual sociodemographic characteristics separated by age groups, survey year (2006, 2011, 2015), as well as Prevalence Ratio, PR (describes the relative change in the prevalence of thinness per year in each subgroup). National Health and Morbidity Survey (NHMS), Malaysia (Table continues onto next page).

Sociodemographic characteristic	Preadolescent					Early Adolescent					Mid-Adolescent				
	Survey year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>
	2006	2011	2015			2006	2011	2015			2006	2011	2015		
	Prevalence (%)					Prevalence (%)					Prevalence (%)				
<b>Thinness, total<sup>c</sup></b>	26.4	25.9	22.8	0.94 (0.85, 1.03)	0.164	18.5	18.2	14.9	0.90 (0.82, 0.99)	0.033	20.8	20.9	17.3	0.91 (0.83, 0.99)	0.046
<b>Ethnicity</b>					0.585					0.811					0.950
<b>Malay</b>	27.1	24.4	24.1	0.94 (0.83, 1.07)		18.9	17.0	15.2	0.90 (0.80, 1.01)		22.0	17.9	18.2	0.91 (0.80, 1.03)	
<b>Chinese</b>	17.1	20.0	18.8	1.05 (0.81, 1.36)		12.2	12.8	13.0	1.04 (0.75, 1.44)		15.4	18.7	11.8	0.87 (0.68, 1.11)	
<b>Indian</b>	38.6	48.5	29.4	0.91 (0.74, 1.11)		26.8	19.2	19.9	0.85 (0.62, 1.17)		28.8	31.7	27.2	0.98 (0.76, 1.27)	
<b>Other Bumiputeras<sup>d</sup></b>	25.8	30.7	17.8	0.86 (0.72, 1.03)		19.8	26.3	14.7	0.87 (0.67, 1.12)		15.6	33.4	15.4	0.94 (0.76, 1.17)	
<b>Residential Area<sup>e</sup></b>					0.079					0.999					0.662
<b>Urban</b>	24.5	26.6	23.4	0.98 (0.86, 1.11)		17.4	17.1	14.5	0.91 (0.80, 1.04)		20.0	21.0	17.2	0.92 (0.82, 1.04)	
<b>Rural</b>	28.7	24.2	20.3	0.85 (0.76, 0.95) **		20.1	18.7	16.6	0.91 (0.79, 1.05)		21.5	22.0	16.6	0.89 (0.77, 1.02)	
<b>Household Size<sup>f</sup></b>					0.923					0.014					0.930
<b>Small</b>	27.9	33.2	26.4	0.97 (0.79, 1.20)		10.8	7.5	20.8	1.51 (1.03, 2.20) *		14.2	17.9	13.7	0.96 (0.76, 1.21)	
<b>Medium</b>	25.8	24.2	22.3	0.94 (0.82, 1.07)		18.6	17.3	13.3	0.85 (0.75, 0.96) **		20.6	21.1	17.3	0.91 (0.81, 1.03)	

Accepted manuscript

<b>Large</b>	26. 4	28. 5	21. 3	0.93 (0.82, 1.05)		20. 5	20. 8	16. 9	0.92 (0.76, 1.10)		23. 3	24. 3	19. 0	0.91 (0.77, 1.09)
<b>Household Income<sup>g</sup></b>					0.83 9					0.95 4				0.12 9
<b>Quintile 1</b>	30. 0	28. 4	28. 7	0.99 (0.74, 1.33)		23. 9	20. 1	17. 0	0.84 (0.68, 1.04)		20. 9	25. 9	20. 0	0.97 (0.80, 1.17)
<b>Quintile 2</b>	29. 9	26. 9	25. 4	0.92 (0.78, 1.09)		19. 3	16. 2	16. 6	0.92 (0.74, 1.15)		25. 4	22. 3	13. 6	0.75 (0.61, 0.91) **
<b>Quintile 3</b>	30. 3	23. 9	20. 4	0.83 (0.70, 0.99) *		16. 0	22. 4	14. 4	0.94 (0.77, 1.15)		18. 9	25. 6	20. 5	1.02 (0.83, 1.24)
<b>Quintile 4</b>	22. 3	24. 9	17. 7	0.90 (0.76, 1.08)		17. 6	11. 1	14. 7	0.90 (0.68, 1.19)		21. 7	14. 2	15. 9	0.86 (0.69, 1.08)
<b>Quintile 5</b>	19. 4	25. 2	14. 8	0.90 (0.75, 1.08)		16. 3	17. 7	12. 9	0.89 (0.70, 1.14)		13. 7	18. 8	16. 2	1.05 (0.83, 1.33)



**Table 2b.** Prevalence of thinness (IOTF criteria) in girls according to individual sociodemographic characteristics separated by age groups, survey year (2006, 2011, 2015), as well as Prevalence Ratio, PR (describes the relative change in the prevalence of thinness per year in each subgroup). National Health and Morbidity Survey (NHMS), Malaysia (Table continued from previous page).

Sociodemographic characteristic	Predolescent					Early Adolescent					Mid-Adolescent				
	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>
	2006	2011	2015			2006	2011	2015			2006	2011	2015		
Prevalence (%)			Prevalence (%)			Prevalence (%)									
<b>Thinness, total<sup>c</sup></b>	30.0	28.9	25.3	0.92 (0.85, 0.99)	0.018	20.6	18.0	16.9	0.91 (0.83, 0.99)	0.034	18.7	21.6	15.8	0.91 (0.83, 0.99)	0.038
<b>Ethnicity</b>					0.284					0.781					0.899
<b>Malay</b>	30.3	28.8	27.2	0.95 (0.87, 1.03)		21.4	18.0	17.1	0.89 (0.80, 1.00)		18.3	20.3	15.5	0.91 (0.81, 1.03)	
<b>Chinese</b>	22.4	29.2	18.0	0.90 (0.73, 1.11)		16.9	15.4	14.7	0.93 (0.70, 1.24)		19.6	17.4	15.6	0.89 (0.69, 1.15)	
<b>Indian</b>	30.0	30.3	35.5	1.06 (0.82, 1.38)		23.5	19.1	14.9	0.81 (0.61, 1.08)		24.0	21.0	25.4	1.03 (0.76, 1.40)	
<b>Other Bumiputeras<sup>d</sup></b>	33.9	29.1	23.1	0.83 (0.69, 1.00)*		19.4	22.0	18.7	0.98 (0.77, 1.24)		15.1	34.4	16.0	0.93 (0.77, 1.12)	
<b>Residential Area<sup>e</sup></b>					0.403					0.758					0.302
<b>Urban</b>	29.1	31.3	25.9	0.94 (0.85, 1.03)		19.6	17.1	16.5	0.92 (0.81, 1.04)		19.9	21.3	16.3	0.89 (0.79, 1.01)	
<b>Rural</b>	30.6	24.0	24.8	0.88 (0.79, 0.98)*		22.2	20.2	17.7	0.89 (0.78, 1.02)		16.7	22.1	16.3	0.98 (0.86, 1.12)	
<b>Household Size<sup>f</sup></b>					0.118					0.594					0.146
<b>Small</b>	24.5	29.4	27.3	1.04 (0.81, 1.33)		16.0	16.5	18.1	1.04 (0.78, 1.38)		14.2	18.3	19.4	1.16 (0.89, 1.49)	
<b>Medium</b>	28.5	27.8	26.6	0.96 (0.87, 1.05)		20.0	17.9	15.9	0.90 (0.79, 1.02)		19.7	21.7	16.3	0.89 (0.79, 0.99)*	

<b>Large</b>	33. 1	31. 1	21. 9	0.83 (0.73, 0.95)**		23. 3	19. 1	18. 5	0.88 (0.75, 1.03)		19. 1	22. 5	13. 7	0.89 (0.72, 1.10)
<b>Household Income<sup>g</sup></b>					0.17 6					0.87 1				0.46 7
<b>Quintile 1</b>	35. 3	23. 2	26. 7	0.83 (0.68, 1.00)		20. 7	20. 5	17. 3	0.92 (0.77, 1.09)		20. 7	17. 9	12. 9	0.78 (0.62, 0.99)*
<b>Quintile 2</b>	31. 6	30. 4	22. 1	0.86 (0.72, 1.02)		24. 7	22. 2	18. 6	0.87 (0.70, 1.08)		19. 1	21. 1	18. 7	0.99 (0.82, 1.19)
<b>Quintile 3</b>	28. 0	34. 6	33. 4	1.08 (0.92, 1.26)		19. 3	20. 8	13. 8	0.85 (0.69, 1.04)		15. 1	23. 6	16. 2	0.98 (0.80, 1.20)
<b>Quintile 4</b>	26. 5	25. 7	25. 2	0.98 (0.82, 1.16)		19. 1	10. 6	18. 4	0.99 (0.75, 1.30)		17. 0	18. 2	17. 6	1.02 (0.81, 1.28)
<b>Quintile 5</b>	27. 0	29. 4	21. 1	0.88 (0.76, 1.02)		19. 5	16. 3	13. 1	0.84 (0.67, 1.04)		20. 8	25. 4	16. 7	0.87 (0.70, 1.07)

Age Groups: Preadolescent 6-9 years old, Early adolescent 10-13 years old and Mid-adolescent 14-17 years old.

<sup>a</sup>95% confidence interval for the prevalence ratio (PR).

<sup>b</sup>Interaction test between sociodemographic variable and survey year. Significance test for each sociodemographic subgroup: \* significance at <5% level, \*\* significance at <1% level, \*\*\* significance at <0.1% level.

<sup>c</sup>Thinness was classified according to the IOTF classification(13)

<sup>d</sup>Other Bumiputeras comprised of indigenous individuals living in Peninsular and East Malaysia.

<sup>e</sup>Urban: Areas with population sizes  $\geq 10,000$ , Rural: Areas with population sizes <10,000.

<sup>f</sup>Household Size was classified based on the number of individuals living in the household into small (maximum three), medium (between four and six) and large (seven or more).

<sup>g</sup>Household Income is the total average earnings of all individuals living in the household per month and was classified into Quintile. Quintile 1 = the lowest income quintile, Quintile 5 = the highest income quintile.

**Table 3a.** Prevalence of overweight (including obesity; IOTF criteria) in boys according to individual sociodemographic characteristics separated by age group, survey year (2006, 2011, 2015), as well as Prevalence Ratio, PR (describes the relative change in the prevalence of overweight individuals per year in each subgroup). National Health and Morbidity Survey (NHMS), Malaysia (Table continues onto next page).

Sociodemographic characteristic	Preadolescent					Early Adolescent					Mid-Adolescent				
	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>
	2006	2011	2015			2006	2011	2015			2006	2011	2015		
	Prevalence (%)					Prevalence (%)					Prevalence (%)				
<b>Overweight, total<sup>c</sup></b>	16.6	19.9	25.8	1.25 (1.14, 1.37)	<0.001	24.7	27.6	29.1	1.08 (1.00, 1.17)	0.043	19.6	22.3	23.3	1.09 (0.99, 1.19)	0.081
<b>Ethnicity</b>					0.151					0.199					0.369
<b>Malay</b>	14.5	20.4	24.3	1.29 (1.15, 1.44)***		24.3	27.8	26.9	1.05 (0.95, 1.16)		19.3	24.1	23.2	1.08 (0.97, 1.21)	
<b>Chinese</b>	27.5	26.9	30.1	1.03 (0.85, 1.26)		30.6	33.4	43.0	1.21 (1.02, 1.43)*		24.3	20.8	28.1	1.11 (0.87, 1.42)	
<b>Indian</b>	21.8	12.1	34.9	1.32 (0.96, 1.80)		29.7	43.2	23.9	0.94 (0.76, 1.17)		21.3	32.5	18.4	0.94 (0.72, 1.21)	
<b>Other Bumiputeras<sup>d</sup></b>	11.8	13.5	25.6	1.55 (1.12, 2.15)**		16.4	18.0	24.4	1.24 (0.94, 1.64)		13.7	14.7	25.0	1.44 (0.99, 2.08)	
<b>Residential Area<sup>e</sup></b>					0.068					0.137					0.014
<b>Urban</b>	19.2	21.1	26.3	1.18 (1.05, 1.32)**		28.4	32.2	30.4	1.03 (0.94, 1.13)		21.8	25.1	23.1	1.02 (0.91, 1.14)	
<b>Rural</b>	13.1	18.0	26.0	1.40 (1.20, 1.62)***		19.4	20.5	26.0	1.16 (1.02, 1.31)*		16.7	17.4	26.9	1.30 (1.11, 1.51)**	
<b>Household Size<sup>f</sup></b>					0.467					0.723					0.819

<b>Small</b>	22. 3	18. 2	38. 2	1.38 (1.08, 1.76)*		30. 0	38. 8	37. 0	1.09 (0.90, 1.31)		22. 4	20. 3	23. 1	1.03 (0.82, 1.30)
<b>Medium</b>	17. 5	21. 4	25. 2	1.19 (1.06, 1.34)**		26. 6	29. 9	29. 2	1.04 (0.95, 1.14)		20. 9	23. 6	25. 3	1.10 (0.97, 1.24)
<b>Large</b>	13. 9	17. 6	23. 3	1.30 (1.09, 1.55)**		19. 9	22. 8	25. 0	1.12 (0.95, 1.33)		16. 5	20. 6	21. 5	1.14 (0.94, 1.37)
<b>Household Income<sup>g</sup></b>					0.11 9					0.20 3				0.32 8
<b>Quintile 1</b>	7.3	16. 0	23. 8	1.72 (1.30, 2.28)***		15. 0	27. 7	27. 1	1.28 (1.06, 1.56)*		16. 2	16. 1	18. 9	1.09 (0.84, 1.40)
<b>Quintile 2</b>	13. 3	16. 1	19. 2	1.22 (0.97, 1.52)		22. 1	22. 3	30. 8	1.19 (0.99, 1.42)		15. 9	21. 4	22. 6	1.18 (0.95, 1.46)
<b>Quintile 3</b>	18. 1	21. 2	29. 3	1.27 (1.04, 1.55)*		25. 2	26. 2	25. 3	1.00 (0.82, 1.22)		17. 9	21. 0	26. 2	1.21 (0.95, 1.55)
<b>Quintile 4</b>	21. 0	25. 1	29. 5	1.19 (0.99, 1.43)		26. 2	30. 9	26. 0	1.00 (0.84, 1.18)		19. 0	24. 8	27. 1	1.18 (0.96, 1.44)
<b>Quintile 5</b>	23. 3	22. 3	28. 4	1.11 (0.92, 1.33)		34. 2	33. 3	35. 8	1.03 (0.89, 1.19)		28. 9	28. 4	25. 9	0.94 (0.79, 1.13)

**Table 3b.** Prevalence of overweight (including obesity; IOTF criteria) in girls according to individual sociodemographic characteristics, separated by age group, survey year (2006, 2011, 2015), as well as Prevalence Ratio, PR (describes the relative change in the prevalence of overweight per year in each subgroup). National Health and Morbidity Survey (NHMS), Malaysia (Table continued from previous page).

Sociodemographic characteristic	Preadolescent			Early Adolescent					Mid-Adolescent						
	Survey Year			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Prevalence Estimates (%)			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>	Prevalence Estimates (%)			PR (95%CI) <sup>a</sup>	<i>p</i> -value <sup>b</sup>
	2006	2011	2015			2006	2011	2015			2006	2011	2015		
	Prevalence (%)			Prevalence (%)					Prevalence (%)						
<b>Overweight, total<sup>c</sup></b>	16.2	17.4	21.1	1.16 (1.06, 1.28)	0.002	20.4	23.5	27.9	1.17 (1.09, 1.27)	<0.001	20.2	18.1	24.0	1.11 (1.01, 1.22)	0.028
<b>Ethnicity</b>					0.490					0.410					0.020
<b>Malay</b>	14.7	20.9	21.5	1.21 (1.08, 1.35)**		20.8	28.1	31.0	1.21 (1.11, 1.32)***		22.5	20.1	26.2	1.09 (0.98, 1.22)	
<b>Chinese</b>	20.7	16.9	19.4	0.98 (0.74, 1.29)		22.1	16.5	22.5	1.03 (0.79, 1.34)		14.5	11.8	22.5	1.34 (1.00, 1.80)	
<b>Indian</b>	27.6	18.8	30.0	1.04 (0.72, 1.50)		23.5	29.6	26.4	1.05 (0.83, 1.33)		20.8	36.1	12.6	0.84 (0.67, 1.06)	
<b>Other Bumiputeras<sup>d</sup></b>	13.8	7.2	18.6	1.23 (0.85, 1.79)		15.2	14.0	24.4	1.32 (0.99, 1.76)		15.4	12.7	27.1	1.49 (1.05, 2.10)*	
<b>Residential Area<sup>e</sup></b>					0.777					0.103					0.201

<b>Urban</b>	18.1	17.7	22.7	1.16 (1.02, 1.32)*	21.9	24.4	27.7	1.13 (1.02, 1.25)*	21.6	18.7	24.0	1.08 (0.96, 1.22)	
<b>Rural</b>	14.2	18.4	17.5	1.13 (0.99, 1.29)	18.6	22.8	30.3	1.28 (1.14, 1.43)***	18.0	18.0	26.2	1.23 (1.06, 1.42)**	
<b>Household Size<sup>f</sup></b>					0.27				0.90			0.75	
					1				6			6	
<b>Small</b>	17.4	22.8	24.7	1.21 (0.88, 1.67)	26.2	29.3	36.2	1.17 (0.97, 1.42)	24.5	14.0	24.1	1.02 (0.79, 1.33)	
<b>Medium</b>	18.3	19.3	20.8	1.09 (0.97, 1.21)	22.1	24.5	29.7	1.17 (1.06, 1.29)**	20.7	20.3	25.9	1.14 (1.01, 1.29)*	
<b>Large</b>	12.6	14.5	21.1	1.32 (1.05, 1.66)*	16.2	20.8	20.5	1.13 (0.96, 1.32)	17.1	16.3	20.9	1.10 (0.91, 1.34)	
<b>Household Income<sup>g</sup></b>					0.23				0.009			0.38	
					2							1	
<b>Quintile 1</b>	12.1	15.2	17.3	1.24 (0.98, 1.58)	13.4	19.9	32.4	1.57 (1.30, 1.91)***	15.9	16.1	23.4	1.28 (0.97, 1.70)	
<b>Quintile 2</b>	15.4	15.4	29.4	1.42 (1.13, 1.78)**	18.4	20.5	25.0	1.18 (0.97, 1.43)	0.03	18.6	21.4	27.0	1.22 (1.03, 1.44)*
<b>Quintile 3</b>	15.9	16.5	15.3	0.98 (0.77, 1.26)	27.1	20.1	24.5	0.96 (0.78, 1.18)	0.00	23.8	21.0	22.4	0.96 (0.79, 1.17)
<b>Quintile 4</b>	20.5	24.6	24.0	1.08 (0.88, 1.31)	25.2	29.9	31.1	1.11 (0.94, 1.31)	0.00	24.0	24.8	29.1	1.16 (0.92, 1.45)
<b>Quintile 5</b>	18.0	18.3	21.8	1.18 (0.96, 1.46)	20.9	29.6	31.7	1.23 (1.06, 1.42)**	0.04	19.1	28.4	24.0	1.15 (0.91, 1.46)

Age Groups: Preadolescent 6-9 years old, Early adolescent 10-13 years old and Mid-adolescent 14-17 years old.

<sup>a</sup>95% confidence interval for the prevalence ratio (PR).

<sup>b</sup>Interaction test between sociodemographic variable and survey year. Significance test for each sociodemographic subgroup: \* significance at <5% level, \*\* significance at <1% level, \*\*\* significance at <0.1% level.

<sup>c</sup>Overweight (including obesity) were classified according to the IOTF classification(13).

<sup>d</sup>Other Bumiputeras comprised of indigenous individuals living in Peninsular and East Malaysia.

<sup>e</sup>Urban: Areas with population sizes  $\geq 10,000$ , Rural: Areas with population sizes  $< 10,000$ .

<sup>f</sup>Household Size was classified based on the number of individuals living in the household into small (maximum three), medium (between four and six) and large (seven or more).

<sup>g</sup>Household Income is the total average earnings of all individuals living in the household per month and was classified into Quintile. Quintile 1 = the lowest income quintile, Quintile 5 = the highest income quintile