

Implementation, Adoption and Use of the Kanta Services in Finland 2010–2022

Vesa JORMANAINEN^{a,b,1}, Tuulikki VEKHO^a, Marina LINDGREN^c,
Ilmo KESKIMÄKI^{a,d} and Minna KAILA^b

^a*Finnish Institute for Health and Welfare (THL)*

^b*University of Helsinki, Faculty of Medicine, Department of Public Health*

^c*Social Insurance Institution of Finland (Kela)*

^d*Tampere University, Faculty of Social Sciences*

ORCID ID: Vesa Jormanainen <https://orcid.org/0000-0002-1906-8043>

Abstract. Nationwide implementation and adoption of the Prescription Centre and the Patient Data Repository services required 5.5 years since May 2010 in Finland. The Clinical Adoption Meta-Model (CAMM) was applied in the post-deployment assessment of the Kanta Services in its four dimensions (availability, use, behavior, clinical outcomes) over time. The CAMM results on the national level in this study suggest ‘Adoption with Benefits’ as the most appropriate CAMM archetype.

Keywords. Implementation, adoption, use, Kanta Services, Finland, Clinical Adoption Meta-Model

1. Introduction

Healthcare systems are currently facing several challenges, such as population aging, economic constraints and rapid technological change, which call for comprehensive reforms. Internationally a prevailing solution offered for these challenges is to deploy large-scale information and communication technology systems [1,2]. However, adopting new national health information (HIS) or health information exchange (HIE) systems are often challenging ventures: a large-scale, complex and costly endeavor, taking many years to develop and build, involving multiple public and private stakeholders and that has impact on millions of people [3–7].

HIS integrates the data collection, processing, reporting, and use of the information necessary for improving health service effectiveness and efficiency through better management at all levels of health services [8,9]. HIE essentially includes the electronic transfer of patient data and health information between healthcare service providers or institutions [10–13].

Current literature is still dominated by reports of single organizations and the punctuality of their HIS implementation [14–17]. Moreover, most HIS/HIE system and healthcare reforms are not properly followed up and their outcomes are rarely evaluated [2,17].

¹ Corresponding Author: Dr. Vesa Jormanainen, Finnish Institute for Health and Welfare, P.O. Box 30, 00271 Helsinki, Finland; E-mail: vesa.jormanainen[at]gov.fi.

This research aimed to assess the implementation and adoption of the *Act on Electronic Prescription* (61/2007) and the *Act on Processing Customer Data in Health and Social Care* (159/2007). The assessment utilized log-based register data on use of the national healthcare Kanta Services since May 2010 in Finland.

2. Methods

The Kanta Services is the name of Finland's centralized, shared, and integrated electronic data system services introduced in phases since May 20, 2010 [18–20]. The implementation processes and adoption efforts in healthcare were prospectively followed-up by utilizing records accumulated in the Finnish Institute for Health and Welfare's (THL) national operative coordinating unit from May 2010 to December 2016. Use of the Kanta Services from May 2010 to December 2022 was followed-up by rigorously utilizing log-based register data on descriptive performance indicators' time series provided by the Social Insurance Institution of Finland (Kela).

We used Clinical Adoption Meta-Model's (CAMM) four dimensions (availability, use, behavior, clinical outcomes) over the fifth dimension of time [21,22]. According to the CAMM, a data system will become available, it will be used which in turn will lead to changes in behavior, that will produce clinical or health outcomes over time.

In the 'No Deployment' archetype of the CAMM, the HIS fails to reach end-users. In the 'Low Adoption' archetype, the HIS is deployed and available, but with minimal or rapidly declining use. In the 'Adoption without Benefit' archetype, a HIS is deployed, available and used by end-users, but it fails to achieve the intended behavior changes or the expected outcomes. In turn, the 'Behavior Change without Outcome Benefit' archetype occurs when an adopted HIS produces the expected behavior change but fails to produce the expected outcomes. The 'Adoption with Benefits' archetype is characterized by a clear progression of HIS availability that leads to ongoing HIS use, which then causes observable changes in clinical and health behavior that, in turn, result in improvements in measured outcomes. In the 'Benefit without Use' archetype, the expected behavior changes and/or outcomes occur without HIS use.

3. Results

3.1. Availability of the nationwide healthcare Kanta Services

The first certificate of interoperability acceptance for healthcare and pharmacy data systems for the electronic Prescription Centre services was granted in May 18, 2010, and in public healthcare, it took 859 days to reach the 50% national population coverage and 1,258 days for the 100% point.

The first certificate of interoperability acceptance for healthcare data systems for the electronic Patient Data Repository was granted in October 28, 2013, and in public healthcare, it took 376 days to reach the 50% point and 760 days for the 100% point.

The mean implementation and adoption time varied considerably (Table 1).

Table 1. Mean number of years from the start of the first implementation and adoption project to the start of the last implementation and adoption project (min. and max. times in years in brackets) of the healthcare Kanta Services in Finland in 2010–2015. PHCs refers to public primary healthcare centres.

Kanta Services to be implemented	Hospital districts Mean (min.–max.)	University-hospital-specific catchment areas Mean (min.–max.)	National Mean
Prescription Centre			
Pharmacies, years	0.6 (0.1–0.2)	1.3 (0.9–2.0)	2.4
PHCs, years	0.6 (<0.01–2.0)	2.0 (0.9–3.4)	3.4
Pharmacies+PHCs, years	1.1 (0.02–2.4)	2.3 (1.8–3.4)	3.4
Patient Data Repository			
PHCs, years	0.6 (<0.01–1.6)	1.5 (1.1–2.0)	2.0
Both Kanta Services			
PHCs, years	3.0 (1.6–5.1)	4.3 (3.6–5.1)	5.5
Pharmacies+PHCs, years	3.6 (2.5–4.3)	4.6 (4.4–5.1)	5.5

3.2. Use of the national healthcare Kanta Services in Finland since May 2010

Citizen users have accessed the web based My Kanta Pages since May 2010, sent electronic repeat prescription (eP) renewal requests to healthcare organizations since November 2015, and have made visits on behalf of their children since October 2016 (Table 2). In addition, citizen users have recorded consents, consent restrictions, organ donation testaments and living wills into the Patient Data Management Service.

Table 2. Performance indicators by healthcare Kanta Services in 2022 and cumulatively 2010–2022 in Finland (numbers in millions). ePs refers to electronic prescriptions.

Kanta Service	2022	2010–2022
Performance indicator	Millions	Millions
My Kanta Pages		
Sign-ins	37.2	179
Repeat eP renewal requests	3.4	18.7
Visits on behalf of children	2.5	12.2
Prescription Centre		
ePs	28.2	274
Dispensation documents	76.7	580
Patient Data Repository		
Documents	477	3174
Service events	230	1581
Persons	6.5	6.5
Information notices	0.24	7.9
Consents	0.24	4.8
Consent restrictions	0.12	0.16
Organ testaments	0.05	0.88
Living wills	0.04	0.28

Healthcare professionals (community pharmacy professionals since 2017) have recorded ePs, and community pharmacy professionals medication dispensing documents into the Prescription Centre since May 2010 (Table 2). Healthcare professionals have recorded documents related to service events into the Patient Data Repository since November 2013, and information notices into the Patient Data Management Service.

3.3. Behavior of healthcare professionals

Healthcare professionals quickly learned to issue and use ePs, and pharmaceutical professionals in community pharmacies learned quickly to record paper-based and

telephone prescriptions into the Prescription Centre after the ‘big bang’ change in January 2017, when ePs became mandatory in Finland.

The Kela launched free-of-charge Kelain web-based service in September 2016 to support the start of mandatory ePs in January 2017. The number of registered healthcare professional Kelain service users rose rapidly to 18,000 and the number of ePs issued via Kelain service rose to 0.301 million in 2018.

3.4. CAMM archetypes and the Kanta Services in Finland

The Kanta Services already reached end-users in a clinical setting in May 2010, and, thus, escaped the CAMM ‘No Deployment’ archetype. The Kanta Services’ implementation and adoption matured and escaped the archetypes ‘Low Adoption’, ‘Adoption without Benefit’, ‘Behavioral Change without Outcome Benefit’ and the ‘Benefit without Use’. Based on the results of this study, the most accurate CAMM archetype is ‘Adoption with Benefits’.

4. Discussion and conclusions

The results of this research suggest that it is possible to implement and adopt two large-scale national HIS/HIE in 5.5 years covering all public primary healthcare centres, community pharmacies and hospitals together with most private healthcare service providers in a country with 5.5 million inhabitants.

An implementation strategy combining top-down and bottom-up approaches [2,11,23–26] employed in Finland proved an appropriate strategy for the implementation and adoption of the two healthcare Kanta Services.

The Prescription Centre services were implemented and adopted first and thereafter the Patient Data Repository services [18,19]. Public healthcare providers implemented and adopted the Kanta Services first and thereafter private healthcare service providers. The initial hospital district based implementation strategy for the Prescription Centre services informed the strategy to a certified Kanta Services compatible HIS/HIE strategy for the Patient Data Repository services. The THL’s operative coordination unit – with appropriate legal mandate in the permanent legislation – supported the large-scale implementation processes of the two healthcare Kanta Services in close cooperation with the Kela.

The results from a previous study showed that the implementation of HIS/HIE systems in Finland has successfully passed several milestones in terms of the CAMM archetypes [27]. The case of Kanta Services in Finland suggested that the healthcare-specific CAMM theoretical construct can be used on the national level HIS implementation assessments. Results of this study repeat the observation. In addition, results of this study suggest that the most appropriate CAMM archetype is ‘Adoption with Benefits’.

Even rigorously utilizing log-based register data of the Kanta Services, however, must accompany parallel and independently conducted HIS/HIE research (e.g. general public and/or professionals’ surveys) in order to gain insights and experiences of the whole [28]. Together, ‘log-based data’ and the ‘research-based in-depth data’ arms likely complement each other, and form a rich database for further elaboration of outcomes, benefits and harms, and other effects to inform development of new functionalities and better user interfaces for both citizen and professional users.

References

- [1] Ludwig DA, Doucette J. Adopting electronic medical records in primary care: lessons learned from health information systems implementation experience in seven countries. *Int J Med Inform* 2009;78:22–31..
- [2] Coiera E. Building a national health IT system from the Middle Out. *JAMIA* 2009;16:271–273.
- [3] Deutsch E, Duftschmid G, Dorda W. Critical areas of national electronic health record programs: is our focus correct? *Int J Med Inform* 2010;79:211–222.
- [4] Flyvbjerg B. What you should know about mega-projects and why: an overview. *Project Manage J* 2014;45:6–19.
- [5] Ross J, Stevenson F, Lau F, et al. Factors that influence the implementation of e-health: a systematic review of reviews (an update). *Implement Sci* 2016;11:146.
- [6] Fennelly O, Cunningham C, Grogan L, et al. Successfully implementing a national electronic health record: a rapid umbrella review. *Int J Med Inform* 2020;144:104281.
- [7] Ellingsen G, Hertzum M, Melby L. The tension between national and local concerns in preparing for large-scale generic systems in healthcare. *Computer Supported Cooperative Work* 2022; published online.
- [8] Sligo J, Gould R, Roberts V, et al. A literature review for large-scale health information system project planning, implementation and evaluation. *Int J Med Inform* 2017;97:86–97.
- [9] Lippeweld T, Sauerborn R, Bodart C (Eds.). *Design and implementation of health information systems*. Geneva: World Health Organization; 2000.
- [10] Finn N. Health information exchange: a stepping stone toward continuity of care and participatory medicine. *J Particip Med* 2011;3:e47.
- [11] Bowden T, Coiera E. Comparing New Zealand’s ‘middle out’ health information technology strategy with other OECD nations. *Int J Med Inform* 2013;82:e87–e95.
- [12] Adler-Milstein J, Ronchi E, Cohen GR, et al. Benchmarking health IT among OECD countries: better data for better policy. *JAMIA* 2014;21:111–116.
- [13] Esmaeilzadeh P, Sambasivan M. Health information exchange (HIE): a literature review, assimilation pattern and a proposed classification for a new policy approach. *J Biomed Inform* 2016;64:74–86.
- [14] Robertson A, Cresswell K, Takian A, et al. Implementation and adoption of nationwide electronic health records in secondary care in England: qualitative analysis of interim results from a prospective national evaluation. *BMJ* 2010;341:c4564.
- [15] Klecun E, Zhou Y, Kankanhalli A, et al. The dynamics of institutional pressures and stakeholder behavior in national electronic health record implementations: a tale of two countries. *J Inform Technol* 2019;34:292–332.
- [16] Luz R, Carneiro Mussi C, Dutra A, et al. Implementation of large-scale health information systems. *Revista de Gestao* 2021;28:106–132.
- [17] Rigby M. *Evaluation: 16 powerful reasons why not to do it and 6 over-riding imperatives*. MEDINFO 2001. IOT Press 2001;1198–1202.
- [18] Jormanainen V. Large-scale implementation and adoption of the Finnish national Kanta services in 2010–2017: a prospective, longitudinal, indicator-based study. *Finnish J eHealth eWelfare* 2018;10:381–395.
- [19] Jormanainen V, Parhiala K, Niemi A, et al. Half of the Finnish population accessed their own data: comprehensive access to personal information online is a corner-stone of digital revolution in Finnish health and social care. *Finnish J eHealth eWelfare* 2019;11:298–310.
- [20] Kanta Services. [Web pages in English](#). Accessed January 5, 2022.
- [21] Price M, Lau F. The clinical adoption meta-model: a temporal meta-model describing the clinical adoption of health information systems. *BMC Med Inform Decision Mak* 2014;14:43.
- [22] Price M. *Clinical Adoption Meta-Model*. In: Lau F, Kuziemsky C (Eds). *Handbook of eHealth evaluation: an evidence-based approach*. Victoria (B.C., Canada): University of Victoria 2016;77–91.
- [23] Greenhalgh T, Bowden T. Moving beyond the cathedral model of IT development. *BMJ* 2011;343:d6480.
- [24] Morrison Z, Robertson A, Cresswell K, et al. Understanding contrasting approaches to nationwide implementation of electronic health record systems: England, the USA and Australia. *J Healthcare Engin* 2011;2:25–41.
- [25] Eason K, Dent M, Waterson P, et al. Bottom-up and middle-out approaches to electronic patient information systems: a focus on healthcare pathways. *Inform Prim Care* 2012;20:51–56.
- [26] Tsai CH, Koch S. Towards a framework for national eHealth evaluation and monitoring: a combined top-down and bottom-up approach using Sweden as example. *Stud Health Technol Inform* 2019;264:954–958.
- [27] Jormanainen V, Reponen J. CAF and CAMM analyses on the first 10 years of national Kanta services in Finland. *Finnish J eHealth eWelfare* 2020;12:302–315.
- [28] Vehko T, Ruotsalainen S, Hyppönen H (Eds.). *E-health and e-welfare of Finland: check point 2018*. THL Report 7/2019. Helsinki: National Institute for Health and Welfare (THL); 2018.