

PLA2G7 associates with hormone receptor negativity in clinical breast cancer samples and regulates epithelial–mesenchymal transition in cultured breast cancer cells

Laura Lehtinen,^{1*} Paula Vainio,¹ Harriet Wikman,² Heini Huhtala,³ Volkmar Mueller,⁴ Anne Kallioniemi,⁵ Klaus Pantel,² Pauliina Kronqvist,¹ Olli Kallioniemi,^{6,8} Olli Carpèn^{1,9} and Kristiina Iljin⁷

¹ Department of Pathology, Turku University and Turku University Hospital, Turku, Finland

² Institute of Tumour Biology, Centre of Experimental Medicine, University Medical Centre Hamburg-Eppendorf, Germany

³ School of Health Sciences, University of Tampere, Tampere, Finland

⁴ Department of Gynecology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

⁵ BioMediTech, University of Tampere, Tampere, Finland

⁶ FIMM, Institute for Molecular Medicine Finland, University of Helsinki, Finland

⁷ VTT Technical Research Centre of Finland, Espoo, Finland

⁸ Present address: Department of Oncology-Pathology, Science for Life Laboratory, Karolinska Institutet, Solna, Sweden

⁹ Present address: Department of Pathology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

*Correspondence to: Laura Lehtinen, Department of Pathology, Turku University and Turku University Hospital, Turku, Finland.

E-mail: lauleh@utu.fi

Breast cancer is the leading cause of cancer-related deaths in women due to distinct cancer subtypes associated with early recurrence and aggressive metastatic progression. High lipoprotein-associated phospholipase A2 (*PLA2G7*) expression has previously been associated with aggressive disease and metastasis in prostate cancer. Here, we explore the expression pattern and functional role of *PLA2G7* in breast cancer. First, a bioinformatic analysis of genome-wide gene expression data from 970 breast samples was carried out to evaluate the expression pattern of *PLA2G7* mRNA in breast cancer. Second, the expression profile of *PLA2G7* was studied in 1042 breast cancer samples including 89 matched lymph node metastasis samples using immunohistochemistry. Third, the effect of *PLA2G7* silencing on genome-wide gene expression profile was studied and validated in cultured breast cancer cells expressing *PLA2G7* at high level. Last, the expression pattern of *PLA2G7* mRNA was investigated in 24 nonmalignant tissue samples and 65 primary and 7 metastatic tumour samples derived from various organs using qRT-PCR. The results from clinical breast cancer samples indicated that *PLA2G7* is overexpressed in a subset of breast cancer samples compared to its expression in benign breast tissue samples and that high *PLA2G7* expression associated with hormone receptor negativity as well as with poor prognosis in a subset of breast cancer samples. *In vitro* functional studies highlighted the putative role of *PLA2G7* in the regulation of epithelial–mesenchymal transition (EMT)–related signalling pathways, vimentin and E-cadherin protein expression as well as cell migration in cultured breast cancer cells. Furthermore, supporting the findings in breast and prostate cancer, high *PLA2G7* mRNA expression was associated with metastatic cancer in four additional organs of origin. In conclusion, our results indicate that *PLA2G7* is highly expressed in a subset of metastatic and aggressive breast cancers and in metastatic samples of various tissues of origin and promotes EMT and migration in cultured breast cancer cells.

Keywords: *PLA2G7*; epithelial–mesenchymal transition; EMT; vimentin; breast cancer; prognosis; metastasis; invasion

Received 2 January 2017; Accepted 10 March 2017

Microarray data: Data can be accessed at ArrayExpress (E-MTAB-5397 and E-MTAB-5398)

No conflicts of interest were declared.

Introduction

Breast cancer is the most frequent cancer in women worldwide accounting for 25% of all cancer cases among females [1]. Although survival rates of breast cancer patients have improved significantly during the last decade breast cancer is still the leading cause of cancer-related deaths in women due to distinct subtypes of breast cancer that are associated with early recurrence and an aggressive metastatic progression of the disease [1–4]. Identification of biomarkers and development of targeted therapies especially for these malignant tumours is important. Recently, the regulators of metastasis initiating epithelial-mesenchymal transition (EMT) have been suggested as important factors and possible diagnostic markers for aggressive and metastatic breast cancers, including triple negative breast cancers (TNBC) [5–9].

EMT, a vital process during embryogenesis and tissue repair, is characterised by specific gene expression changes, reduced intercellular adhesion, gain of mesenchymal characteristics and increased cell mobility to distant locations. Loss of immotile epithelial phenotype and gain of mesenchymal properties via EMT has recently been shown to be a key metastasis-promoting step in many cancers [10]. In addition to enabling metastasis EMT has been suggested to promote the generation and function of cancer stem cells as well as development of drug resistance [11]. EMT is induced by multiple oncogenic events and signalling pathways and requires the expression of a variety of regulators, including members of the Twist, Snail/Slug and Zeb transcription factor families. In addition, Tiwari *et al.* recently reported the SOX4-EZH2 pathway as one of the master regulators of EMT in metastatic breast cancer [12]. Two major hallmarks of EMT are the loss of expression of the epithelial cell adhesion molecule E-cadherin (CDH1) and gain of expression of the major mesenchymal cell cytoskeletal component vimentin (VIM). The cytoplasmic intermediate filament protein vimentin is a component of the cytoskeleton normally found in embryonic or mesenchymal cells. However, in cancer vimentin is a marker of EMT and is expressed in neoplastic epithelial cells with metastatic properties [13–15]. The expression of vimentin correlates with invasiveness and poor prognosis in clinical cancer samples [16–18].

PLA2G7, also known as lipoprotein-associated phospholipase A2 or platelet-activating factor (PAF) acetyl hydrolase, is a phospholipase that hydrolyses PAF and truncated phospholipids generated by oxidative attack [19]. PLA2G7 was recently reported as a novel

biomarker in 50% of primary and 70% of metastatic prostate cancers and associated with aggressive disease in prostate cancer [20]. Furthermore, PLA2G7 was identified as a potential drug target especially in *ERG* oncogene positive prostate cancers and the alterations induced by PLA2G7 silencing highlighted the potential of PLA2G7 inhibition as an anti-proliferative, pro-apoptotic and anti-migratory therapeutic approach [20,21]. Interestingly, in addition to *ERG* oncogene expressing prostate cancers, high PLA2G7 expression has also been proposed to play a causal role in colon tumourigenesis downstream of loss-of-function *TP53* mutations and activating *RAS* mutations [22]. Supporting this finding, deletion of PLA2G7 in *Apc^{Min/+}* mice has been shown to decrease both intestinal polyposis as well as colon tumourigenesis [23]. In recent years, PLA2G7 has also been under intensive research in the area of cardiovascular diseases. High PLA2G7 amount and enzyme activity have been associated with an increased risk of cardiovascular disease, and PLA2G7 inhibitor darapladib has been studied in the prevention and treatment of coronary heart disease [24–29].

In this study, we focus on analysing the expression pattern of PLA2G7 in breast cancer and investigate the functional role of PLA2G7 in cultured breast cancer cells using genome-wide gene expression analysis, immunohistochemistry and functional assays. The results indicate a role for PLA2G7 in the regulation of metastasis-enabling EMT and cell migration, associate high PLA2G7 mRNA expression with hormone receptor negative aggressive disease and indicate that high PLA2G7 protein expression in lymph node metastases associates with unfavourable prognosis in breast cancer.

Methods

Transcriptomics analysis

In order to evaluate the *in vivo* relevance of PLA2G7 in breast cancer, we analysed PLA2G7 mRNA expression levels in normal (n = 13) and malignant (n = 957) breast tissue samples using GeneSapiens database (www.genesapiens.org) [30]. Survival information was available for 395 and recurrence data for 159 breast cancer samples.

Clinical material

Breast tumour samples were obtained from patients at Hamburg University Medical Centre (Hamburg, Germany) between 1999 and 2007 and Tampere University Hospital (Tampere, Finland) between 1990 and

1999. Patient data and tumour characteristics have been described previously [31,32]. Informed consent for the scientific use of tissue materials was obtained from the patients and ethical permission was obtained from the local ethical committees. All procedures were in accordance with the Helsinki Declaration.

The patient records contained information on clinicopathological parameters, primary treatments, site and time of tumour recurrences and survival. The maximum follow-up period was 19.8 years. Relapse-free survival was defined as time to any disease recurrence (local, regional or distant). The end point in breast cancer specific survival was death from breast cancer and in overall survival death from any cause.

Immunohistochemistry

The construction of tissue microarrays (TMA) has been described previously [31,32]. The paraffin embedded breast cancer TMAs containing in total 1042 samples from 897 breast cancer patients, including 89 cases with matched primary and lymph node metastatic samples, were stained with *PLA2G7* (1:150, rabbit polyclonal, Cayman Chemical) antibody. Immunostaining was performed with the automated immunostaining device BenchMark XT (Roche Diagnostics/Ventana Medical Systems, Tucson, AZ, USA) using ultraView Universal DAB Detection Kit (Roche Diagnostics/Ventana Medical Systems). *PLA2G7* protein expression level in the epithelial breast cancer cells in each sample was evaluated independently in blinded fashion by two pathologists and graded into two groups based on staining intensity: low expression (no staining or weak staining), high expression (moderate or strong staining). After excluding the cases without residual tissue or tumour cells, 618 breast tumours and 65 lymph node metastasis samples remained eligible for evaluation.

Cell culture

The breast carcinoma cell lines MDA-MB-468, MCF-7, MDA-MB-231 and ZR-75-1 cell lines were from American Type Culture Collection (Manassas, VA, USA). BT-549 and KPL-1 cell lines were from Cell Lines Service (Eppelheim, Germany). All cell lines were maintained according to the distributor's instructions.

Gene silencing

For transient knockdown of *PLA2G7*, siRNA (SI00072177; siPLA2G7, Qiagen, Valencia, CA) was transfected into MDA-MB-468 cells using siLentFect

Lipid Reagent (Bio-Rad Laboratories, CA) in Opti-MEM (Invitrogen, Carlsbad, CA) at a final concentration of 13 nM. Cells were incubated for 48 hours before further experiments. To generate stable *PLA2G7* deficient MDA-MB-468 cells (MDA-468/*PLA2G7*⁻ cells), two lentiviral shRNA constructs (NM_005084.2-1046s1c1; shPLA2G7_1 and NM_005084.2-684s1c1; shPLA2G7_2, Sigma-Aldrich, St. Louis, MO) were transduced into MDA-MB-468 cells as described [33]. All Stars scrambled siRNA (siScr, Qiagen) and non-targeting construct (SHC002V; shScr: Sigma-Aldrich) were used as negative control.

Quantitative RT-PCR

The confirmation of effective target gene silencing was performed with Taqman reverse transcriptase PCR (qRT-PCR). Total RNA from 70 000 cells was extracted using RNeasy kit (Qiagen) according to the manufacturer's protocol and processed to cDNA with Applied Biosystems cDNA synthesis kit. TaqMan gene expression probes and primers from the Universal Probe Library (Roche Diagnostics, Espoo, Finland) were used to study *PLA2G7* (forward: tgctctaccttagaacctga; reverse: ttttgcctttgc cgtacct), and β -actin (ACTB, forward: ccaaccgcgagaagatga; reverse: ccagaggcgtacaggatag) mRNA expression. The quantitative PCR was done using ABI Prism 7900 (Applied Biosystems, Foster City, CA). Quantitation was carried out using the $\Delta\Delta$ CT method with RQ manager 1.2 software (Applied Biosystems). Average expression of the control samples was considered for the calculation of the fold changes and β -actin was used as an endogenous control. Three or more replicate samples were studied for detection of mRNA expression.

Gene expression analysis using bead arrays

500 ng of total purified RNA was used for amplification with Illumina RNA TotalPrep Amplification kit (Ambion, Austin, TX), and biotin labelled cRNA was hybridised to Sentrix HumanRef-12 vs.3 Expression Bead Chips (Illumina, San Diego, CA). Analysis was done as described [20]. Microarray data are available in the ArrayExpress database (www.ebi.ac.uk/array-express, siRNA treatment: E-MTAB-5397 and shRNA treatments: E-MTAB-5398). Two replicate samples were studied for each treatment and gene expression profiles of the *PLA2G7* silenced samples were compared with the scrambled control-treated samples. Ingenuity Pathway Analysis (IPA) Software (Ingenuity Systems Inc., Redwood City, CA, USA), DAVID Bioinformatics Resources [34,35] and

Molecular Signatures Database (MSigDB) Gene Set Enrichment Analysis Software [36,37] were used to assess the enrichment of functional and pathway annotations as well as gene sets for the genes differentially expressed by *PLA2G7* siRNA or both *PLA2G7* shRNAs.

Cell viability and apoptosis assays

The effects of *PLA2G7* silencing on cancer cell viability and induction of apoptosis were assessed with the CellTiter-Glo cell viability assay and ApoONE apoptosis assay (Promega Inc, Madison, WI, USA) according to the manufacturer's instructions. The EnVision multilabel plate reader (Perkin-Elmer, Waltham, MA) was used for signal quantification. Scrambled siRNA or shRNA was used as the negative control and *KIF11* siRNA as the positive control.

Western blotting

Western blot analysis was performed using antibodies against vimentin (1:1000, V6630, Sigma-Aldrich), E-cadherin (1:1000, Cell Signalling Technology, Danvers, MA), EGFR (1:100, Cell Signalling Technologies) and β -actin (1:5000, Sigma-Aldrich). The signals obtained were visualised with the enhanced chemiluminescence (ECL) detection system (Amersham) and normalised to β -actin signal.

Immunofluorescence staining

For the immunofluorescence staining, MDA-468/*PLA2G7*- cells were fixed with 4% paraformaldehyde (PFA) in PBS, permeabilised with 0.2% Triton X-100, and blocked with 3% bovine serum albumin in PBS. Cells were stained with vimentin antibody and Alexa-conjugated secondary antibody was used for secondary staining (1:300 dilution; Molecular Probes, Thermo Fisher Scientific, Waltham, MA, USA). Nuclei were stained with Vectashield mounting medium (Vector Laboratories, Burlingame, CA) containing DAPI. Images were taken with Zeiss Axiovert 200M fluorescence microscope (Carl Zeiss AG, Oberkochen, Germany).

Wound healing invasion assay

MDA-468/*PLA2G7*- cells were utilized in the wound healing invasion assay using the IncuCyte real-time imaging system (Essen BioScience, Ann Arbor, Michigan, USA). The cells were plated on 96-well plates (ImageLock plate, Essen BioScience, MI) coated with 50 μ l 10% Growth Factor Reduced Matrigel (BD Biosciences) after which the cells

were allowed to attach overnight at +37°C. A wound was scratched across each well (Wound Maker, Essen BioScience) and the growth medium was removed. The cells were then carefully covered with 50 μ l 25% Matrigel in normal growth medium and incubated in 37°C for 2–3 hours to allow gelling, after which 100 μ l of growth medium was carefully added to each well. The rate of invasion (wound closure through the matrix) was monitored hourly with Incucyte imaging software (Essen BioScience) for 72 hours. Invasion efficiency was determined as percentage of the relative wound confluence compared to respective negative control (regarded as 100%).

cDNA tumour panel

TissueScan™ Cancer Survey cDNA Array 96 – I (Origene, Rockville, MD, USA) consisting of 96 samples from 65 primary and 7 metastatic tumour samples as well as 24 non-malignant tissue samples derived from 8 different primary organs and provided with clinical information was utilised according to the manufacturer's instructions to analyse the expression pattern of *PLA2G7* mRNA in multiple malignancies. The expression of *PLA2G7* was quantified using qRT-PCR.


Statistical analysis

Statistical analysis of immunohistochemical samples was carried out using the SPSS program (SPSS Inc., Chicago, IL). Comparisons between categorical variables were performed using the χ^2 test. Kaplan-Meier curves were used to estimate survival probability and patient survival differences were analysed by a log-rank test. Other statistical analyses were performed using Student's t-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$) and the Pearson correlation coefficient. The results from cultured cells are presented as the mean \pm SD of at least three replicates unless otherwise stated.

Results

Correlation of *PLA2G7* mRNA expression with hormone receptor expression and prognosis in breast cancer

To evaluate the expression pattern and possible association of *PLA2G7* with poor survival in breast cancer, the GeneSapiens database was utilised to analyse *PLA2G7* mRNA expression in 13 normal breast



samples and 957 breast cancer samples [30]. Comparison of *PLA2G7* expression levels in normal breast tissues and breast cancer samples indicated high *PLA2G7* mRNA expression in a subset of breast cancers (Figure 1A). Co-expression analysis indicated *PLA2G7* mRNA to be overexpressed especially in breast cancer samples with low oestrogen and progesterone receptor (ESR1 and PGR) expression ($p = 0$ and $p = 3.4E-10$, respectively; Figure 1B) whereas the association between high *PLA2G7* mRNA and low *HER2/ERBB2* expression was not statistically significant (supplementary material, Figure S1). To investigate whether *PLA2G7* may have prognostic significance, Kaplan–Meier analysis was performed. The results revealed a significant association between high *PLA2G7* mRNA expression and poor breast cancer survival ($p = 0.004$) as well as increased risk of disease relapse ($p = 0$; Figure 1C).

To validate the association between high *PLA2G7* expression and hormone receptor negativity as well as poor prognosis in breast cancer, two additional independent sets of clinical breast cancer tissue microarrays were studied with immunohistochemical staining. *PLA2G7* expression was analysed in a total of 683 breast cancer samples, including 65 samples derived from lymph node metastases (Figure 2A). High *PLA2G7* expression was detected in 21.7% ($n = 148$) of all breast cancer samples. Association between high *PLA2G7* expression and standard clinicopathological parameters is presented in Table 1. In accordance with the results from bioinformatic analysis of *PLA2G7* mRNA expression, high *PLA2G7* expression showed significant association with hormone receptor negativity in primary the breast cancer samples analysed. Furthermore, even though no significant association with survival or relapse was seen in the samples obtained from primary tumours, high *PLA2G7* protein expression was associated with poor

Figure 1. High *PLA2G7* mRNA expression is associated with hormone receptor negativity and poor prognosis in clinical breast cancer samples. (A) Box plot analysis of normalized *PLA2G7* mRNA expression values in normal and malignant tissues. The box refers to quartile distribution (25–75%) range, with the median shown as a vertical line. Data observations which lie more than 1.5*inter-quartile range higher than third quartile are considered as outliers and indicated separately. (B) The mRNA co-expression pattern of *PLA2G7* and oestrogen receptor (ESR1) or progesterone receptor (PGR) in breast cancer samples ($n = 957$). (C) Kaplan–Meier plot of breast cancer specific survival and risk of relapse based on *PLA2G7* mRNA expression in breast cancer. A log-rank test was performed between samples with high (50%) and low (50%) expression to evaluate the prognostic significance of *PLA2G7* mRNA expression level.




Figure 2. *PLA2G7* is expressed in a subset of breast cancers and associates with poor prognosis in breast cancer lymph node metastasis samples. (A) Staining intensity of *PLA2G7* in primary breast cancer and breast cancer lymph node metastasis was scored as follows: Low expression (0 or +), high expression (++ to +++). Representative section of staining intensities is presented. The areas presented at higher magnification have been indicated in the core images. (B) Kaplan-Meier curve presentation of overall survival and risk of relapse in the patient groups with no or weak *PLA2G7* staining ($n = 57$) or positive *PLA2G7* staining ($n = 8$) in the lymph node metastasis samples.

Table 1. PLA2G7 expression and standard clinicopathological parameters in 618 breast cancer samples.

Variable	TMA 1					TMA 2 (primary tumours)				
	All (n)	All (%)	High PLA2G7 (n)	High PLA2G7 (%)	<i>p</i>	All (n)	All (%)	High PLA2G7 (n)	High PLA2G7 (%)	<i>p</i>
All samples	400	100	84	21.0		218	100	56	25.7	
Tumour type	309					218				
ILC	135	43.7	21	15.6	0.028	27	12.4	5	18.5	0.048
IDC	174	56.3	45	25.9		165	75.7	43	26.1	
Hormone receptor status	328					218				
negative	53	16.2	16	30.2	0.041	44	20.2	18	40.9	0.010
positive	275	83.8	51	18.0		174	79.8	38	21.8	
HER2 expression	330					187				
negative	285	86.4	54	18.9	0.061	160	85.6	41	25.6	0.706
positive	45	13.6	14	31.1		27	14.4	6	22.2	
WHO grade	260					213				
I	81	31.2	12	14.8	0.244	10	4.7	4	40.0	0.018
II	126	48.5	27	21.4		116	54.5	21	18.1	
III	53	20.4	14	26.4		87	40.8	30	34.5	
pT stage	342					218				
pT1	218	63.7	51	23.4	0.227	109	50.0	29	26.6	0.973
pT2	98	28.7	14	14.3		91	41.7	23	25.3	
pT3	12	3.5	1	8.3		10	4.6	2	20.0	
pT4	14	4.1	3	21.4		8	3.7	2	25.0	
Lymph node metastasis	322					217				
pN0	205	63.7	38	18.5	0.538	129	59.4	31	24.0	0.590
pN1	117	36.3	25	21.4		88	40.6	24	27.3	
Metastasis	341					214				
M0	330	96.8	68	20.6	0.350	205	95.8	50	24.4	0.543
M1	11	3.2	1	9.1		9	4.2	3	33.3	
Recurrence	345					208				
no	316	91.6	67	21.2	0.065	161	77.4	40	24.8	0.211
yes	29	8.4	2	6.9		47	22.6	16	34.0	

ILC, invasive lobular carcinoma; IDC, invasive ductal carcinoma. Numbers in bold indicate statistically significant *P* values.

survival (*p* = 0.002) as well as increased risk of disease relapse (*p* = 0.005) in the samples obtained from lymph node metastases (Figure 2B).

No significant difference was detected in PLA2G7 expression between the paired primary tumour and metastasis samples. A staining result was obtained for 39 paired primary tumours and lymph node metastases and 72% (28) of the pairs showed similar staining intensity in primary and metastatic samples.

PLA2G7 mRNA expression in cultured breast cancer cells

In order to find a suitable breast cancer cell model for functional experiments the mRNA expression level of PLA2G7 was studied in six breast cancer cell lines with qRT-PCR. Three of the cell lines studied were ER-positive (MCF-7, KPL-1 and ZR-75-1) and three were ER-negative (MDA-MB-231, BT-549 and MDA-MB-468). The results from qRT-PCR analysis are in accordance with the clinical data and show high PLA2G7 expression in two hormone receptor

negative cell lines, MDA-MB-468 and BT-549 (Figure 3A).

The effect of PLA2G7 silencing on genome-wide gene expression in MDA-MB-468 breast cancer cells

To investigate the molecular processes regulated by PLA2G7 in breast cancer cells, the effect of PLA2G7 silencing on genome-wide gene expression profile was studied in MDA-MB-468 cells expressing PLA2G7 at high level. The gene expression analysis was conducted using one siRNA and two different shRNA molecules silencing PLA2G7 efficiently compared to the respective scrambled control (Figure 3B and supplementary material, Figure S2A). For MDA-468/PLA2G7- cells the genes uniformly differentially regulated by both of the shRNAs were selected for further analysis.

DAVID Bioinformatics Resources and Ingenuity Pathway Analysis (IPA) software was used to assess enrichment of functional and pathway annotations for




Figure 3. *PLA2G7* mRNA is highly expressed in triple negative breast cancer cell lines. (A) *PLA2G7* mRNA expression in oestrogen receptor positive and oestrogen receptor negative breast cancer cell lines. (B) Validation of stable *PLA2G7* gene silencing in MDA-MB-468 breast cancer cells at the mRNA level.

the genes differentially expressed by *PLA2G7* impairment. In accordance with earlier functional studies in prostate cancer cells [20], in transiently silenced *PLA2G7* siRNA transfected breast cancer cells, the gene expression profiling highlighted the effect of *PLA2G7* on proliferation (cell cycle) and apoptosis in addition to cell motion and adhesion (supplementary material, Table S1). However, functional cell viability and apoptosis assays with siRNA treated MDA-468 breast cancer cells showed only a minor decrease in cell viability at the 48 hours time point (supplementary material, Figure S2B). Interestingly, the results from genome-wide gene expression profiling using MDA-468/*PLA2G7*⁻ cells indicated that *PLA2G7* silencing decreases the expression of genes associated with regulation of cell migration, epithelial cell differentiation and EMT (Table 2). Supporting the role of *PLA2G7* in the regulation of EMT as well as migration and invasion, *PLA2G7* impairment induced gene expression changes

associated with IL-17A (interleukin 17A), Notch and oncostatin M signalling. These processes have all been previously associated with tumour metastasis or EMT [38–42].

Validation of the effect of *PLA2G7* impairment on vimentin, E-cadherin and EGFR expression

Two major hallmarks of EMT are the loss of expression of the epithelial cell adhesion molecule E-cadherin and gain of expression of the major mesenchymal cell cytoskeletal component vimentin. To validate the role of *PLA2G7* in EMT, MDA-468/*PLA2G7*⁻ cells were analysed for vimentin and E-cadherin protein expression. The results indicated clearly reduced vimentin and increased E-cadherin protein levels (Figure 4A). In addition, immunofluorescence staining of MDA-468/*PLA2G7*⁻ cells showed a clear decrease in vimentin staining intensity in comparison to the corresponding control (Figure 4B).

Table 2. The effect of stable *PLA2G7* silencing on gene expression profile in MDA-MB-468 breast cancer cells

Down (n = 19)	
Biological processes	<i>P</i> value
Epithelial cell differentiation	6.3E-3
Regulation of cell migration	9.4E-3
Epidermis development	1.1E-2
Regulation of cell motion	1.2E-2
Ectoderm development	1.3E-2
Canonical Pathways	
Bile Acid Biosynthesis, Neutral Pathway	1.28E-2
Regulation of the Epithelial-Mesenchymal Transition Pathway	1.36E-2
Methylglyoxal Degradation III	1.48E-2
IL-17A Signalling in Gastric Cells	2.45E-2
Notch Signalling	3.7E-2
Up (n = 39)	
Biological processes	
Immune response	2.5E-3
Canonical Pathways	<i>P</i> value
Interferon Signalling	1.35E-13
Role of Pattern Recognition Receptors in Recognition of Bacteria and Viruses	2.34E-5
Activation of IRF by Cytosolic Pattern Recognition Receptors	1.83E-4
Oncostatin M Signalling	1.63E-3
Role of IL-17A in Psoriasis	2.26E-2

The functional gene ontology and pathway annotations were analysed for the sets of uniformly differentially expressed genes (FC >1.5 or FC <0.666) in response to two shRNAs. The analysis was conducted using DAVID Bioinformatics Resources and Ingenuity Pathway Analysis software




Figure 4. PLA2G7 impairment decreases the protein expression of vimentin and increases the protein expression of E-cadherin. (A) The effect of stable PLA2G7 silencing on vimentin and E-cadherin protein expression in MDA-MB-468 cells. β -actin expression was used as an endogenous control. (B) Immunofluorescence staining of MDA-MB-468 cells without (shScrambled) or with stable PLA2G7 silencing (shPLA2G7) using vimentin antibody (red). DAPI staining (blue) was used to visualize nuclei. 63 \times objective magnification.




Figure 5. PLA2G7 impairment reduces breast cancer cell invasion. (A) The effect of stable PLA2G7 silencing on MDA-MB-468 cell motility in wound healing invasion assay at 12, 24 and 48 hours after wound scratch. The relative wound closure effect was calculated as wound confluence in comparison to shScrambled control. (B) Wound closure at 48 hours in comparison to 0 hours. Scale bar = 300 μ m. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.




Figure 6. Relative PLA2G7 mRNA expression in non-malignant and malignant breast, colon, kidney, liver and lung tissue samples based on TissueScan™ Cancer Survey cDNA panel. M, metastasis; S, sarcomatoid.

Epidermal growth factor receptor (EGFR) is known to be frequently overexpressed and suggested as a potential drug target in aggressive TNBCs [43,44]. In the current study, gene expression analysis suggested *PLA2G7* as a regulator of EGFR expression in MDA-468 breast cancer cells expressing EGFR at high level. Western blot analysis of EGFR showed a significant decrease in protein expression in response to stable *PLA2G7* silencing (supplementary material, Figure S3).

Analysis of the effect of *PLA2G7* impairment on breast cancer cell invasion

To validate the putative regulatory role of *PLA2G7* on breast cancer cell migration and invasion, a wound healing invasion assay was performed with MDA-468/*PLA2G7*- cells. The results showed that stable *PLA2G7* knockdown significantly reduced cell invasion in comparison to scrambled control at 12, 24 and 48 hours (Figure 5A and B). To confirm that the effect was not due to changes in cell viability the effect of stable *PLA2G7* silencing on cell viability was studied (supplementary material, Figure S4). In accordance with the experiments conducted with *PLA2G7* siRNA transfected MDA-MB-468 cells (supplementary material, Figure S2B) stable *PLA2G7* knockdown only moderately affected MDA-468 cell viability thus confirming that the effect on cell invasion was not due to reduced cell viability.

PLA2G7 mRNA expression in cancers of various origins

The expression of *PLA2G7* mRNA was further analysed in a cDNA tumour panel consisting of 65 primary and 7 metastatic tumour samples as well as 24 non-malignant tissue samples derived from 8 different primary organs. Interestingly, the results from breast, colon, kidney, liver and lung derived samples showed the highest *PLA2G7* expression in the aggressive and metastatic cancer samples (Figure 6A–E). In ovarian cancer samples, no similar pattern was seen (supplementary material, Figure S5A). In accordance with earlier reports [20], the prostate cancer sample with the highest expression was the only true cancer sample in the set (supplementary material, Figure S5B). Surprisingly, in thyroid samples the expression pattern was the opposite and normal tissues showed higher *PLA2G7* expression than tumour tissues (supplementary material, Figure S5C).

Discussion

Breast cancer is the most frequent cancer in women worldwide and the leading cause of cancer-related deaths in women due to distinct cancer subtypes associated with aggressive progression [1–4]. Here, we evaluated the expression pattern of *PLA2G7* in breast cancer and investigated the molecular processes regulated by *PLA2G7* in cultured breast cancer cells.

The results from three independent clinical breast cancer sample sets showed that *PLA2G7* is highly expressed in hormone receptor negative breast cancers. Analysis of transcriptomic data indicated a significant association between high *PLA2G7* mRNA expression and poor prognosis in primary breast cancer samples. However, the association between high *PLA2G7* protein expression and unfavourable disease outcome was not seen at the protein level in samples derived from primary breast cancers. Surprisingly, when evaluating the protein expression of *PLA2G7* in the lymph node metastases, high *PLA2G7* protein expression was significantly associated with poor prognosis. The reason for the inconsistent results in survival analyses of primary breast cancer samples using either *PLA2G7* mRNA or protein expression remain to be investigated. The qRT-PCR primers and antibody used for evaluation of *PLA2G7* expression detect both of the transcripts listed in the current release of the Ensembl database [45]. It can be hypothesised that the results may be affected by possible *PLA2G7* protein secretion or the expression of *PLA2G7* by stromal and inflammatory cells present in the samples. However, the association between high *PLA2G7* expression and poor prognosis using samples from lymph node metastases was acquired using the protein expression of *PLA2G7* only in the tumour cells, supporting the role for *PLA2G7* in cancer progression. Furthermore, *PLA2G7* was also highly expressed in metastatic or aggressive tumours derived from cancers of other origins, indicating a wider range of significance in cancer progression. However, although high *PLA2G7* expression associates with aggressive tumours and metastasis in many tissues, the results were opposite in thyroid tissue, suggesting that the role of *PLA2G7* is multifaceted and dependent on the nature of the disease and biological context.

To gain biological insights into the role of *PLA2G7* in breast cancer we performed functional studies with breast cancer cells and identified a regulatory link between *PLA2G7* and metastasis-enabling EMT. The genome-wide gene expression profile

indicated that *PLA2G7* impairment altered IL-17A (interleukin 17A), Notch and oncostatin M signalling, supporting the role of *PLA2G7* in the regulation of EMT as well as migration and invasion. The pro-inflammatory cytokine IL-17A has recently emerged as a critical promoter of breast cancer metastasis, chemoresistance, angiogenesis as well as cell proliferation, whereas Notch signalling has been associated with the activation of EMT in cancer cells [38–42]. Furthermore, inhibition of Notch signalling causes growth arrest and inhibition of EMT in both breast cancer stem cells and breast cancer cells [46]. Cytokine oncostatin M has been associated with poor outcome in breast cancer [47]. In addition to regulating oestrogen receptor expression and breast cancer metastasis to bone, oncostatin M has been reported to mediate mesenchymal and stem cell-like differentiation in breast cancer [47–49].

The effect of *PLA2G7* depletion on the protein levels of key EMT regulators, vimentin and E-cadherin as well as the invasive phenotype was then monitored in cultured breast cancer cells expressing *PLA2G7* at high level. The MDA-468 and BT-549 cell lines were derived from aggressive triple-negative breast tumours harbouring inactivating mutations in tumour suppressor genes *PTEN*, *RBI* and *TP53* [50]. Mutations in *TP53* are found in the majority of TNBCs and targeted deletion of both *Trp53* and *Rbl* in mouse mammary stem progenitor cells has been reported to induce aggressive EMT-type mammary tumours [51–53]. Inhibition of *PLA2G7* expression resulted in decreased expression of vimentin, increased expression of E-cadherin as well as reduced cell invasion. The cytoplasmic intermediate filament protein vimentin is a central marker of EMT in cancer. Vimentin is expressed in neoplastic breast epithelial cells with metastatic properties as well as in aggressive and metastatic clinical breast cancers, supporting the association between *PLA2G7* and aggressive disease in breast cancer [14–16]. Furthermore, these results indicate that *PLA2G7* is required for cytoskeletal alterations enabling cell migration and EMT and highlight a significant role for *PLA2G7* in breast cancer metastasis. Regulators of EMT have been recently suggested as major therapeutic targets of aggressive and metastatic breast cancers, including triple negative breast cancers (TNBC) [5–9,12]. Further studies are needed to understand the mechanisms behind the ability of *PLA2G7* to regulate EMT and invasion in more detail.

In addition to regulating EMT and invasion, *PLA2G7* impairment was found to inhibit EGFR expression *in vitro*. Although recent results from clinical trials studying the combinatorial effect of monoclonal anti-EGFR antibody cetuximab and carboplatin or cisplatin in the

treatment of TNBC have not been successful, combining different therapies has been suggested as a necessary step to reach significant therapeutic progress [54–59]. Interestingly, darapladib, a novel *PLA2G7* inhibitor, has recently been under intensive research in the area of cardiovascular diseases. In addition, lipid-lowering statins are known to reduce *PLA2G7* activity in plasma, atherosclerotic plaques and cultured prostate cancer cells [20,25,60,61]. Thus, although the first phase III trials with darapladib did not produce much anticipated promising results in the treatment of chronic coronary heart disease, pharmacologically well-studied and well-known molecules inhibiting *PLA2G7*, such as darapladib and statins, could be readily translated to *in vivo* preclinical studies and clinical trials also in the area of oncology. Our results propose a possible beneficial effect from the combination of *PLA2G7* inhibition with other therapies (ie EGFR targeted therapy) in advanced breast cancer.

In conclusion, results from this study propose *PLA2G7* as an EMT and metastasis-promoting enzyme in a subset of breast cancers and link the expression of *PLA2G7* with hormone receptor negativity and poor prognosis especially in metastatic breast cancer. Furthermore, in addition to breast and prostate cancer, results from this study also suggest a possible association between high *PLA2G7* expression and metastasis in four other malignancies; namely colon, kidney, liver and lung cancer. Recent studies further link *PLA2G7* with nasopharyngeal carcinoma invasiveness and early relapse in ovarian cancer [62,63]. Thus, these results present a rationale for further exploring the role of *PLA2G7* in cancer progression and the putative potential of *PLA2G7* targeted therapy in cancer management.

Acknowledgements

We thank Sinikka Kollanus for performing immunohistochemical staining and Taina Korpela for excellent technical assistance. We thank the Finnish DNA Microarray Centre for performing microarray and TaqMan qRT-PCR analysis, and Vidal Fey as well as Arho Virkki for the normalisation of gene expression analysis results. John-Patrick Mpindi is acknowledged for help in bioinformatics analyses from clinical sample data. This work was supported by Cancer Organizations of Finland, Sigrid Juselius Foundation, TIME collaborative project, Maud Kuistila Memorial Foundation, K. Albin Johansson's Stiftelse, Research Foundation for Clinical Chemistry, Finnish Cultural Foundation and the Academy of Finland.

Author contributions

Conceived and designed the experiments: LL, PV, KI. Performed the experiments: LL, PV. Analysed the data: LL, PV, HW, HH, PK. Contributed reagents/materials/analysis tools: HW, KP, VM, AK, KI, OK, OC. Wrote the paper: LL, PV, KI. All authors reviewed the manuscript and had final approval of the submitted version.

References

- Torre LA, Bray F, Siegel RL, et al. Global cancer statistics, 2012. *CA: A Cancer Journal for Clinicians* 2015; **65**: 87–108.
- Lang JE, Wechsler JS, Press MF, et al. Molecular markers for breast cancer diagnosis, prognosis and targeted therapy. *J Surg Oncol* 2015; **111**: 81–90.
- Oakman C, Viale G, Di Leo A. Management of triple negative breast cancer. *Breast (Edinburgh, Scotland)* 2010; **19**: 312–321.
- Woodward WA, Strom EA, Tucker SL, et al. Changes in the 2003 American Joint Committee on Cancer staging for breast cancer dramatically affect stage-specific survival. *Journal of Clinical Oncology* 2003; **21**: 3244–3248.
- Gupta GP, Massagué J. Cancer metastasis: building a framework. *Cell* 2006; **127**: 679–695.
- Jeong H, Ryu Y, An J, et al. Epithelial-mesenchymal transition in breast cancer correlates with high histological grade and triple-negative phenotype. *Histopathology* 2012; **60**: E87–E95.
- Sarrió D, Rodríguez-Pinilla S, Hardisson D, et al. Epithelial-mesenchymal transition in breast cancer relates to the basal-like phenotype. *Cancer Res* 2008; **68**: 989–997.
- Yamashita N, Tokunaga E, Kitao H, et al. Vimentin as a poor prognostic factor for triple-negative breast cancer. *J Cancer Res Clin Oncol* 2013; **139**: 739–746.
- Zagouri F, Bago-Horvath Z, Rössler F, et al. High MET expression is an adverse prognostic factor in patients with triple-negative breast cancer. *Br J Cancer* 2013; **108**: 1100–1105.
- Gavert N, Ben-Ze'ev A. Epithelial-mesenchymal transition and the invasive potential of tumours. *Trends Mol Med* 2008; **14**: 199–209.
- Mallini P, Lennard T, Kirby J, et al. Epithelial-to-mesenchymal transition: what is the impact on breast cancer stem cells and drug resistance. *Cancer Treat Rev* 2014; **40**: 341–348.
- Tiwari N, Tiwari VK, Waldmeier L, et al. Sox4 is a master regulator of epithelial-mesenchymal transition by controlling Ezh2 expression and epigenetic reprogramming. *Cancer Cell* 2013; **23**: 768–783.
- Chu YW, Runyan RB, Oshima RG, et al. Expression of complete keratin filaments in mouse L cells augments cell migration and invasion. *Proc Natl Acad Sci U S A* 1993; **90**: 4261–4265.
- Thompson EW, Paik S, Br \bar{A} \bar{A} \bar{A} \bar{A} Nner N, et al. Association of increased basement membrane invasiveness with absence of estrogen receptor and expression of vimentin in human breast cancer cell lines. *J Cell Physiol* 1992; **150**: 534–544.
- Zajchowski DA, Bartholdi MF, Gong Y, et al. Identification of gene expression profiles that predict the aggressive behavior of breast cancer cells. *Cancer Res* 2001; **61**: 5168–5178.
- Domagala W, Lasota J, Bartkowiak J, et al. Vimentin is preferentially expressed in human breast carcinomas with low estrogen receptor and high Ki-67 growth fraction. *Am J Pathol* 1990; **136**: 219–227.
- Sabo E, Miselevich I, Bejar J, et al. The role of vimentin expression in predicting the long-term outcome of patients with localized renal cell carcinoma. *Br J Urol* 1997; **80**: 864–868.
- Gilles C, Polette M, Piette J, et al. Vimentin expression in cervical carcinomas: association with invasive and migratory potential. *J Pathol* 1996; **180**: 175–180.
- Stafforini DM. Biology of platelet-activating factor acetylhydrolase (PAF-AH, lipoprotein associated phospholipase A2). *Cardiovasc Drugs Ther* 2009; **23**: 73–83.
- Vainio P, Lehtinen L, Mirtti T, et al. Phospholipase PLA2G7, associated with aggressive prostate cancer, promotes prostate cancer cell migration and invasion and is inhibited by statins. *Oncotarget* 2011; **2**: 1176–1190.
- Vainio P, Gupta S, Ketola K, et al. Arachidonic acid pathway members PLA2G7, HPGD, EPHX2, and CYP4F8 identified as putative novel therapeutic targets in prostate cancer. *Am J Pathol* 2011; **178**: 525–536.
- McMurray HR, Sampson ER, Compitello G, et al. Synergistic response to oncogenic mutations defines gene class critical to cancer phenotype. *Nature* 2008; **453**: 1112–1116.
- Xu C, Reichert EC, Nakano T, et al. Deficiency of phospholipase A2 group 7 decreases intestinal polyposis and colon tumorigenesis in Apc(Min/+) mice. *Cancer Res* 2013; **73**: 2806–2816.
- May HT, Horne BD, Anderson JL, et al. Lipoprotein-associated phospholipase A2 independently predicts the angiographic diagnosis of coronary artery disease and coronary death. *Am Heart J* 2006; **152**: 997–1003.
- O'Donoghue M, Morrow DA, Sabatine MS, et al. Lipoprotein-associated phospholipase A2 and its association with cardiovascular outcomes in patients with acute coronary syndromes in the PROVE IT-TIMI 22 (PRavastatin Or atorVastatin Evaluation and Infection Therapy-Thrombolysis In Myocardial Infarction). *Circulation* 2006; **113**: 1745–1752.
- Oei HS, van der Meer IM, Hofman A, et al. Lipoprotein-associated phospholipase A2 activity is associated with risk of coronary heart disease and ischemic stroke: the Rotterdam Study. *Circulation* 2005; **111**: 570.
- Packard CJ, O'reilly DS, Caslake MJ, et al. Lipoprotein-associated phospholipase A2 as an independent predictor of coronary heart disease. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 2000; **343**: 1148–1155.
- Wilensky RL, Shi Y, Mohler ER, et al. Inhibition of lipoprotein-associated phospholipase A2 reduces complex coronary atherosclerotic plaque development. *Nat Med* 2008; **14**: 1059–1066.
- Serruys PW, García-García HM, Buszman P, et al. Effects of the direct lipoprotein-associated phospholipase A(2) inhibitor darapladib on human coronary atherosclerotic plaque. *Circulation* 2008; **118**: 1172–1182.
- Kilpinen S, Autio R, Ojala K, et al. Systematic bioinformatic analysis of expression levels of 17,330 human genes across 9,783

- samples from 175 types of healthy and pathological tissues. *Genome Biol* 2008; **9**: R139–R139.
31. Lehtinen L, Vainio P, Wikman H, *et al.* 15-Hydroxyprostaglandin dehydrogenase associates with poor prognosis in breast cancer, induces epithelial-mesenchymal transition, and promotes cell migration in cultured breast cancer cells. *J Pathol* 2012; **226**: 674–686.
 32. Alarmo E, Huhtala H, Korhonen T, *et al.* Bone morphogenetic protein 4 expression in multiple normal and tumour tissues reveals its importance beyond development. *Modern Pathology* 2013; **26**: 10–21.
 33. Pollari S, Kakonen SM, Edgren H, *et al.* Enhanced serine production by bone metastatic breast cancer cells stimulates osteoclastogenesis. *Breast Cancer Res Treat* 2011; **125**: 421–430.
 34. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 2009; **4**: 44–57.
 35. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009; **37**: 1–13.
 36. Subramanian A, Tamayo P, Mootha VK, *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005; **102**: 15545–15550.
 37. Mootha VK, Lindgren CM, Eriksson K, *et al.* PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 2003; **34**: 267–273.
 38. Brabletz S, Bajdak K, Meidhof S, *et al.* The ZEB1/miR-200 feedback loop controls Notch signalling in cancer cells. *Embo J* 2011; **30**: 770–782.
 39. Du J, Xu K, Fang L, *et al.* Interleukin-17, produced by lymphocytes, promotes tumour growth and angiogenesis in a mouse model of breast cancer. *Mol Med Rep* 2012; **6**: 1099–1102.
 40. Benevides L, Cardoso CRB, Tiezzi DG, *et al.* Enrichment of regulatory T cells in invasive breast tumour correlates with the upregulation of IL-17A expression and invasiveness of the tumour. *Eur J Immunol* 2013; **43**: 1518–1528.
 41. Cochaud S, Giustiniani J, Thomas C, *et al.* IL-17A is produced by breast cancer TILs and promotes chemoresistance and proliferation through ERK1/2. *Sci Rep* 2013; **3**: 3456–3456.
 42. Roy LD, Sahraei M, Schettini JL, *et al.* Systemic neutralization of IL-17A significantly reduces breast cancer associated metastasis in arthritic mice by reducing CXCL12/SDF-1 expression in the metastatic niches. *BMC Cancer* 2014; **14**: 225.
 43. Livasy CA, Karaca G, Nanda R, *et al.* Phenotypic evaluation of the basal-like subtype of invasive breast carcinoma. *Modern Pathol* 2006; **19**: 264–271.
 44. Liu D, He J, Yuan Z, *et al.* EGFR expression correlates with decreased disease-free survival in triple-negative breast cancer: a retrospective analysis based on a tissue microarray. *Med Oncol* 2012; **29**: 401–405.
 45. Cunningham F, Amode MR, Barrell D, *et al.* Ensembl 2015. *Nucleic Acids Res* 2014; **43**: D662–D669.
 46. Suman S, Das TP, Damodaran C. Silencing NOTCH signaling causes growth arrest in both breast cancer stem cells and breast cancer cells. *Br J Cancer* 2013; **109**: 2587–2596.
 47. West NR, Murphy LC, Watson PH. Oncostatin M suppresses oestrogen receptor- α expression and is associated with poor outcome in human breast cancer. *Endocr Relat Cancer* 2012; **19**: 181–195.
 48. Bolin C, Tawara K, Sutherland C, *et al.* Oncostatin M promotes mammary tumour metastasis to bone and osteolytic bone degradation. *Genes Cancer* 2012; **3**: 117–130.
 49. West NR, Murray JI, Watson PH. Oncostatin-M promotes phenotypic changes associated with mesenchymal and stem cell-like differentiation in breast cancer. *Oncogene* 2014; **33**: 1485–1494.
 50. Lehmann BD, Bauer JA, Chen X, *et al.* Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 2011; **121**: 2750–2767.
 51. Langerod A, Zhao H, Borgan O, *et al.* TP53 mutation status and gene expression profiles are powerful prognostic markers of breast cancer. *Breast Cancer Res* 2007; **9**: R30.
 52. Jiang Z, Deng T, Jones R, *et al.* Rb deletion in mouse mammary progenitors induces luminal-B or basal-like/EMT tumour subtypes depending on p53 status. *J Clin Invest* 2010; **120**: 3296–3309.
 53. Dumay A, Feugeas JP, Wittmer E, *et al.* Distinct tumour protein p53 mutants in breast cancer subgroups. *Int J Cancer* 2013; **132**: 1227–1231.
 54. Carey LA, Rugo HS, Marcom PK, *et al.* TBCRC 001: randomized phase II study of cetuximab in combination with carboplatin in stage IV triple-negative breast cancer. *J Clin Oncol* 2012; **30**: 2615–2623.
 55. Baselga J, Gómez P, Greil R, *et al.* Randomized phase II study of the anti-epidermal growth factor receptor monoclonal antibody cetuximab with cisplatin versus cisplatin alone in patients with metastatic triple-negative breast cancer. *J Clin Oncol* 2013; **31**: 2586–2592.
 56. Kim EJ, Jeong J, Bae S, *et al.* mTOR inhibitors radiosensitize PTEN-deficient non-small-cell lung cancer cells harboring an EGFR activating mutation by inducing autophagy. *J Cell Biochem* 2013; **114**: 1248–1256.
 57. Koumangoye RB, Nangami GN, Thompson PD, *et al.* Reduced annexin A6 expression promotes the degradation of activated epidermal growth factor receptor and sensitizes invasive breast cancer cells to EGFR-targeted tyrosine kinase inhibitors. *Molecular Cancer* 2013; **12**: 167–167.
 58. Morrison G, Fu X, Shea M, *et al.* Therapeutic potential of the dual EGFR/HER2 inhibitor AZD8931 in circumventing endocrine resistance. *Breast Cancer Res Treat* 2014; **144**: 263–272.
 59. Brand TM, Iida M, Dunn EF, *et al.* Nuclear epidermal growth factor receptor is a functional molecular target in triple-negative breast cancer. *Molecular Cancer Therapeutics* 2014; **13**: 1356–1368.
 60. Schaefer EJ, McNamara JR, Asztalos BF, *et al.* Effects of atorvastatin versus other statins on fasting and postprandial C-reactive protein and lipoprotein-associated phospholipase A2 in patients with coronary heart disease versus control subjects. *Am J Cardiol* 2005; **95**: 1025–1032.
 61. Racherla S, Arora R. Utility of Lp-PLA2 in lipid-lowering therapy. *Am J Ther* 2012; **19**: 115–120.

62. Reinartz S, Finkernagel F, Adhikary T, *et al.* A transcriptome-based global map of signaling pathways in the ovarian cancer microenvironment associated with clinical outcome. *Genome Biol* 2016; **17**: 108.
63. Low HB, Png CW, Li C, *et al.* Monocyte-derived factors including *PLA2G7* induced by macrophage-nasopharyngeal carcinoma cell interaction promote tumour cell invasiveness. *Oncotarget* 2016; **7**: 55473–55490.

SUPPLEMENTARY MATERIAL ONLINE

Supplementary figure legends

Table S1. The effect of transient *PLA2G7* silencing on breast cancer cell gene expression profile

Figure S1. The mRNA co-expression pattern of *PLA2G7* and *HER2/neu* (*ERBB2*) in breast cancer samples (n = 957)

Figure S2. (A) Validation of transient *PLA2G7* gene silencing in MDA-MB-468 breast cancer cells at the mRNA level. (B) The effect of 48 h *PLA2G7* and *KIF11* silencing on MDA-MB-468 cell viability. (C) The effect of 48 h *PLA2G7* and *KIF11* silencing on induction of apoptosis in MDA-MB-468 cells

Figure S3. The effect of stable *PLA2G7* silencing on EGFR protein expression in MDA-MB-468 cells. actin expression was used as an endogenous control

Figure S4. The effect of stable *PLA2G7* silencing and shScrambled expression on MDA-MB-468 cell viability at 48 h after plating equal numbers of cells

Figure S5. Relative *PLA2G7* mRNA expression in non-malignant and malignant ovarian, prostate and thyroid tissue samples based on TissueScan™ Cancer Survey cDNA panel