

# Sensitivity of Zebrafish (*Danio rerio*) Embryos to Hospital Effluent Compared to *Daphnia magna* and *Aliivibrio fischeri*

Martina WITTLEROVÁ<sup>1</sup>, Gabriela JÍROVÁ<sup>1,2</sup>, Alena VLKOVÁ<sup>1,2</sup>, Kristina KEJLOVÁ<sup>1</sup>, Marek MALÝ<sup>1</sup>, Tuula HEINONEN<sup>3</sup>, Zdeňka WITTLINGEROVÁ<sup>2</sup>, Magdaléna ZIMOVÁ<sup>1,2</sup>

<sup>1</sup>National Institute of Public Health, Prague, Czech Republic, <sup>2</sup>Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic, <sup>3</sup>Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

Received October 8, 2020

Accepted November 26, 2020

## Summary

The Fish Embryo Acute Toxicity (FET) Test was adopted by the Organisation for Economic Co-operation and Development as OECD TG 236 in 2013. The test has been designed to determine acute toxicity of chemicals on embryonic stages of fish and proposed as an alternative method to the Fish Acute Toxicity Test performed according to OECD TG 203. In recent years fish embryos were used not only in the assessment of toxicity of chemicals but also for environmental and wastewater samples. In our study we investigated the acute toxicity of treated wastewater from seven hospitals in the Czech Republic. Our main purpose was to compare the suitability and sensitivity of zebrafish embryos with the sensitivity of two other aquatic organisms commonly used for wastewater testing – *Daphnia magna* and *Aliivibrio fischeri*. For the aim of this study, in addition to the lethal endpoints of the FET test, sublethal effects such as delayed heartbeat, lack of blood circulation, pericardial and yolk sac edema, spinal curvature and pigmentation failures were evaluated. The comparison of three species demonstrated that the sensitivity of zebrafish embryos is comparable or in some cases higher than the sensitivity of *D. magna* and *A. fischeri*. The inclusion of sublethal endpoints caused statistically significant increase of the FET test efficiency in the range of 1-12 %. Based on our results, the FET test, especially with the addition of sublethal effects evaluation, can be considered as a sufficiently sensitive and useful additional tool for ecotoxicity testing of the acute toxicity potential of hospital effluents.

## Key words

Fish Embryo Toxicity (FET) test • Hospital wastewater • Acute toxicity • Aquatic organisms • Sublethal endpoints

## Corresponding author

G. Jírová, National Institute of Public Health, Šrobárova 49/48, 100 00 Praha 10, Czech Republic. E-mail: gabriela.jirova@szu.cz

## Introduction

The Fish Embryo Acute Toxicity Test (FET test) was adopted by the Organisation for Economic Co-operation and Development as OECD TG 236 in 2013 (OECD 2013). The method using fish embryos as the testing organisms has been primarily designed to determine acute toxicity of chemicals. Fish embryos as non-feeding developmental stages are not categorized as protected vertebrates according to the European Directive 2010/63/EU on the protection of animals used for scientific purposes (European Parliament and Council 2010). Therefore, the FET test is considered as an alternative to experiments with adult fish with a good correlation with the Fish Acute Toxicity Test by OECD TG 203 (OECD 2019) as reported in studies by Lammer *et al.* (2009) and Dang *et al.* (2017). Recently, its use has been extended also to the assessment of complex mixtures (e.g. environmental samples, wastewater (WW), construction products). The zebrafish embryos represent a useful model with a wide range of possible applications in environmental hazard and risk assessment (Scholz *et al.* 2008). However, the FET test sensitivity to substances is variable and its applicability domain has not yet been fully defined (Dang *et al.* 2017, Sobanska *et al.* 2017).

Hospital wastewater (HWW) is a source of

diverse pollutants which may have negative impact mainly on the aquatic environment (Pérez-Alvarez *et al.* 2018). Hospital effluents are prone to higher ecotoxicological potential than urban wastewater (Laquaz *et al.* 2017). This fact is due to the content of wider spectrum and higher quantity of pharmaceutical compounds (Santos *et al.* 2010, Wiest *et al.* 2018) and other chemicals. In their review Orias and Perrodin (2013) highlighted a great diversity of the substances present within hospital effluents, and in some cases their high ecotoxicity. Due to the requirements of the EU Water Framework Directive (European Parliament and Council 2000), the quality of surface waters in the EU has been constantly improving in recent decades. One of the main objectives is to improve the treatment processes at WW treatment plants aiming to significantly reduce the discharge of undesirable substances, including pharmaceuticals, pesticides, industrial substances and human care products. Although HWW is mostly treated by WW treatment plants before discharge into the sewage system or the environment, numerous pharmaceuticals and other chemicals are insufficiently removed during treatment and end up in surface water (Wigh *et al.* 2016, Vålitalo *et al.* 2017). Therefore, in our research we investigated HWW to evaluate the efficacy of the treatment processes within the Czech Republic. Moreover, WW samples were selected from different hospital types and sizes with the intention to consider the variability of their composition.

In the review focused on conventional and alternative bioassays suitable for ecotoxicological evaluation of WW, Jirova *et al.* (2016) highlighted the need to supplement the conventional methods (based on bacteria, algae, crustaceans, fish and seeds) with the FET test as an alternative approach to vertebrate ecotoxicity tests and a useful tool for efficient detection of acute toxicity. However, scientific data regarding the sensitivity and suitability of the FET test for hospital effluent testing are scarce so far. Therefore, the primary objective of our study was to extend knowledge of this issue and make a comparative assessment of the tests based on the zebrafish embryos and other used aquatic species.

For the determination of ecotoxicity of WW from health care facilities, a battery of three bioassays with *Daphnia magna* (ISO 6341), *Aliivibrio fischeri* (ISO 11348-2) and *Desmodesmus subspicatus* (ISO 8692) is recommended by the Czech standard ČSN 756406 (2020). In the study investigating treated effluents from

five different hospitals, Jirova *et al.* (2018) concluded that the battery of three species consisting of *D. magna*, *A. fischeri* and *D. subspicatus* may be appropriate for routine testing of ecotoxicological potential of hospital effluents. For our purpose, based on the conclusions of foregoing and numerous other studies (Abbas *et al.* 2018, Ellepola *et al.* 2020, Laquaz *et al.* 2017, Li *et al.* 2017, Vasconcelos *et al.* 2017), *D. magna* and *A. fischeri* were selected as sufficiently sensitive comparative test organisms.

The international standard ISO 15088 was adopted in 2007 and its merit is to determine acute toxicity of WW to zebrafish eggs after 48 hpf (hours post fertilization) of exposure. Therefore, in the present study we compared the results of two final test exposure times 48 hpf (according to the ISO 15088) and 96 hpf (according to the FET test).

Based on their experience with the FET test, many authors recommend the inclusion of further endpoints to increase the sensitivity of the assay. Braunbeck *et al.* (2015) recommended more research to better define the domain of applicability of the FET test and suggested modifications of the method with addition of more endpoints for the detection of a multitude of toxic effects. In a comparative study, Stelzer *et al.* (2018) indicated the need to complement the FET test with sublethal metrics which would increase its efficiency, and also highlighted the necessity to validate this statement by further studies using other WW samples. For the aim of our study, in addition to the lethal endpoints of the FET test (coagulated embryos, lack of somite formation, non-detachment of the tail and lack of heartbeat), sublethal effects comprising delayed heartbeat, lack of blood circulation, pericardial and yolk sac edema, spinal curvature and pigmentation failures were evaluated.

In this study the WW samples from seven different hospitals were investigated to compare the suitability and sensitivity of zebrafish embryos with the sensitivity of two other aquatic organisms – *D. magna* and *A. fischeri*. Other purposes were to evaluate the sensitivity of the FET test after inclusion of sublethal parameters and to confirm the increase in efficiency of the method after prolongation of exposure time.

## Methods

### *Wastewater samples*

The samples were collected in 2019 from the outlets of WW treatment plants of seven different

hospitals located in the central, southern and eastern regions of the Czech Republic. Table 1 shows general characteristics of the hospitals and their WW treatment plants. The sample H1 was a mixture of treated HWW and untreated hospital laundry WW. One sample was

taken from each hospital. The samples were collected after treatment activities before discharging into the urban sewer system, stored in cooling boxes, transported to the laboratory and deep frozen at  $\leq 18$  °C prior to analysis.

**Table 1.** Characteristics of hospitals and their wastewater treatment plants.

Hospital	Type of hospital	Sizing (beds)	Wastewater treatment process	Wastewater generation ( $\text{m}^3 \cdot \text{d}^{-1}$ )	Disinfection process
H1	General	476	Mechanical-Biological	10	$\text{Cl}_2$
H2	University	1006	Mechanical-Biological	50	NaClO
H3	University	2199	Mechanical-Biological	50-100	NaClO
H4	General	500	Mechanical-Biological	30-50	$\text{Cl}_2$
H5	General	423	Mechanical-Biological	40	Do not chlorinate
H6	General	1447	Mechanical-Biological	360	$\text{Cl}_2$
H7	University	1600	Mechanical-Biological	100-120	NaClO

#### Ecotoxicological tests

Ecotoxicological potential of hospital WW was described using three organisms which represented three trophic levels: *A. fischeri* (formerly *Vibrio fischeri*) as a decomposer, *D. magna* as a primary consumer and *Danio rerio* as a secondary consumer. Characteristics of bioassays used in the study are summarized in Table 2.

As a first step, undiluted samples were analyzed by means of *D. magna* and *D. rerio* assays. In the case of *A. fischeri* test, 80 % concentration of the sample was used in accordance with the method procedure. The

results were expressed as percentage of toxic effect on each species.

If the values of toxic effect exceeded 50 %,  $\text{EC}_{50}$  or  $\text{LC}_{50}$  (effective or lethal concentration of the sample that caused negative impact in 50 % of the tested organisms) were calculated by the probit method. The experiments were conducted using a series of increasing concentrations depending on the level of HWW toxicity.

Every test was performed in two independent runs and the results were expressed as mean.

**Table 2.** Characteristics of bioassays used in the study.

Method	Organism	Standard	Exposure time	Endpoint
<i>Daphnia immobilization test</i> ( <i>D. magna</i> )	<i>Daphnia magna</i>	ISO 6341	48 h	Immobility [%] $\text{EC}_{50}$ [%]
<i>Luminescence inhibition test</i> ( <i>A. fischeri</i> 15 min, 30 min)	<i>Aliivibrio fischeri</i>	ISO 11348-2	15 min 30 min	Luminescence inhibition [%] $\text{EC}_{50}$ [%]
FET test (FET 48 h)	<i>Danio rerio</i> embryo	OECD TG 236	48 hpf	Mortality [%] $\text{LC}_{50}$ [%]
FET test (FET 96 h)	<i>Danio rerio</i> embryo	OECD TG 236	96 hpf	Mortality [%] $\text{LC}_{50}$ [%]
FET test (FET 96 h+sub)	<i>Danio rerio</i> embryo	OECD TG 236 modified	96 hpf	Mortality + sublethal effects [%] $\text{LC}_{50}$ [%]

hpf: hours post fertilization.

*Fish Embryo Acute Toxicity (FET) test*

The FET test was performed according to OECD TG 236 (OECD 2013). Three different protocols of the method were carried out: the test with basic exposure time 96 hpf (hours post fertilization), the test with reduction of exposure time to 48 hpf and the test with addition of five sublethal endpoints with exposure time 96 hpf.

The breeding stock of AB line of *D. rerio* wild type was maintained in Zebrafish Housing System ZebTEC Stand-Alone (Tecniplast) with continuous treatment and monitoring of important water parameters including pH, conductivity and temperature. Mature zebrafish aged between 6 and 18 months were used to obtain fertilized fish eggs, which were collected from a minimum of three breeding groups and selected for testing at random.


Spawning took place in two-liter breeding tanks which capitalize on the fish natural tendency to spawn in shallow water. After spawning, the eggs were rinsed with maintenance water adjusted to pH 6.5 to 7.5 and final hardness 70 to 100 mg·l<sup>-1</sup> CaCO<sub>3</sub>. The eggs were checked under inverted microscope. Viable embryos maximally at the 16 cell-stage without obvious irregularities were individually placed into polystyrene 24-well plates and incubated under 26±1 °C with 16:8 light:dark cycle. Each well contained 2 ml of test solution. Five sample concentrations were prepared in each test run. 20 eggs per concentration were exposed to the samples. Dilution water with a final concentration of 294.0 mg·l<sup>-1</sup> CaCl<sub>2</sub> × 2H<sub>2</sub>O, 123.3 mg·l<sup>-1</sup> MgSO<sub>4</sub> × 7H<sub>2</sub>O, 63.0 mg·l<sup>-1</sup> NaHCO<sub>3</sub>, 5.5 mg·l<sup>-1</sup> KCl (as defined in ISO 7346-2) was used for the preparation of the test concentrations and negative controls. The sensitivity of embryos was controlled by

3,4-dichloroaniline (4 mg/l) as positive control in each test run. All experiments were carried out in a static way without change of exposure solution. The test was considered as valid when oxygen concentration was maintained above 80 % during the test and mortality in the negative controls did not exceed 10 %.

Embryos were observed for four lethal and five sublethal effects. Lethal parameters described in the OECD TG 236 (OECD 2013) comprised coagulation of embryos, lack of somite formation, non-detachment of the tail and lack of heartbeat. Added sublethal endpoints included delayed heartbeat, lack of blood circulation, presence of edema (pericardial or yolk sac), spinal curvature and pigmentation failures. Examples of the observed sublethal effects are illustrated in Figure 1. Embryos placed in the culture plates were examined separately using inverted microscope Olympus CKX53 equipped with a digital camera Canon EOS 2000D. For determination of heart rate, embryos were video recorded and heartbeat was counted when the video was slowed down. The heart rate of embryos exposed to the samples was evaluated in relation to the negative control groups.

The cumulative mortality of zebrafish embryos was recorded at 48 hpf (termed FET 48 h) and 96 hpf (termed FET 96 h) in experimental and control groups. The results were calculated as percentage of mortality (toxic effect) of the tested embryos, or as LC<sub>50</sub>.

After the addition of sublethal effects to the lethal endpoints, the embryos development was evaluated at 96 hpf (termed FET 96 h+sub). Sublethal criteria were counted only if lethal parameters were not observed. The results were calculated as percentage of toxic effects on the tested embryos, or as EC<sub>50</sub>.



**Fig. 1.** Representative sublethal abnormalities of zebrafish embryos after 96 hpf (hours post fertilization) exposure to the hospital wastewater samples. PE=pericardial edema, YSE=yolk sac edema, SP=spinal curvature, PF=pigmentation failure.

### *Daphnia immobilization test*

The acute toxicity test with *D. magna* Straus was carried out according to ISO 6341 (2012). Neonates (less than 24 h old specimens of *D. magna*) of at least third generation originated from laboratory culture were exposed to five designated dilutions of samples. 20 organisms were placed in each concentration and negative control (dilution water according to ISO 6341 was identical to the dilution water used in the FET test mentioned above) and incubated 48 h under the following test conditions: temperature  $20 \pm 2$  °C, 16:8 light:dark cycle, oxygen concentration  $\geq 2$  mg·l<sup>-1</sup>, no feeding. For the experiment validity, the inhibition of mobility in the negative controls had to be  $\leq 10$  %. The sensitivity of test organisms was monitored by regular tests with K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> used as positive control.

The results were expressed as percentage of *Daphnia* immobilization (toxic effect), or as EC<sub>50</sub> after 48 h exposure.

### *Aliivibrio bioluminescence inhibition test*

The luminescent bacteria test followed ISO 11348-2 (2007). The test was performed using liquid-dried bacteria *A. fischeri* NRRL-B-11177 (HACH LANGE). Bacteria were activated by rehydration with reactivation solution. For the experiments, the conductivity of the samples was  $>10$  mS·cm<sup>-1</sup>, pH 6.0 to 8.5 and oxygen concentration  $>3$  mg·l<sup>-1</sup>. 2 % NaCl solution was used as dilution water and negative control. The suspensions of bacteria and diluted samples (ten sample concentrations in replicates) were incubated at  $15 \pm 1$  °C. Luminescence was measured after 15 min and 30 min of exposure to the sample concentrations series by luminometer Sirius (Berthold Detection Systems). The bacteria sensitivity was controlled using positive control (ZnSO<sub>4</sub> × 7H<sub>2</sub>O).

The results were presented as percentage of inhibition of the bacteria light emission (toxic effect) with respect to the negative control, or as EC<sub>50</sub>.

### *Statistical analysis*

Linear mixed model was used to evaluate the results of the monitoring of wastewater ecotoxicity and to assess the difference between methods (fixed effect) while taking into account the variability between hospitals (random effect). The dependent variable was angularly transformed percentage (of toxic effect). Transformed percentages more closely approximate the normal distribution (Sokal and Rohlf 1995). When

statistically significant effects were identified, Šidák's multiple comparisons procedure was applied to ascertain which specific methods differed. Values of  $p < 0.05$  were considered as indicating a statistically significant result. All statistical analyses were performed using Stata software package, release 14.2 (Stata Corp LP, College Station, TX, USA).

## Results

### *Comparison of the methods sensitivity based on the toxic effects values*

To compare the sensitivity of three aquatic species to the acute toxicity of HWW, the samples from seven different hospitals were investigated. In the first part of the study, the percentage of immobility (*D. magna*), luminescence inhibition (*A. fischeri*), mortality and sublethal effects (*D. rerio* embryos) after exposure to HWW samples were expressed as toxic effects. The results are shown in Figure 2a. The obtained data demonstrated highly variable toxic impact of hospital effluents on the studied organisms. The tests FET 96 h and FET 96 h+sub indicated high sensitivity of zebrafish embryos to five samples out of seven samples (i.e. H1, H3-H6) with toxic effect values in the range of 62.5-100 %. The results of other methods for these five samples were more variable with values ranging from 0 % to 100 % of crustacean immobility and fish embryo mortality (FET 48 h) and from 0 % to 98.4 % of bacteria luminescence inhibition. Based on the results of all samples, no statistically significant differences were found between FET tests (FET 48 h, FET 96 h, FET 96 h+sub) and *D. magna* test ( $p=0.123$ ,  $p=0.507$ ,  $p=0.275$  respectively). However, the results showed statistically significant differences between FET 96 h and *A. fischeri* 15 min, 30 min ( $p=0.018$ ,  $p=0.024$  respectively) and between FET 96 h+sub and *A. fischeri* 15 min, 30 min ( $p=0.006$ ,  $p=0.008$  respectively).

### *Comparison of methods sensitivity based on the EC<sub>50</sub>, LC<sub>50</sub> values*


In the following study, if the values of toxic effect exceeded 50 % (H1, H3-H6), EC<sub>50</sub> or LC<sub>50</sub> values were calculated. The results are presented in Figure 2b. In the case of samples which were assessed as the most toxic and which negatively affected the majority of the test species (H1, H3, H6), *D. magna* was the most sensitive organism with EC<sub>50</sub> values ranging from 4.5 to 37.6 %. The method sensitivity to the samples H1 and

H3 decreased in the following order: *D. magna*>FET 96 h+sub>FET 96 h>*A. fischeri* 15 min>FET 48 h and to sample H6: *D. magna*>*A. fischeri* 15 min>*A. fischeri* 30 min>FET tests.

*Comparison within the FET tests*

When we made a comparison within the different protocols of the FET test, a decreasing sensitivity of each method was observed as follows: FET 96 h+sub>FET 96 h>FET 48 h. Mortality was manifested most frequently by lack of heartbeat, growth retardation, and less commonly by coagulation of embryos. As shown in Figure 2b, the inclusion of sublethal endpoints has reduced the LC<sub>50</sub> values for all samples with statistically

significant increment of the FET test sensitivity (p=0.026) in the range of 1-12 %. The presence of sublethal effects was variable not only between different samples but it also varied depending on the exposure time (Table 3). Prolonged exposure time caused a statistically significant increase of the zebrafish embryos sensitivity (p<0.001) in 57 % of the tested samples. The frequency of occurrence of sublethal criteria was: spinal curvature 34.5 %, lack of blood circulation 26.4 %, delayed heartbeat 25.7 %, pigmentation failures 12.7 %, occurrence of edema 0.7 % of all sublethal parameters we observed. According to Kimmel *et al.* (1995), embryos development in control groups was normal during all the experiments.



**Fig. 2.** Ecotoxicity of wastewater samples from different hospitals (H1-H7) expressed as: (a) toxic effect [%] of undiluted samples and (b) EC<sub>50</sub>, LC<sub>50</sub> values [%]. Comparison between *Daphnia magna* (Dm), *Aliivibrio fischeri* (Af) and *Danio rerio* embryos. FET=Fish Embryo Acute Toxicity Test.

**Table 3.** Overview of the observed sublethal effects on zebrafish embryos after 48 hpf and 96 hpf (hours post fertilization) exposure to the wastewater samples from different hospitals (H1-H7). Sublethal effects were measured only if lethal effects were not observed.

Sublethal effects	H1	H2	H3	H4	H5	H6	H7
	48 hpf/96 hpf						
<i>Delayed heartbeat</i>	-/+	-/-	+/+	-/+	-/+	+/+	-/-
<i>Lack of blood circulation</i>	-/+	-/-	+/+	-/+	-/+	+/+	-/-
<i>Presence of edema</i>	-/+	-/-	+/+	-/+	-/+	-/-	-/-
<i>Spinal curvature</i>	-/+	-/-	+/+	-/+	-/+	-/+	-/-
<i>Pigmentation failures</i>	-/+	-/-	+/+	-/+	-/+	+/+	-/-

- not observed, + observed.

## Discussion

The problem of the discharge of chemical contaminants from hospitals into wastewater and subsequently into the aquatic ecosystems is well-known (Datel *et al.* 2020). The presence of micropollutants such as pharmaceuticals, disinfectants and personal care products in treated effluents poses considerable ecological risk for the aquatic environment (Meza *et al.* 2020, Rogowska *et al.* 2020). Numerous studies have been carried out to determine ecotoxicity of target substances commonly detected in HWW (Halling-Sørensen *et al.* 2000, Cleuvers 2003, Cleuvers 2005, Brandhof *et al.* 2010, Martins *et al.* 2012, Li *et al.* 2016, Romanucci *et al.* 2019). However, examination of individual target compounds, causing ecotoxicity does not enable to determine the interactions between all known and unknown hazardous substances which are present in HWW and to evaluate their overall impact on aquatic organisms. The mixture of two or more substances may produce synergistic, antagonistic and additive interactions which may increase or decrease the resulting WW ecotoxicity (Emmanuel *et al.* 2005, Godoy *et al.* 2019).

Therefore, it is necessary to investigate the whole effluent as a complex mixture of substances. In our present study a high variability of toxic effects of WW samples from individual hospitals on the used aquatic organisms was demonstrated. Similarly to these conclusions, Jírová *et al.* (2018) confirmed considerable different levels of ecotoxicity of treated wastewater from five hospitals. This variability relates to the different wastewater quality which may be influenced by a number of factors comprising current therapeutic procedures, the type and specialization of the hospital, its location, number of inpatients, flow rate, season, the day of the week and the daily period (Boillot *et al.* 2008). Good correlation was demonstrated between the amount of antibiotics in the wastewater and their volume currently being administered to patients (Hamjinda *et al.* 2015). Additionally, the insufficient degradability of certain substances during the cleaning process in the wastewater treatment plants causes the presence of residues of pollutants in wastewater (Li *et al.* 2017). A literary review showed that the quality of treatment processes is another important factor which affects the treated wastewater composition (Verlicchi *et al.* 2015).

A specific type of effluents is generated from the hospital laundry. It has a different composition from WW

produced by other hospital departments having a high concentration of organic and microbial loads depending on the washing stage. In the study of Kern *et al.* (2015), acute toxicity of hospital laundry WW was investigated by *D. magna* immobilization test and adult *D. rerio* lethality test. The obtained data showed extremely toxic impact on *D. magna* (48 h EC<sub>50</sub>=2.01 %) and lower acute toxicity for *D. rerio* (48 h LC<sub>50</sub>=29.25 %). In our study, the sample that can be considered as the most toxic (H6) because of its low EC<sub>50</sub> and LC<sub>50</sub> values (Fig. 1), was the mixture of treated hospital WW and untreated laundry WW before discharging to the sewage system. It can be assumed that the high level of acute toxicity was caused by the complement of untreated laundry effluent to common HWW.

Ecotoxicological evaluation of WW should be performed using the battery of indicator organisms from different trophic levels as different species may show diverse levels of sensitivity to the pollutants. The results of our study, as well as those of Jirova *et al.* (2018), showed the suitability of *D. magna* and *A. fischeri* for routine testing of HWW ecotoxicity. To our knowledge, the information on effects of HWW on fish early life stages using the FET test, is scarce. Stelzer *et al.* (2018) compared the FET test and other standard fish protocols used worldwide for WW analyses. LC<sub>50</sub> value of untreated hospital effluent determined by the FET test was 53.5 %. Based on findings of this study, the zebrafish embryos may not represent the most sensitive developmental stage of *D. rerio* compared to larvae and juvenile. In another study, a mixture of urban and hospital effluents was evaluated for ecotoxicity by Wigh *et al.* (2016). After 96 h of exposure to the untreated WW, 100 % of zebrafish embryos died. The mortality decreased to 13 % after the WW treatment. The addition of sublethal criteria for evaluation, such as the presence of edema, blood circulation defects, malformation of the heart, yolk sac, tail, head, spine, an abnormal eye development and pigmentation, resulted in a considerable improvement of the test efficiency.

By adding sublethal parameters, we wanted to explore the potential of the FET test for the evaluation of hospital effluent toxicity. Our conclusions are in agreement with findings of numerous studies which have reported an increase in the FET test sensitivity after addition of sublethal endpoints (Babić *et al.* 2017, Krzykwa *et al.* 2019, Cedron *et al.* 2020). We selected five sublethal effects, which may be easily determined: delayed heartbeat, lack of blood circulation, presence of



edema (pericardial or yolk sac), spinal curvature and pigmentation failures. Tenorio-Chávez *et al.* (2020) investigated treated WW from one hospital using zebrafish embryos and obtained LC<sub>50</sub> value 6.1 %. EC<sub>50</sub> value of sublethal malformations was 2.5 % which means higher sensitivity of sublethal parameters compared to lethal endpoints. The main effects identified were pericardial edema, yolk sac malformation, hypopigmentation, hatching abnormalities, tail and chorda deformation, without fin, chorion and craniofacial malformation. Stelzer *et al.* (2018) observed an increase in embryo test sensitivity of >30 % after addition of three sublethal parameters comprising immobility, formation of edema and nonhatching.

According to our results, when we draw a comparison between FET 48 h and FET 96 h, we can see the increase in efficiency of the method caused by the prolonged exposure time from 48 hpf to 96 hpf. In our study, the LC<sub>50</sub> values of FET 48 h were constantly higher than those of FET 96 h. We are of the opinion that these findings could be related, besides other factors, to the chorion as a possible barrier for some chemicals, mainly substances with very high molecular weight (Braunbeck *et al.* 2015, Sobanska *et al.* 2017). This limit disappears during the hatching period of embryonic development from about 72 hpf to 96 hpf (Kimmel *et al.* 1995). Similar results were presented by Stelzer *et al.* (2018), who, based on analyses of HWW samples, discovered less sensitivity of FET 48 h (LC<sub>50</sub>=56.5 %) compared to FET 96 h (LC<sub>50</sub>=53.5 %).

## Conclusions

The Fish Embryo Acute Toxicity Test is used worldwide for ecotoxicological research as an alternative method to the experiments with adult fish. Considering the lack of scientific data regarding the suitability of *D. rerio* embryos as the testing organisms for HWW

investigation, the main purpose of our study was to make a comparison between zebrafish embryos and two other aquatic organisms commonly used for WW analyses.

The results of our study demonstrated highly variable toxic impacts of WW samples from different hospitals on all the test species. With respect to determining acute toxicity of HWW, the obtained data indicate that the sensitivity of zebrafish embryos is comparable or in some cases higher than the sensitivity of *D. magna* and *A. fischeri* and the FET test may be considered as a sufficiently effective and appropriate method for routine HWW testing.

The prolonged exposure time of the test from 48 hpf to 96 hpf significantly improved the sensitivity of zebrafish embryos in 57 % of the tested samples.

In addition to the lethal endpoints, five sublethal criteria were evaluated as a modification of the FET test. The inclusion of sublethal parameters comprising delayed heartbeat, lack of blood circulation, pericardial and yolk sac edema, spinal curvature and pigmentation failures caused statistically significant increase of the FET test sensitivity ( $p=0.026$ ) in the range of 1-12 %.

Our study highlights the necessity of further validation of the FET test applicability for the assessment of hospital effluents by examination of WW samples from a larger set of hospitals. The choice of a suitable battery of sublethal effects in addition to the lethal endpoints of the FET test may be recommended to extend the method efficiency.

## Conflict of Interest

There is no conflict of interest.

## Acknowledgements

Supported by ERDF/ESF project "International competitiveness of NIPH in research, development and education in alternative toxicological methods" (No. CZ.02.1.01/0.0/0.0/16\_019/0000860).

## References

- ABBAS M, ADIL M, EHTISHAM-UL-HAQUE S, MUNIR B, YAMEEN M, GHAFAR A, SHAR GA, TAHIR MA, IQBAL M: *Vibrio fischeri* bioluminescence inhibition assay for ecotoxicity assessment: A review. *Sci Total Environ* 626: 1295-1309, 2018. <https://doi.org/10.1016/j.scitotenv.2018.01.066>
- BABIĆ S, BARIŠIĆ J, VIŠIĆ H, KLOBUČAR RS, TOPIĆ POPOVIĆ N, STRUNJAK-PEROVIĆ I, ČOŽ-RAKOVAC R, KLOBUČAR G: Embryotoxic and genotoxic effects of sewage effluents in zebrafish embryo using multiple endpoint testing. *Water Res* 115: 9-21, 2017. <https://doi.org/10.1016/j.watres.2017.02.049>
- BOILLOT C, BAZIN C, TISSOT-GUERRAZ F, DROGUET J, PERRAUD M, CETRE JC, TREPO D, PERRODIN Y: Daily physicochemical, microbiological and ecotoxicological fluctuations of a hospital effluent according to technical and care activities. *Sci Total Environ* 403: 113-129, 2008. <https://doi.org/10.1016/j.scitotenv.2008.04.037>



- BRANDHOF EJ, MONTFORTS M: Fish embryo toxicity of carbamazepine, diclofenac and metoprolol. *Ecotoxicol Environ Saf* 73: 1862-1866, 2010. <https://doi.org/10.1016/j.ecoenv.2010.08.031>
- BRAUNBECK T, KAIS B, LAMMER E, OTTE J, SCHNEIDER K, STENGEL D, STRECKER R: The fish embryo test (FET): origin, applications, and future. *Environ Sci Pollut Res* 22: 16247-16261, 2015. <https://doi.org/10.1007/s11356-014-3814-7>
- CEDRON VP, WEINER AMJ, VERA M, SANCHEZ L: Acetaminophen affects the survivor, pigmentation and development of craniofacial structures in zebrafish (*Danio rerio*) embryos. *Biochem Pharmacol* 174: 113816, 2020. <https://doi.org/10.1016/j.bcp.2020.113816>
- CLEUVERS M: Aquatic ecotoxicity of pharmaceuticals including the assessment of combination effects. *Toxicol Lett* 142: 185-194, 2003. [https://doi.org/10.1016/S0378-4274\(03\)00068-7](https://doi.org/10.1016/S0378-4274(03)00068-7)
- CLEUVERS M: Initial risk assessment for three beta-blockers found in the aquatic environment. *Chemosphere* 59: 199-205, 2005. <https://doi.org/10.1016/j.chemosphere.2004.11.090>
- ČSN 75 6406: Wastewater discharge and treatment from health facilities, 2020.
- DANG Z, VAN DER VEN LTM, KIENHUIS AS: Fish embryo toxicity test, threshold approach, and moribund as approaches to implement 3R principles to the acute fish toxicity test. *Chemosphere* 186: 677-685, 2017. <https://doi.org/10.1016/j.chemosphere.2017.08.047>
- DATEL JV, HRABANKOVA A: Pharmaceuticals load in the Svihov water reservoir (Czech Republic) and impacts on quality of treated drinking water. *Water* 12: 1387, 2020. <https://doi.org/10.3390/w12051387>
- ELLEPOLA N, OGAS T, TURNER DN, GURUNG R, MALDONADO-TORRES S, TELLO-ABURTO R, PATIDAR PL, ROGELJ S, PIYASENA ME: A toxicological study on photo-degradation products of environmental ibuprofen: Ecological and human health implications. *Ecotoxicol Environ Saf* 188: 109892, 2020. <https://doi.org/10.1016/j.ecoenv.2019.109892>
- EMMANUEL E, HANNA K, BAZIN C, KECK G, CLÉMENT B, PERRODIN Y: Fate of glutaraldehyde in hospital wastewater and combined effects of glutaraldehyde and surfactants on aquatic organisms. *Environ Int* 31: 399-406, 2005. <https://doi.org/10.1016/j.envint.2004.08.011>
- EUROPEAN PARLIAMENT AND COUNCIL: DIRECTIVE 2000/60/EC of the European Parliament and of the Council establishing a framework for Community action in the field of water policy (EU Water Framework Directive). *Official Journal of the European Union L* 327: 1-73, 2000.
- EUROPEAN PARLIAMENT AND COUNCIL: DIRECTIVE 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes. *Official Journal of the European Union L* 276: 33-79, 2010.
- GODOY AA, OLIVEIRA ÁC, SILVA JGM, JESUS AZEVEDO CC, DOMINGUES I, NOGUEIRA AJA, KUMMROW F: Single and mixture toxicity of four pharmaceuticals of environmental concern to aquatic organisms, including a behavioural assessment. *Chemosphere* 235: 373-382, 2019. <https://doi.org/10.1016/j.chemosphere.2019.06.200>
- HALLING-SØRENSEN B, HOLTEN LÚTZHØFT H-C, ANDERSEN HR, INGERSLEV F: Environmental risk assessment of antibiotics: comparison of mecillinam, trimethoprim and ciprofloxacin. *J Antimicrob Chemother* 46: 53-58, 2000. [https://doi.org/10.1093/jac/46.suppl\\_1.53](https://doi.org/10.1093/jac/46.suppl_1.53)
- HAMJINDA NS, CHIEMCHAI SRI W, WATANABE T, HONDA R, CHIEMCHAI SRI CH: Toxicological assessment of hospital wastewater in different treatment processes. *Environ Sci Pollut Res* 25: 7271-7279, 2015. <https://doi.org/10.1007/s11356-015-4812-0>
- ISO 11348-2: Water quality – Determination of the inhibitory effect of water samples on the light emission of *Vibrio fischeri* (Luminescent bacteria test) - Part 2: Method using liquid-dried bacteria, 2007.
- ISO 15088: Water quality – Determination of the acute toxicity of waste water to zebrafish eggs (*Danio rerio*), 2007.
- ISO 6341: Water quality – Determination of the inhibition of the mobility of *Daphnia magna* Straus (Cladocera, Crustacea) – Acute toxicity test, 2012.
- ISO 7346-2: Water quality – Determination of the acute lethal toxicity of substances to a freshwater fish [*Brachydanio rerio* Hamilton-Buchanan (Teleostei, Cyprinidae)]-Part 2: Semi-static method, 1999.
- ISO 8692: Water quality – Fresh water algal growth inhibition test with unicellular green algae, 2012.

- JIROVA G, VLKOVA A, WITTLEROVA M, DVORAKOVA M, KASPAROVA L, CHRZ J, KEJLOVA K, WITTLINGEROVA Z, ZIMOVA M, HOSIKOVA B, JIRAVOVA J, KOLAROVA H: Toxicity of wastewater from health care facilities assessed by different bioassays. *Neuro Endocrinol Lett* 39: 441-453, 2018.
- JIROVA G, WITTLINGEROVA Z, ZIMOVA M, VLKOVA A, WITTLEROVA M, DVORAKOVA M, JIROVA D: Bioindicators of wastewater ecotoxicity. *Neuro Endocrinol Lett* 37: 17-24, 2016.
- KERN DI, OLIVEIRA SCHWAICKHARDT R, LUTTERBECK CA, KIST LT, ALCAYAGA EAL, MACHADO ÊL: *Arch Environ Contam Toxicol* 68: 64-73, 2015. <https://doi.org/10.1007/s00244-014-0072-0>
- KIMMEL CHB, BALLARD WW, KIMMEL SR, ULLMANN B, SCHILLING TF: Stages of embryonic development of the zebrafish. *Dev Dyn* 203: 253-310, 1995. <https://doi.org/10.1002/aja.1002030302>
- KRZYKWA JC, SAEID A, JEFFRIES MKS: Identifying sublethal endpoints for evaluating neurotoxic compounds utilizing the fish embryo toxicity test. *Ecotoxicol Environ Saf* 170: 521-529, 2019. <https://doi.org/10.1016/j.ecoenv.2018.11.118>
- LAMMER E, CARR GJ, WENDLER K, RAWLINGS JM, BELANGER SE, BRAUNBECK T: Is the fish embryo toxicity test (FET) with the zebrafish (*Danio rerio*) a potential alternative for the fish acute toxicity test? *Comp Biochem Phys C* 149: 196-209, 2009. <https://doi.org/10.1016/j.cbpc.2008.11.006>
- LAQUAZ M, DAGOT CH, BAZIN CH, BASTIDE T, GASCHET M, PLOY M-C, PERRODIN Y: Ecotoxicity and antibiotic resistance of a mixture of hospital and urban sewage in a wastewater treatment plant. *Environ Sci Pollut Res* 25: 9243-9253, 2017. <https://doi.org/10.1007/s11356-017-9957-6>
- LI Q, WANG P, CHEN L, GAO H, WU L: Acute toxicity and histopathological effects of naproxen in zebrafish (*Danio rerio*) early life stages. *Environ Sci Pollut Res* 23: 18832-18841, 2016. <https://doi.org/10.1007/s11356-016-7092-4>
- LI S-W, WANG Y-H, LIN AY-C: Ecotoxicological effect of ketamine: Evidence of acute, chronic and photolysis toxicity to *Daphnia magna*. *Ecotoxicol Environ Saf* 143: 173-179, 2017. <https://doi.org/10.1016/j.ecoenv.2017.05.040>
- MARTINS N, PEREIRA R, ABRANTES N, GONÇALVES F, MARQUES CR: Ecotoxicological effects of ciprofloxacin on freshwater species: data integration and derivation of toxicity thresholds for risk assessment. *Ecotoxicology* 21: 1167-1176, 2012. <https://doi.org/10.1007/s10646-012-0871-x>
- MEZA LC, PIOTROWSKI P, FARNAN J, TASKER TL, XIONG B, WEGGLER B, MURRELL K, DORMAN FL, HEUVEL JPV, BURGOS WD: Detection and removal of biologically active organic micropollutants from hospital wastewater. *Sci Total Environ* 700: 134469, 2020. <https://doi.org/10.1016/j.scitotenv.2019.134469>
- OECD: Test Guideline No. 203: Fish, Acute Toxicity Test. OECD Guidelines for Testing of Chemicals, Section 2, Paris, France, 2019.
- OECD: Test Guideline No. 236: Fish Embryo Acute Toxicity (FET) Test. OECD Guidelines for Testing of Chemicals, Section 2, Paris, France, 2013.
- ORIAS F, PERRODIN Y: Characterisation of the ecotoxicity of hospital effluents: A review. *Sci Total Environ* 454-455: 250-276, 2013. <https://doi.org/10.1016/j.scitotenv.2013.02.064>
- PÉREZ-ALVAREZ I, ISLAS-FLORES H, GÓMEZ-OLIVÁN LM, BARCELÓ D, DE ALDA ML, SOLSONA SP, SÁNCHEZ-ACEVES L, SANJUAN-REYES N, GALAR-MARTÍNEZ M: Determination of metals and pharmaceutical compounds released in hospital wastewater from Toluca, Mexico, and evaluation of their toxic impact. *Environ Pollut* 240: 330-341, 2018. <https://doi.org/10.1016/j.envpol.2018.04.116>
- ROGOWSKA J, CIESZYNSKA-SEMENOWICZ M, RATAJCZYK W, WOLSKA L: Micropollutants in treated wastewater. *Ambio* 49: 487-503, 2020. <https://doi.org/10.1007/s13280-019-01219-5>
- ROMANUCCI V, SICILIANO A, GALDIERO E, GUIDA M, LUONGO G, LIGUORI R, FABIO G, PREVITERA L, ZARRELLI A: Disinfection by-products and ecotoxic risk associated with hypochlorite treatment of tramadol. *Molecules* 24: 693, 2019. <https://doi.org/10.3390/molecules24040693>
- SANTOS LHMLM, ARAÚJO AN, FACHINI A, PENA A, DELERUE-MATOS C, MONTENEGRO MCBSM: Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. *J Hazard Mater* 175: 45-95, 2010. <https://doi.org/10.1016/j.jhazmat.2009.10.100>
- SCHOLZ S, FISCHER S, GÜNDEL U, KÜSTER E, LUCKENBACH T, VOELKER D: The zebrafish embryo model in environmental risk assessment-applications beyond acute toxicity testing. *Environ Sci Pollut Res* 15: 394-404, 2008. <https://doi.org/10.1007/s11356-008-0018-z>

- SOBANSKA M, SCHOLZ S, NYMAN A-M, CESNAITIS R, ALONSO SG, KLÜVER N, KÜHNE R, TYLE H, KNECHT J, DANG Z, LUNDBERGH I, CARLON C, COEN W: Applicability of the Fish Embryo Acute Toxicity (FET) Test (OECD 236) in the Regulatory Context of Registration, Evaluation, Authorisation, and Restriction of Chemicals (REACH). *Environ Toxicol Chem* 37: 657-670, 2017. <https://doi.org/10.1002/etc.4055>
- SOKAL RR, ROHLF FJ: *Biometry: The Principles and Practice of Statistics in Biological Research. 3rd Edition.* W.H. Freeman and Co., New York, 1995, 887 p.
- STELZER JAA, ROSIN CK, BAUER LH, HARTMANN M, PULGATI FH, ARENZON A: Is Fish Embryo Test (FET) According to OECD 236 sensible enough for delivering quality data for effluent risk assessment? *Environ Toxicol Chem* 37: 2925-2932, 2018. <https://doi.org/10.1002/etc.4215>
- TENORIO-CHÁVEZ P, CERRO-LÓPEZ M, CASTRO-PASTRANA LI, RAMÍREZ-RODRIGUES MM, OROZCO-HERNÁNDEZ JM, GÓMEZ-OLIVÁN LM: Effects of effluent from a hospital in Mexico on the embryonic development of zebrafish, *Danio rerio*. *Sci Total Environ* 727: 1-11, 2020. <https://doi.org/10.1016/j.scitotenv.2020.138716>
- VASCONCELOS EC, DALKE CR, OLIVEIRA CMR: Influence of select antibiotics on *Vibrio fischeri* and *Desmodesmus subspicatus* at  $\mu\text{g L}^{-1}$  concentrations. *J Environ Manage* 60: 157-164, 2017. <https://doi.org/10.1007/s00267-017-0841-4>
- VÄLITALO P, MASSEI R, HEISKANEN I, BEHNISCH P, BRACK W, TINDALL AJ, PASQUIER D, KÜSTER E, MIKOLA A, SCHULZE T, SILLANPÄÄ M: Effect-based assessment of toxicity removal during wastewater treatment. *Water Res* 126: 153-163, 2017. <https://doi.org/10.1016/j.watres.2017.09.014>
- VERLICCHI P, AL AUKIDY M, ZAMBELLO E: What have we learned from worldwide experiences on the management and treatment of hospital effluent? - An overview and a discussion on perspectives. *Sci Total Environ* 514: 467-491, 2015. <https://doi.org/10.1016/j.scitotenv.2015.02.020>
- WIEST L, CHONOVA T, BERGÉ A, BAUDOT R, BESSUEILLE-BARBIER F, AYOUNI-DEROUCHE L, VULLIET E: Two-year survey of specific hospital wastewater treatment and its impact on pharmaceutical discharges. *Environ Sci Pollut Res* 25: 9207-9218, 2018. <https://doi.org/10.1007/s11356-017-9662-5>
- WIGH A, DEVAUX A, BROSELIN V, GONZALES-OSPINA A, DOMENJOUR B, AÏT-AÏSSA S, CREUSOT N, GOSSET A, BAZIN CH, BONY S: Proposal to optimize ecotoxicological evaluation of wastewater treated by conventional biological and ozonation processes. *Environ Sci Pollut Res* 23: 3008-3017, 2016. <https://doi.org/10.1007/s11356-015-5419-1>
-