

Kwashiorkor — New evidence in the puzzle of oedema formation



André Briend^{a,b}

^aDepartment of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen, Frederiksberg, Denmark

^bCenter for Child Health Research, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

Kwashiorkor, a type of severe malnutrition associated with oedema, has been known for centuries but its mechanism remains elusive.¹ The low level of plasma albumin and the associated decreased plasma oncotic pressure has been for years the main explanation of oedema of kwashiorkor: an excess fluid flow from the capillaries to the interstitium was assumed to result from an imbalance of Starling's forces.² This mechanism seems plausible, as a low level of plasma albumin is a hallmark of kwashiorkor³ and the Starling's principle shows that plasma oncotic pressure, largely determined by plasma albumin, is amongst the factors influencing fluid flow from the capillaries towards the interstitium.⁴ For years, however, the implication of other mechanisms has been postulated, as there is an imperfect match between albumin levels and the degree of oedema.⁵ Arguably, albumin level is not the only determinant of the plasma oncotic pressure which itself is only one of many variables determining the balance of the Starling's forces and this may explain some of the mismatch between albumin and the degree of oedema. Another explanation could be that the drainage of fluid from the interstitium to the lymphatic system is impaired. It is now accepted that, as a rule, fluid does not flow back from the interstitium to the venous end of the capillaries as initially assumed by Starling,⁶ giving a leading role to the lymphatic system in regulating the volume of interstitial fluid. This important role of the lymphatic system has been a blind spot in the discussion about the mechanisms of oedema formation in kwashiorkor. This is surprising, as it seems that lymphatic drainage has the capacity to increase by a factor of 10 to 50 in case of excess fluid accumulation in the interstitium, providing a strong protection against oedema formation.⁷ It can be argued that a moderate increase of fluid flow from the capillaries to the interstitial fluid cannot lead to oedema without a simultaneous dysfunction of this protecting mechanism.

The paper by Gonzales et al. in recent issue of *eBioMedicine*⁸ gives important clues on oedema formation. First, by comparing children with and without oedema

and children with similar albumin levels but different degrees of oedema, it confirms that low plasma albumin levels are necessary for oedema formation but that it is not the only explanation. Second, it gives some evidence that lymphatic drainage may be impaired, by showing an association between markers of the lymphatic function and oedema. This is an important piece in the puzzle of oedema formation.

Lymphatic drainage is an active and energy-consuming process involving mechanical fluid pumping by lymphatic valves, and thus, unsurprisingly, it is affected in malnutrition. It is not clear why lymphatic drainage is specifically affected in kwashiorkor, but evidence that markers of lymphatic function may be associated with oedema provides a new important insight in oedema formation.

The Gonzales et al. paper also shows evidence of extracellular matrix degradation in kwashiorkor patients. This factor has been suggested to be a possible contributing factor in oedema formation: it has been postulated that the interstitium behaves like a chromatography column, which can collapse in two phases when the chemical structure of the extracellular matrix is altered.⁹ The role of such a mechanism should be taken with caution, however. The physical structure of the interstitium makes it very resistant to compression but easily disrupted when its hydrostatic pressure increases: a free liquid phase appears, leading to oedema, even in absence of any change in the chemical structure of the interstitium, as soon as this pressure becomes positive.⁷

The treatment recommended for kwashiorkor by the World Health Organization is the same as for other forms of severe malnutrition.¹⁰ The new evidence provided by the paper of Gonzales et al. should prompt investigations on factors leading to this alteration of the lymphatic function to develop new approach really adapted to treat this life-threatening condition.

Contributors

André Briend wrote this commissioned commentary.

Declaration of interests

The author declares no competing interests.

**eBioMedicine 2022;80:
104070**
Published online 19 May
2022
[https://doi.org/10.1016/j.
ebiom.2022.104070](https://doi.org/10.1016/j.ebiom.2022.104070)

E-mail address: andre.briend@gmail.com

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

References

- 1 Briend A. Kwashiorkor: still an enigma—the search must go on. *CMAM Forum Technical Brief*. 2014. [Internet][cited 2015 Nov 24]. Available from: <http://fr.cmamforum.org/Pool/Resources/Kwashiorkor,-still-an-enigma-CMAM-Forum-Dec-2014.pdf>.
- 2 Waterlow JC. Kwashiorkor revisited: the pathogenesis of oedema in kwashiorkor and its significance. *Trans R Soc Trop Med Hyg*. 1984;78:436–441.
- 3 Coulthard MG. Oedema in kwashiorkor is caused by hypoalbuminaemia. *Paediatr Int Child Health*. 2015;35:83–89.
- 4 Levick JR. Revision of the Starling principle: new views of tissue fluid balance. *J Physiol*. 2004;557:704.
- 5 Golden MH, Golden BE, Jackson AA. Albumin and nutritional oedema. *Lancet*. 1980;1:114–116.
- 6 Michel CC. Fluid exchange in the microcirculation. *J Physiol*. 2004;557:701–702.
- 7 Guyton Arthur C, Hall JE. *Textbook of Medical Physiology*. 11th edition Philadelphia: Elsevier Saunders; 2006.
- 8 Gonzales Gerard Bryan, Njunge James M, Gichuki Bonface M, et al. The role of albumin and the extracellular matrix on the pathophysiology of oedema formation in severe malnutrition. *EBioMedicine*. 2022;7:(79) 103991.
- 9 Golden Michael H. Nutritional and other types of oedema, albumin, complex carbohydrates and the interstitium - a response to Malcolm Coulthard's hypothesis: oedema in kwashiorkor is caused by hypoalbuminaemia. *Paediatr Int Child Health*. 2015. 2046905515Y000000010.
- 10 World Health Organization. *Pocket Book of Hospital Care For Children*. Geneva: World Health Organization; 2013.