




ELSEVIER

Contents lists available at ScienceDirect

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laa

Some further applications of a lattice theoretic method in the study of singular LCM matrices

Mika Mattila^{a,*}, Pentti Haukkanen^a, Jori Mäntysalo^b^a Faculty of Information Technology and Communication Sciences, Tampere University, Finland^b IT services, Tampere University, Finland

ARTICLE INFO

Article history:

Received 15 December 2022

Accepted 26 April 2023

Available online xxxx

Submitted by P. Semrl

MSC:

11C20

15B36

06A07

Keywords:

Bourque-Ligh conjecture

LCM matrix

GCD matrix

Smith determinant

ABSTRACT

In 1876 H. J. S. Smith defined an LCM matrix as follows: let $S = \{x_1, x_2, \dots, x_n\}$ be a set of positive integers with $x_1 < x_2 < \dots < x_n$. The LCM matrix $[S]$ on the set S is the $n \times n$ matrix with $\text{lcm}(x_i, x_j)$ as its ij entry. During the last 30 years singularity of LCM matrices has interested many authors. In 1992 Bourque and Ligh ended up conjecturing that if the GCD closedness of the set S (which means that $\text{gcd}(x_i, x_j) \in S$ for all $i, j \in \{1, 2, \dots, n\}$), suffices to guarantee the invertibility of the matrix $[S]$. However, a few years later this conjecture was proven false first by Haukkanen et al. and then by Hong. It turned out that the conjecture holds only on GCD closed sets with at most 7 elements but not in general for larger sets. However, the given counterexamples did not give much insight on why does the conjecture fail exactly in the case when $n = 8$. This situation was later improved in a couple of articles, where a new lattice theoretic approach was introduced (the method is based on the fact that because the set S is assumed to be GCD closed, the structure $(S, |)$ actually forms a meet semilattice). For example, it has been shown that in the case when the set S has 8 elements and the matrix $[S]$ is singular, there is only one option for the semilattice structure of $(S, |)$, namely the cube structure.

Since the cases $n \leq 8$ have been thoroughly studied in various articles, the next natural step is to apply the methods to

* Corresponding author.

E-mail addresses: mika.mattila@tuni.fi (M. Mattila), pentti.haukkanen@tuni.fi (P. Haukkanen), jori.mantysalo@tuni.fi (J. Mäntysalo).<https://doi.org/10.1016/j.laa.2023.04.027>0024-3795/© 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

the case $n = 9$. This was done by Altınışık and Altıntaş as they consider the different lattice structures of $(S, |)$ with nine elements that can result in a singular LCM matrix $[S]$. However, their investigation leaves two open questions, and the main purpose of this presentation is to provide solutions to them. We shall also give a new lattice theoretic proof for a result referred to as Sun's conjecture, which was originally proven by Hong via number theoretic approach.

© 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The concept of an LCM matrix, as well as the concept of a GCD matrix, was originally defined by H. J. S. Smith [12] in his seminal paper from the year 1876. Assuming the set $S = \{x_1, x_2, \dots, x_n\}$ to be a finite subset of \mathbb{Z}^+ with distinct elements, Smith defined the GCD matrix (S) of the set S to be the $n \times n$ matrix with $\gcd(x_i, x_j)$ as its ij element. Similarly, the LCM matrix $[S]$ of the set S was defined to be the $n \times n$ matrix with $\text{lcm}(x_i, x_j)$ as its ij element. Besides calculating determinant formulas for several GCD and LCM type matrices, Smith was also considered the invertibility of GCD and LCM matrices. For example, he showed that if the set S is factor closed (i.e. the implication $(y | x \text{ for some } x \in S) \Rightarrow y \in S$ holds for all $y \in \mathbb{Z}^+$), then both of the matrices (S) and $[S]$ are invertible.

Surprisingly it took more than 110 years until LCM matrices got as get much attention in mathematical literature. Although GCD-type matrices had been studied in several papers over the years, it was not until 1992 that LCM matrices were reintroduced by Bourque and Ligh [2]. Among other things, they showed that it is actually quite easy to find singular LCM matrices by considering the LCM matrix of the four element set $S = \{1, 2, 15, 42\}$ (see [2, p. 68]). The authors then turned the attention to finding conditions less restricting than factor closedness of the set S that would suffice to guarantee the invertibility of the matrix $[S]$. They ended up conjecturing that the GCD-closedness of the set S suffices to guarantee the invertibility of $[S]$.

In 1997 Haukkanen et al. [6] managed to find a GCD closed set S with 9 elements whose LCM matrix is singular, which disproved the Bourque-Ligh conjecture and showed that GCD closedness of the set S does not actually suffice to guarantee the invertibility of the matrix $[S]$. Two years later Hong [7] was able to find another similar counterexample in which there was only 8 elements in the set S . By using number theoretic methods Hong [10] also showed that the conjecture holds for GCD closed sets with at most 7 elements and that it does not hold in general for larger sets, which in some sense meant that the conjecture was solved completely. However, if we assume the set S is GCD closed, it also means that the structure $(S, |)$ itself forms a meet semilattice, which makes it possible to study the conjecture from an entirely lattice theoretic point of view. In [11] Korkee et al. showed (via investigation of all possible semilattice structures with at most 7 elements) that the LCM matrix $[S]$ is invertible for any GCD closed set S with $|S| \leq 7$. In [4]

this same lattice theoretic approach is utilized to show that if the matrix $[S]$ is singular and the set S is GCD closed with 8 elements, then $(S, |)$ has unique, cube-like structure. These same methods were also adapted by Altınışık et al. in [1], where they study the singularity of the matrix $[S]$ in the case when S is a GCD closed set with 9 elements. Finally, in 2020 Haukkanen et al. [5] refined the lattice theoretic method by defining a property that was in common for all meet semilattices with at most 8 elements that sufficed to guarantee the invertibility of the corresponding LCM matrix $[S]$. At the same time it also turned out that in many cases the lattice theoretic structure of $(S, |)$ not only guarantees the invertibility but also completely determines the inertia of the matrix $[S]$.

In the present article we continue utilizing our lattice theoretic method and find a couple of new ways to apply it. In Section 3 we reconsider the work done by Altınışık and Altıntaş in [1] from a slightly different perspective and we classify all possible 9 element semilattice structures for which there exists a gcd closed set of this type such that the corresponding LCM matrix is singular. In this context we also solve one open conjecture about this 9 element case raised in their paper. In Section 4 we turn our attention to another conjecture raised by Altınışık and Altıntaş and disprove it by presenting a couple of counterexamples. In Section 5 we give a novel lattice theoretic proof for a result known as Sun's conjecture, and finally in Section 6 we give an example on how our method can give some useful information in the study of the so-called power LCM matrices as well.

2. Preliminaries

First we need to introduce a couple of notations and concepts developed in [5] and bring them to our lattice theoretic context. Let $S = \{x_1, x_2, \dots, x_n\}$ be a GCD closed set with $x_i \leq x_j \Rightarrow i \leq j$. Let us denote

$$C_S(x) = \{y \in S \mid y | x \text{ and for all } z \in S : (y | z \text{ and } z | x \Rightarrow y = z)\}$$

and

$$\text{meetcl}_S(C_S(x)) = \{\text{gcd}(y_1, y_2, \dots, y_k) \mid k \in \mathbb{Z}^+ \text{ and } y_1, y_2, \dots, y_k \in C_S(x)\}.$$

In other words, $C_S(x)$ is the set of all elements of the set S that are covered by x in the meet semilattice $(S, |)$ and $\text{meetcl}_S(C_S(x))$ is the smallest GCD closed subset of S that contains all the elements of the set $C_S(x)$.

Definition 2.1 (cf. [5], Definition 2.2). An element $x \in S$ generates a double-chain set in S if the set $\text{meetcl}_S(C_S(x)) \setminus C_S(x)$ can be expressed as a union of two disjoint sets A and B that are chains in $(S, |)$.

By the simple fact that $\text{gcd}(x_i, x_j)\text{lcm}(x_i, x_j) = x_i x_j$ we may decompose the matrix $[S]$ as

$$[S] = \text{diag}(x_1, x_2, \dots, x_n) \left(\frac{1}{\gcd(x_i, x_j)} \right) \text{diag}(x_1, x_2, \dots, x_n).$$

Next we use Möbius inversion and define the function Ψ_S on S as

$$\Psi_S(x_i) = \sum_{x_j | x_i} \frac{\mu_S(x_j, x_i)}{x_j}, \tag{2.1}$$

where the Möbius function values $\mu_S(x_j, x_i)$ are defined by using the recursive formula

$$\begin{aligned} \mu_S(x_i, x_i) &= 1, \\ \mu_S(x_j, x_i) &= - \sum_{x_j \prec x_k \preceq x_i} \mu_S(x_k, x_i) = - \sum_{x_j \preceq x_k \prec x_i} \mu_S(x_j, x_k). \end{aligned}$$

Now the matrix $\left(\frac{1}{\gcd(x_i, x_j)} \right)$ may be written as

$$\left(\frac{1}{\gcd(x_i, x_j)} \right) = E \text{diag}(\Psi_S(x_1), \Psi_S(x_2), \dots, \Psi_S(x_n)) E^T,$$

where $E = (e_{ij})$ is the 0, 1 incidence matrix of the set S with

$$e_{ij} = \begin{cases} 1 & \text{if } x_j | x_i, \\ 0 & \text{otherwise.} \end{cases}$$

Putting all together we have

$$[S] = \Delta E \Lambda E^T \Delta = (\Delta E) \Lambda (\Delta E)^T,$$

where $\Delta = \text{diag}(x_1, x_2, \dots, x_n)$ and $\Lambda = \text{diag}(\Psi_S(x_1), \Psi_S(x_2), \dots, \Psi_S(x_n))$.

Since the matrix ΔE is clearly invertible (triangular matrix with nonzero diagonal elements), the matrix $[S]$ is invertible if and only if the matrix Λ is invertible. Moreover,

$$\det \Lambda = \Psi_S(x_1) \Psi_S(x_2) \cdots \Psi_S(x_n).$$

From this we easily obtain the following fundamental result.

Proposition 2.1. *If the set S is GCD closed, then the LCM matrix $[S]$ is invertible if and only if $\Psi_S(x_i) \neq 0$ for all $i = 1, 2, \dots, n$.*

The following theorem is one of the key results of [5] and it plays a pivotal role in this article as well.

Theorem 2.1 (cf. [5], Theorem 4.1). *Let S be a GCD closed set. If the element $x_i \in S$ generates a double-chain set in S , then $\Psi_S(x_i) \neq 0$.*

The following result is important to keep in mind when one tries to construct a set S whose LCM matrix $[S]$ is singular.

Theorem 2.2. *Suppose that the element $x_i \in S$ and that there is at least one element in $C_S(x_i)$. If $x_i > \text{lcm}(C_S(x_i))$, then $\Psi_S(x_i) \neq 0$.*

Proof. Suppose that $x_i = a(\text{lcm}(C_S(x_i)))$, where $a \in \mathbb{Z}$ and $a > 1$. Then

$$\begin{aligned} \Psi_S(x_i) &= \sum_{x_j \preceq x_i} \frac{\mu_S(x_j, x_i)}{x_j} \\ &= \frac{1}{\text{lcm}(C_S(x_i))} \left(\frac{\text{lcm}(C_S(x_i))}{x_i} + \sum_{x_j \prec x_i} \frac{\text{lcm}(C_S(x_i))}{x_j} \mu_S(x_j, x_i) \right) \\ &= \frac{1}{\text{lcm}(C_S(x_i))} \left(\underbrace{\frac{1}{a}}_{\notin \mathbb{Z}} + \sum_{x_j \prec x_i} \underbrace{\frac{\text{lcm}(C_S(x_i))}{x_j}}_{\in \mathbb{Z}} \underbrace{\mu_S(x_j, x_i)}_{\in \mathbb{Z}} \right) \neq 0. \quad \square \end{aligned}$$

3. Solving the 9 element case completely

The method used by Altınışık and Altıntaş in [1] is based on constructing a GCD closed set $S_8 = \{x_1, x_2, \dots, x_8\}$ for which the matrix $[S_8]$ is singular. They then go through all the different ways to insert a new positive integer a in the set so that the set $S_9 := S_8 \cup \{a\}$ would still remain GCD closed. If this turns out to be possible, the LCM matrix $[S_9]$ is also guaranteed to remain singular as it is similar to a matrix that has one of its leading principal minors equal to zero. Unfortunately, however, this method does not find the singular LCM matrices of order 9 whose every proper principal submatrix is nonsingular. This means that if we are interested in finding all possible semilattice structures with 9 elements that correspond to some singular LCM matrix $[S]$, the method of Altınışık and Altıntaş only gives a partial solution and cannot find all of them.

There are 5994 different meet semilattice structures with 9 elements (in comparison to only 1078 meet semilattice structures with 8 elements and 37622 structures with 10 elements). Since we are only interested in those structures that can be used to produce singular LCM matrices, Theorem 2.1 and Proposition 2.1 allow us to sieve out all the semilattice structures where every element generates a double-chain set. We are going to utilize program Sage in order to achieve this. Very slow but working way to generate all semilattices is

```
SLall = [MeetSemilattice(P) for P in Posets(9) if P.is_meet_semilattice()]
And then we select only those possessing an element e such that it covers at least three elements, and the subposet of the meet semilattice generated by those elements with generating elements removed has width greater than or equal to three:
```




Fig. 1. The 13 semilattice structures that need to be studied separately.

```

SLspecial = []
for SL in SLall:
    for e in SL:
        if len(SL.lower_covers(e)) >= 3:
            elms = SL.lower_covers(e)
            SL1 = SL.submeetsemilattice(elms)
            P = SL1.subposet([e for e in SL1 if e not in elms])
            if P.width() >= 3:
                Lspecial.append(L)
                break
    
```

This leaves us with 13 semilattice structures shown in Fig. 1 that we need to consider separately.

3.1. Inserting a new maximal element

Let us consider the first four structures 9_A , 9_B , 9_C and 9_D . These four structures can also be obtained by using the methods in [1], but by using a slightly more general approach to construct them.

- Start from an 8 element set S_8 whose LCM matrix $[S_8]$ is singular. We may use, e.g., the set $S_8 = \{1, 2, 3, 5, 66, 70, 255, 39270\}$ illustrated in Fig. 2 (see also [5], Example 5.3). The figure also displays the indexing of each element $x_i \in S_8$.




Fig. 2. An example of a GCD closed set with eight elements such that the corresponding LCM matrix is singular.

- Choose an integer a such that $\gcd(x_i, a) = 1$ for all $x_i \in S_8$. For example in the case of the above mentioned set we may choose $a = 13$.
- In the case 9_A define $x_9 = ax_1$ and $S_9 = S_8 \cup \{x_9\}$.
- In the case 9_B define $x_9 = ax_4$ and $S_9 = S_8 \cup \{x_9\}$.
- In the case 9_C define $x_9 = ax_7$ and $S_9 = S_8 \cup \{x_9\}$.
- In the case 9_D define $x_9 = ax_8$ and $S_9 = S_8 \cup \{x_9\}$.
- In any case we have $\mu_{S_8}(x_i, x_8) = \mu_{S_9}(x_i, x_8)$ for all $i = 1, 2, \dots, 8$ and $x_9 \nmid x_8$, which implies that $\mu_{S_8}(x_9, x_8) = 0$. Thus $\Psi_{S_9}(x_8) = \Psi_{S_8}(x_8) = 0$ and we may conclude that the matrix $[S_9]$ is singular.

3.2. Inserting a new minimum element

Let us now consider the semilattice structure 9_E . This kind of approach was introduced by Hong [7] in 1999 as it enables one to construct a singular LCM matrix of arbitrary size $n \geq 9$. Also the method used by Altınışık and Altıntaş finds this semilattice structure possible. As we did earlier, we are going to address this special case from our lattice theoretic point of view.

- Again we start from any 8 element set S_8 for which the matrix $[S_8]$ is singular.
- Choose an arbitrary integer $a > 1$.
- Define $x'_1 = a, x'_{i+1} = ax_i$ for all $i = 1, 2, \dots, 8$ and $S_9 = \{x'_1, x'_2, \dots, x'_8, x'_9\}$
- We have $\mu_{S_8}(x_i, x_8) = \mu_{S_9}(x'_{i+1}, x'_9)$ for all $i = 1, 2, \dots, 8$ and $\mu_{S_9}(x'_1, x'_9) = 0$, which implies that $\Psi_{S_9}(x_9) = \frac{1}{a}\Psi_{S_8}(x_8) = 0$. Hence the matrix $[S_9]$ is singular.

3.3. Inserting a new element between two comparable elements

Let us now focus on the semilattices $9_F, 9_G$ and 9_H . The method of Altınışık and Altıntaş finds all three structures possible. We shall again provide a slightly more general solution.

- Start from any 8 element set S_8 such that $x_1 = 1$ and the matrix $[S_8]$ is singular.
- Index the elements so that x_5 is an upper bound for x_2 and x_3 .

- It can be shown that $\text{lcm}(x_2, x_3) \mid x_5$, but $x_5 > \text{lcm}(x_2, x_3)$, see e.g. [1, Theorem 2.4].
- Let us denote $x_5 = a\text{lcm}(x_2, x_3)$, where $a > 1$.
- In many (possibly in all) known examples the number a turns out to be a prime number that does not divide either x_2 or x_3 , which means that $\text{gcd}(a, x_i) = 1$ for all $x_i \in S_8 \setminus \{x_8\}$. In the unlikely event when it would not be possible to apply this method to the element x_5 we can always try again with the elements x_6 and x_7 and redraw the lattice by putting either of these elements on the left side of the cube. In the example presented in Fig. 2 this method can be applied to any of the elements x_5, x_6 or x_7 since we have $x_5 = 66 = 11 \cdot \text{lcm}(2, 3)$, $x_6 = 70 = 7 \cdot \text{lcm}(2, 5)$ and $x_7 = 255 = 17 \cdot \text{lcm}(3, 5)$.
- In the case 9_F define $x'_1 = x_1 = 1$, $x'_2 = a$ and $x'_i = x_{i-1}$ for $i = 3, 4, \dots, 8, 9$. We have $\mu_{S'}(x'_2, x'_9) = 0$ and therefore $\Psi_{S_9}(x'_9) = \Psi_{S_8}(x_8) = 0$ and thus $[S_9]$ is singular.
- In the case 9_G define $x'_i = x_i$ for $i = 1, 2, 3, 4$, $x'_5 = ax_2$ and $x'_i = x_{i-1}$ for $i = 6, 7, 8, 9$. Also in this case $\mu_{S_9}(x'_5, x'_9) = 0$ and furthermore $\Psi_{S_9}(x'_9) = \Psi_{S_8}(x_8) = 0$ making the matrix $[S_9]$ is singular.
- In the case 9_H define $x'_i = x_i$ for $i = 1, 2, 3, 4, 6, 7$, $x'_5 = \text{lcm}(x_2, x_3)$, $x'_8 = x_5 = a\text{lcm}(x_2, x_3)$ and $x'_9 = x_8$. Again $\mu_{S_9}(x'_5, x'_9) = 0$, $\Psi_{S_9}(x'_9) = \Psi_{S_8}(x_8) = 0$ and the matrix $[S_9]$ is singular.

3.4. Two irreducible cases

Let us now take the semilattices 9_I and 9_J into consideration.

- With the method used by Altınışık and Altıntaş it is impossible to obtain these two structures.
- Nevertheless, both of these structures are actually possible although they cannot be reduced to the 8 element case.
- In 1997 Haukkanen, Wang and Sillanpää [6] disproved the Bourque-Ligh conjecture by providing the counterexample $S = \{1, 2, 3, 4, 5, 6, 10, 45, 180\}$, which belongs to the class 9_I .
- Also the structure 9_J is possible, for example the set

$$S = \{1, 5, 11, 17, 19, 748, 1463, 2907, 4476780\}$$

is of this type and its LCM matrix is singular. This example is illustrated in Fig. 3. The singularity of the matrix $[S_9]$ follows from the calculation

$$\Psi_{S_9}(x_9) = \frac{1}{4476780} - \frac{1}{2907} - \frac{1}{1463} - \frac{1}{748} - \frac{1}{255} + \frac{1}{19} + \frac{1}{17} + \frac{1}{11} + \frac{0}{1} = 0.$$




Fig. 3. An example of a GCD closed set belonging to the category 9_J such that the corresponding LCM matrix is singular.

3.5. One negative example of inserting a new element into the middle

We shall now focus on the structure 9_K . It would essentially belong to the earlier category “inserting a new element between two comparable elements”, but there are good reasons why we should study it separately.

- Altınışık and Altıntaş attempted to find an example that would belong to this class, but they were not successful. They ended up conjecturing that this structure is not possible at all, see [1, Conjecture 4.1].
- Since $\mu_{S_9}(x_2, x_9) = 0$, there appears to be no difference between this and the corresponding 8 element case.
- However, it turns out that the existence of the element x_2 changes the situation quite drastically.
- Without loss of generality we may assume that $x_1 = 1$ and that the elements x_2 and x_3 are named as shown in Fig. 4. Because we have $x_1 | x_2 | x_3$, the element x_3 cannot be a prime number. We have

$$\Psi_{S_9}(x_9) = \underbrace{\frac{1}{x_9} - \frac{1}{x_8} - \frac{1}{x_7} - \frac{1}{x_6}}_{<0} + \underbrace{\frac{1}{x_5} + \frac{1}{x_4} + \frac{1}{x_3} - \frac{1}{x_1}}_{:=K < 0} < 0,$$

since

$$K \leq -1 + \max\left(\frac{1}{2} + \frac{1}{3} + \frac{1}{25}, \frac{1}{4} + \frac{1}{3} + \frac{1}{5}, \frac{1}{2} + \frac{1}{9} + \frac{1}{5}\right) < 0.$$

This shows that the class 9_K is not possible, which also solves the Conjecture 4.1 by Altınışık and Altıntaş (the fact that x_3 cannot be a prime number also prevents the LCM matrix of the set $S'_8 := S_9 \setminus \{x_2\}$ to be singular since the values $\Psi_{S_9}(x_9)$ and $\Psi_{S'_8}(x'_8)$ are exactly the same).

3.6. Two more impossible cases

Finally, we are left with the structures 9_L and 9_M shown in Fig. 5.




Fig. 4. The structure 9_K , where the different elements x_i have been specified.




Fig. 5. Illustrations of the structures 9_L and 9_M , where the choices for different elements x_i are also displayed.

- In both of these structures every element except for the maximum element x_9 generates a double-chain set, which means that we only need to check the value of $\Psi_S(x_9)$.
- In both cases we have

$$\begin{aligned} \Psi_S(x_9) &= \frac{1}{x_9} - \frac{1}{x_8} - \frac{1}{x_7} - \frac{1}{x_6} - \frac{1}{x_5} + \frac{1}{x_4} + \frac{1}{x_3} + \frac{1}{x_2} + \frac{0}{x_1} \\ &= \underbrace{\frac{1}{x_9}}_{>0} + \underbrace{\left(\frac{1}{x_4} - \frac{1}{x_7} - \frac{1}{x_8}\right)}_{>0} + \underbrace{\left(\frac{1}{x_3} - \frac{1}{x_6}\right)}_{>0} + \underbrace{\left(\frac{1}{x_2} - \frac{1}{x_5}\right)}_{>0} > 0, \end{aligned}$$

which implies that the matrix $[S_9]$ has to be nonsingular.

3.7. Some conclusions

We were able to completely solve the 9 element case and show that there are exactly 10 possible semilattice structures that can be used to construct a GCD closed 9-element set whose LCM matrix is singular. Based on this investigation we can also say that if S is a GCD closed set with 9 elements and the matrix $[S]$ is singular, then the semilattice $(S, |)$ has the cube as its subsemilattice. Altınışık and Altıntaş actually conjecture that this is the case with all sets S whose LCM matrix is singular, and we shall consider this conjecture in the next section. In theory it would now be possible to proceed to the next case and try to find all possible semilattice structures with 10 elements such that there is a corresponding set of positive integers whose LCM matrix is singular. However, as




Fig. 6. Illustrations of the three semilattice structure S_{13} of Example 4.1. The choices for different elements x_i are also shown in the picture.

mentioned earlier there are 37622 meet semilattice structures with 10 elements and it turns out that there are 166 structures where at least one element does not generate a double-chain set within the structure. In other words, solving the 10 element case would require us to either do a huge amount of tedious calculations or to develop more sophisticated methods than we currently have at our disposal.

4. A couple of interesting examples with more than 9 elements

In many cases the 8 element cube semilattice can be found as a subsemilattice of the semilattice $(S, |)$ when the LCM matrix is singular, and this may even happen in quite unexpected ways. We shall illustrate this with a couple of examples.

Example 4.1. Let us consider the GCD closed sets

$$S_{13} = \{1, 2, 3, 13, 23, 25, 41, 75, 369, 533, 6877, 16675, 3679538850\},$$

$$S_{14} = \{1, 2, 3, 6, 7, 11, 13, 19, 56, 147, 209, 1859, 196859, 33105384312\}$$

and

$$S_{16} = \{1, 2, 3, 5, 7, 14, 20, 35, 54, 231, 255, 1820, 45738, 137445, 39308760, 3029801294520\}.$$

The Hasse diagrams of these three semilattices and also the indexing of different elements x_i in each case are shown in Figs. 6, 7 and 8.

The matrices $[S_{13}]$, $[S_{14}]$ and $[S_{16}]$ are all singular since we have

$$\Psi_{S_{13}}(x_{13}) = \frac{1}{3679538850} - \frac{1}{16675} - \frac{1}{6877} - \frac{1}{533} - \frac{1}{369} - \frac{1}{75} - \frac{1}{2} + \frac{1}{41} + \frac{1}{25} + \frac{1}{23} + \frac{1}{13} + \frac{1}{3} + \frac{0}{1} = 0,$$




Fig. 7. Illustrations of the semilattice structure S_{14} considered in Example 4.1. The choices for different elements x_i are also shown in the picture.




Fig. 8. Illustration of the semilattice structure S_{16} of Example 4.1. The choices for different elements x_i are also shown in the picture.

$$\begin{aligned} \Psi_{S_{14}}(x_{14}) &= \frac{1}{33105384312} - \frac{1}{196859} - \frac{1}{1859} - \frac{1}{209} - \frac{1}{147} - \frac{1}{56} - \frac{1}{6} \\ &\quad + \frac{1}{19} + \frac{1}{13} + \frac{1}{11} + \frac{1}{7} + \frac{1}{3} + \frac{1}{2} - \frac{1}{1} = 0 \end{aligned}$$

and

$$\begin{aligned} \Psi_{S_{16}}(x_{16}) &= \frac{1}{3029801294520} - \frac{1}{39308760} - \frac{1}{137445} - \frac{1}{45738} - \frac{1}{1820} \\ &\quad + \frac{1}{255} + \frac{1}{231} + \frac{1}{54} + \frac{1}{35} + \frac{1}{20} + \frac{1}{14} - \frac{1}{7} - \frac{1}{5} - \frac{1}{3} - \frac{1}{2} + \frac{1}{1} = 0. \end{aligned}$$

First we should observe that the set S_{13} does not have the 8 element cube semilattice as its subsemilattice. This implies that Conjecture 4.2 by Altınışık and Altıntaş [1] does not hold in general for GCD closed sets that have at least 13 elements (the conjecture remains still open in the cases when there are exactly 10, 11 or 12 elements in the set S). On the other hand, in the sets S_{14} and S_{16} there are actually two different cube-type subsemilattices. In S_{14} these two subsemilattices have the top and bottom elements in common whereas in S_{16} the two subsemilattices are totally disjoint. One can only say that when it comes to GCD closed sets whose LCM matrix is singular, the semilattice structure of $(S, |)$ can sometimes take quite surprising forms.

5. Lattice theoretic approach to Sun's conjecture

According to Hong [9], in January 1997 (just before the Bouque-Ligh conjecture was solved) professor Qi Sun at Sichuan University had a private conversation with Hong and conjectured that if S is a GCD closed set satisfying $\max_{x_i \in S} \{\omega(x_i)\} \leq 2$, then the LCM matrix $[S]$ is nonsingular (ω is the arithmetic function that counts the number of distinct prime divisors of the input element x_i). In 2005 the conjecture was proven right by Hong in the above mentioned article. This number theoretic proof is quite technical and requires the use of several auxiliary results. We are now going to apply our lattice theoretic method once again and give a new proof which is rather short, visually intuitive and easy to understand.

Theorem 5.1 (*Sun's conjecture*). *Let $S = \{x_1, x_2, \dots, x_n\}$ be a GCD closed set, where every element $x_i \in S$ has at most two distinct prime divisors. Then the LCM matrix $[S]$ is invertible.*

Proof. We only need to show that $\Psi_S(x_i) \neq 0$ for all $x_i \in S$. For the element x_1 we have $\Psi_S(x_1) = \frac{1}{x_1} > 0$. If x_i covers exactly one element in S (namely x_j), then we have

$$\Psi_S(x_i) = \frac{1}{x_i} - \frac{1}{x_j} < 0$$

Suppose then that x_i covers at least two elements in S . Because $\max_{x_i \in S} \{\omega(x_i)\} \leq 2$, all the divisors of x_i are of the form $p^k q^l$, where $p, q \in \mathbb{P}$. Let $p^{a_1} q^{b_1}, p^{a_2} q^{b_2}, \dots, p^{a_r} q^{b_r}$ be the elements that are covered by x_i in S . Without the loss of generality we may assume that $a_1 < a_2 < \dots < a_r$. Because the elements $p^{a_k} q^{b_k}$ are incomparable, we must now have $b_1 > b_2 > \dots > b_r$. Since S is GCD closed, for any $k < l$ we have $\gcd(p^{a_k} q^{b_k}, p^{a_l} q^{b_l}) = p^{a_k} q^{b_l} \in S$. It should be noted that there may exist also other elements of type $p^a q^b$ in S such that $p^a q^b | x_i$ and we have either $a_k < a < a_{k+1}$ for some $k \in \{1, \dots, r-1\}$, $b_{l+1} < b < b_l$ for some $l \in \{1, \dots, r-1\}$ or $a < a_1$ and $b < b_r$. However, in these possible cases we have $\mu_S(p^a q^b, x_i) = 0$ by [4, Lemma 3.2] and therefore these types of elements do not produce any nonzero terms to the function $\Psi_S(x_i)$ and also the




Fig. 9. Illustration of the structure of the set $\text{meetcl}(C_S(x_i)) \cup \{x_i\}$ that we use in the proof of Sun's conjecture.




Fig. 10. A figure that gives the Möbius function values $\mu_S(p^{a_k} q^{b_k}, x_i)$ for different elements $p^{a_k} q^{b_k} \in \text{meetcl}(C_S(x_i)) \cup \{x_i\}$.

existence of such elements does not affect on the calculation of the values $\mu_S(p^{a_k} q^{b_l}, x_i)$, where $k, l \in \{1, \dots, r\}$.

The structure of $\text{meetcl}(C_S(x_i)) \cup \{x_i\}$ is illustrated in Fig. 9. We also need to calculate the values $\mu_S(p^{a_k} q^{b_k}, x_i)$ of the Möbius function on the set S , which is done in Fig. 10. We now obtain

$$\begin{aligned} \Psi_S(x_i) &= \sum_{x_k \in S} \frac{\mu_S(x_k, x_i)}{x_k} = \frac{1}{x_i} - \sum_{j=1}^r \frac{1}{p^{a_j} q^{b_j}} + \sum_{j=1}^{r-1} \frac{1}{p^{a_j} q^{b_{j+1}}} \\ &= \frac{1}{x_i} + \left(\frac{1}{p^{a_1} q^{b_2}} - \frac{1}{p^{a_1} q^{b_1}} - \frac{1}{p^{a_2} q^{b_2}} \right) + \sum_{j=3}^{r-1} \left(\frac{1}{p^{a_{j-1}} q^{b_j}} - \frac{1}{p^{a_j} q^{b_j}} \right) \end{aligned}$$

$$= \frac{1}{x_i} + \frac{1}{p^{a_2} q^{b_1}} \underbrace{(p^{a_2-a_1} q^{b_1-b_2} - p^{a_2-a_1} - q^{b_1-b_2})}_{=p^{a_2-a_1}(q^{b_1-b_2}-1)-q^{b_1-b_2}>0} + \sum_{j=3}^{r-1} \frac{1}{p^{a_j} q^{b_j}} \underbrace{(p^{a_j-a_{j-1}} - 1)}_{>0} > 0.$$

Thus our proof is complete. □

6. Extending the lattice theoretic method in order to study power LCM matrices

The LCM matrix of the set S with respect to the arithmetical function N^α , where $N^\alpha(n) = n^\alpha$ for all $n \in \mathbb{Z}^+$, is the $n \times n$ matrix $[S]_{N^\alpha}$ with $(\text{lcm}(x_i, x_j))^\alpha$ as its ij entry. These type of matrices are known as power LCM matrices. If we wish to study the invertibility of the matrix $[S]_{N^\alpha}$ under the assumption that the set S is GCD closed, we may use a similar approach as we did with the usual LCM matrices. By making use of the identity $(\text{gcd}(x_i, x_j))^\alpha (\text{lcm}(x_i, x_j))^\alpha = x_i^\alpha x_j^\alpha$ we may decompose the matrix $[S]_{N^\alpha}$ as

$$[S]_{N^\alpha} = \text{diag}(x_1^\alpha, x_2^\alpha, \dots, x_n^\alpha) \left(\frac{1}{(\text{gcd}(x_i, x_j))^\alpha} \right) \text{diag}(x_1^\alpha, x_2^\alpha, \dots, x_n^\alpha).$$

Next we use Möbius inversion in exactly the same way and define the function $\Psi_{S, \frac{1}{N^\alpha}}$ on S as

$$\Psi_{S, \frac{1}{N^\alpha}}(x_i) = \sum_{x_j | x_i} \frac{\mu_S(x_j, x_i)}{x_j^\alpha}. \tag{6.2}$$

It should be noted that for the special case $\alpha = 1$ we have $\Psi_{S, \frac{1}{N^\alpha}} = \Psi_S$.

Now the so-called reciprocal power GCD matrix $\left(\frac{1}{(\text{gcd}(x_i, x_j))^\alpha} \right)$ may be written as

$$\left(\frac{1}{(\text{gcd}(x_i, x_j))^\alpha} \right) = E \text{diag}(\Psi_{S, \frac{1}{N^\alpha}}(x_1), \Psi_{S, \frac{1}{N^\alpha}}(x_2), \dots, \Psi_{S, \frac{1}{N^\alpha}}(x_n)) E^T,$$

where $E = (e_{ij})$ is the 0, 1 incidence matrix defined in Section 2. Thereby we obtain the factorization

$$[S]_{N^\alpha} = \Delta^\alpha E \Lambda E^T \Delta^\alpha = (\Delta^\alpha E) \Lambda (\Delta^\alpha E)^T,$$

where $\Delta^\alpha = \text{diag}(x_1^\alpha, x_2^\alpha, \dots, x_n^\alpha)$ and

$$\Lambda = \text{diag}(\Psi_{S, \frac{1}{N^\alpha}}(x_1), \Psi_{S, \frac{1}{N^\alpha}}(x_2), \dots, \Psi_{S, \frac{1}{N^\alpha}}(x_n)),$$

which leads us to the analogous conclusion that the power LCM matrix $[S]_{N^\alpha}$ is invertible if and only if $\Psi_{S, \frac{1}{N^\alpha}}(x_i) \neq 0$ for all $i = 1, 2, \dots, n$.

In 2002 Hong [8] raised the following conjecture concerning power LCM matrices.

Please cite this article in press as: M. Mattila et al., Some further applications of a lattice theoretic method in the study of singular LCM matrices, Linear Algebra Appl. (2023), <https://doi.org/10.1016/j.laa.2023.04.027>




Fig. 11. Illustration of the semilattice structure used in the proof of Theorem 6.1 and the Möbius function values $\mu_S(x_i, x_n)$ that are also needed in the calculation of the function $\Psi_{S, \frac{1}{N^\alpha}}(x_n)$.

Conjecture 6.1 (cf. [8], Conjecture). *If t is a given positive integer, then there exists a positive integer $k(t)$ (depending only on t) such that the power LCM matrix $[(\text{lcm}(x_i, x_j))^t]$ defined on any GCD closed set $S = \{x_1, x_2, \dots, x_n\}$ is nonsingular for all $n \leq k(t)$. But for any $n \geq k(t) + 1$ there exists a GCD closed set $S = \{x_1, x_2, \dots, x_n\}$ such that the matrix $[(\text{lcm}(x_i, x_j))^t]$ is singular.*

Known results for usual LCM matrices imply that $k(t) \geq 8$ for all $t \geq 2$, see [3]. The existence of $k(t)$ appears to be a highly nontrivial problem. However, we can use the cube semilattice to prove the following result that is closely related to the above conjecture.

Theorem 6.1. *If $M \geq 1$ is an arbitrary real number, then there exists a GCD closed set $S = \{x_1, x_2, \dots, x_n\}$ and real number α_0 such that $\alpha_0 > M$ and the power LCM matrix $[(\text{lcm}(x_i, x_j))^{\alpha_0}]$ is singular.*

Proof. Let $k \geq 2$ be a positive integer such that $\frac{\ln k}{\ln 2} \geq M$. Let p_1, p_2, \dots, p_{k-1} be sufficiently large prime numbers (to be specified later). Let us consider the set

$$S = \{1, 2, 3, 5, 6, 10, 15, 2p_1, 2p_2, \dots, 2p_{k-1}, 30p_1p_2 \cdots p_{k-1}\}$$

with $n = k + 7$ elements. We obtain the following semilattice structure shown in Fig. 11. The corresponding Möbius function values $\mu_S(x_i, x_n)$ are also shown in the same figure.

Now we write

$$h(\alpha) := \Psi_{S, \frac{1}{N^\alpha}}(x_n) = \sum_{i=1}^n \frac{\mu_S(x_i, x_n)}{x_i^\alpha}$$

$$\begin{aligned}
 &= \frac{-1}{1^\alpha} + \frac{k}{2^\alpha} + \frac{1}{3^\alpha} + \frac{1}{5^\alpha} + \frac{-1}{6^\alpha} + \frac{-1}{10^\alpha} + \frac{-1}{15^\alpha} + \sum_{i=1}^{k-1} \frac{-1}{(2p_i)^\alpha} + \frac{1}{x_n^\alpha} \\
 &= \left(\underbrace{-1 + \frac{k}{2^\alpha}}_{\geq 0 \text{ when } \alpha \leq \ln k / \ln 2} \right) + \frac{1}{30^\alpha} \underbrace{(10^\alpha - 5^\alpha - 3^\alpha - 2^\alpha)}_{\geq 0 \text{ when } \alpha \geq 1} \\
 &\quad + \left(\underbrace{\frac{1}{5^\alpha} - \sum_{i=1}^{k-1} \frac{1}{(2p_i)^\alpha}}_{\geq 0 \text{ for suff. large } p_i} \right) + \frac{1}{x_n^\alpha}.
 \end{aligned}$$

We can see that $h(\alpha) > 0$ for all $1 \leq \alpha \leq \frac{\ln k}{\ln 2}$. In particular, since $1 \leq M \leq \frac{\ln k}{\ln 2}$, we have $h(M) > 0$.

On the other hand $h(\alpha)$ is a continuous function of α and we have

$$\lim_{\alpha \rightarrow \infty} h(\alpha) = -1 + \lim_{\alpha \rightarrow \infty} \sum_{i=2}^n \underbrace{\frac{\mu_S(x_i, x_n)}{x_i^\alpha}}_{\rightarrow 0} = -1.$$

It now follows from Bolzano’s theorem that $h(\alpha_0) = 0$ for some $\alpha_0 > M$. \square

Declaration of competing interest

The authors have no conflicts of interest to declare.

Data availability

No data was used for the research described in the article.

References

- [1] E. Altınışık, T. Altıntaş, A note on the singularity of LCM matrices on GCD-closed sets with 9 elements, *J. Sci. Arts* 3 (40) (2017) 413–422.
- [2] K. Bourque, S. Ligh, On GCD and LCM matrices, *Linear Algebra Appl.* 174 (1992) 65–74.
- [3] W. Cao, On Hong’s conjecture for power LCM matrices, *Czechoslov. Math. J.* 57 (2007) 253–268.
- [4] P. Haukkanen, M. Mattila, J. Mäntysalo, Studying the singularity of LCM-type matrices via semi-lattice structures and their Möbius functions, *J. Comb. Theory, Ser. A* 135 (2015) 181–200.
- [5] P. Haukkanen, M. Mattila, J. Mäntysalo, Studying the inertias of LCM matrices and revisiting the Bourque-Ligh conjecture, *J. Comb. Theory, Ser. A* 171 (2020) 105161.
- [6] P. Haukkanen, J. Wang, J. Sillanpää, On Smith’s determinant, *Linear Algebra Appl.* 258 (1997) 251–269.
- [7] S. Hong, On the Bourque-Ligh conjecture of least common multiple matrices, *J. Algebra* 218 (1999) 216–228.
- [8] S. Hong, Gcd-closed sets and determinants of matrices associated with arithmetical functions, *Acta Arith.* 101 (2002) 321–332.

- [9] S. Hong, Nonsingularity of least common multiple matrices on gcd-closed sets, *J. Number Theory* 113 (2005) 1–9.
- [10] S. Hong, Nonsingularity of matrices associated with classes of arithmetical functions on lcm-closed sets, *Linear Algebra Appl.* 416 (1) (2006) 124–134.
- [11] I. Korkee, M. Mattila, P. Haukkanen, A lattice-theoretic approach to the Bourque-Ligh conjecture, *Linear Multilinear Algebra* 67 (12) (2019) 2471–2487.
- [12] H.J.S. Smith, On the value of a certain arithmetical determinant, *Proc. Lond. Math. Soc.* 7 (1876) 208–212.