



# UNIVERSITY OF TAMPERE

This document has been downloaded from  
Tampub – The Institutional Repository of University of Tampere

*Publisher's version*

Authors: Sumanen Markku, Suominen Sakari, Koskenvuo Markku,  
Sillanmäki Lauri, Mattila Kari

Name of article: Occurrence of symptoms and depressive mood among working-  
aged coronary heart disease patients.

Year of  
publication: 2004

Name of journal: Health and Quality of Life Outcomes

Volume: 2

Number of issue: 60

Pages: 1-7

ISSN: 1477-7525

Discipline: Medical and Health sciences / Forensic science and other medical  
sciences

Language: en

School/Other  
Unit: School of Medicine

URL: <http://www.hqlo.com/content/2/1/60>

URN: <http://urn.fi/urn:nbn:uta-3-621>

DOI: <http://dx.doi.org/10.1186/1477-7525-2-60>

All material supplied via TamPub is protected by copyright and other intellectual property rights, and duplication or sale of all part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorized user.

Research

Open Access

## Occurrence of symptoms and depressive mood among working-aged coronary heart disease patients

Markku PT Sumanen\*<sup>1</sup>, Sakari B Suominen<sup>2</sup>, Markku J Koskenvuo<sup>3</sup>, Lauri H Sillanmäki<sup>3</sup> and Kari J Mattila<sup>4</sup>

Address: <sup>1</sup>Kangasala Health Centre, Finland, <sup>2</sup>Department for Social and Health Services, University of Turku, Finland, <sup>3</sup>Department of Public Health, University of Turku, Finland and <sup>4</sup>Medical School, University of Tampere and Department of General Practice, Hospital District of Pirkanmaa, Finland

Email: Markku PT Sumanen\* - markku.sumanen@kolumbus.fi; Sakari B Suominen - suominen@utu.fi;

Markku J Koskenvuo - markku.koskenvuo@pp.inet.fi; Lauri H Sillanmäki - lahesi@utu.fi; Kari J Mattila - kari.j.mattila@uta.fi

\* Corresponding author

Published: 08 November 2004

Received: 02 June 2004

*Health and Quality of Life Outcomes* 2004, **2**:60 doi:10.1186/1477-7525-2-60

Accepted: 08 November 2004

This article is available from: <http://www.hqlo.com/content/2/1/60>

© 2004 Sumanen et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** The typical symptoms of coronary heart disease (CHD), chest pain and breathlessness, are well-known. They are considered quite dramatic, and can thus be fairly reliably mapped by a survey. However, people might have other clearly unpleasant symptoms impairing quality of life. The aim of this study is to evaluate the appearance of these complaints of working-aged people with self-reported CHD.

**Methods:** The study consists of a postal questionnaire of randomly selected Finns in age groups 30–34, 40–44 and 50–54, a response rate of 39% (N = 15,477). The subjects were asked whether or not a doctor had told them that they had angina pectoris or had had myocardial infarction. Four randomly selected age and sex matched controls were chosen for every patient. The occurrence of self-reported dyspnoea, chest pain during anger or other kind of emotion, palpitation and perspiration without physical exercise, irregular heartbeats, flushing, trembling of hands and voice, jerking of muscles, depression and day-time sleepiness were examined. Odds ratios (OR) with 95% confidence intervals (CI), between occurrence of symptoms and CHD with and without heart infarction, were computed by multivariate logistic regression analysis.

**Results:** The sample eventually comprised 319 CHD patients. Dyspnoea, chest pain during anger or other kind of emotion, palpitation, perspiration without physical exercise, irregular heartbeats daily or almost daily, trembling of hands and voice, and jerking of muscles occurred statistically significantly more frequently among CHD patients than among controls. The CHD patients also reported more depressive mood according to Beck's inventory scores and poorer sleep and more frequent day-time sleepiness than controls. In the multivariate logistic regression analysis chest pain during anger or other kind of emotion (ORs 4.12 and 3.61) and dyspnoea (ORs 2.33 and 3.81) were the symptoms most associated with CHD.

**Conclusions:** Working-aged people with self-reported coronary heart disease evince a number of symptoms limiting the quality of their every day life. This aspect should be paid attention to when evaluating functional capacity of these patients.

## Background

In Finland, as in most industrialised countries mortality from cardiovascular diseases has shown decreasing trends since around 1970 [1,2]. The Mini-Finland survey from the years 1979–80 revealed that the angina pectoris symptom (i.e. reported chest pain under physical strain) may already appear in both sexes at the age of 30, though it was not until the age of 65 that it becomes more common among men compared to women [3].

The typical symptoms of coronary heart disease (CHD), chest pain and breathlessness, are well-known. They are considered quite dramatic, and thus can be fairly reliably mapped by a survey [4]. However, coronary heart disease patients also have other complaints in respect of their health, for example fatigue and sleep problems [5]. It has also been estimated that 17% to 27% of patients with coronary artery disease have major depression and a significantly larger percentage has subsyndromal symptoms of depression [6].

The diagnosis of CHD is usually based on medical examinations or register data and not on what people by themselves experience. However, people act and use health services according to what they experience to suffer from and what they experience to limit their ability to work. Since CHD patients have many other symptoms than the traditional and well-known, it is important to know what the spectrum of symptoms and complaints among working-aged people with self-reported coronary heart disease is in relation to functional capacity.

## Methods

### Design

The Health and Social Support study (HeSSup) is a prospective etiological follow-up study on the psychosocial health of the Finnish working-aged population. The HeSSup population consisted of a random sample of 39,563 individuals drawn from the Finnish Population Register in three age groups: 30–34, 40–44, and 50–54. The survey was carried out by postal questionnaire. Forms were returned by 15,477 individuals (approximately 5,000 in each age group), a response rate of 38.9% (37.6% in 30–34, 37.9% in 40–44, and 41.1% in 50–54).

The sample was subjected to a thorough analysis of non-response [7]. The analysis was made using the official statistics of the Finnish population for the corresponding age groups in 1998 to assess whether the study population adequately represented the Finnish population. Diagnosed epilepsy and diagnosed hypertension were selected to represent chronic diseases. The major reasons for refuse were the length of the questionnaire and above all suspicion of the purpose behind the request for written consent. Less educated, divorced, widowed, unemployed and

those on disability pension were least willing to participate. Differences in physical conditions between the study participants and the whole population were, however, small. It was also noted that people suffering from hypertension returned the questionnaire somewhat less readily than others.

### Material and methods

The subjects were asked whether or not a doctor had told them that they had angina pectoris or had had myocardial infarction. The perceived state of health was determined according to Likert's five-step scale (good, quite good, fair, rather poor and poor). In order to avoid small frequencies in the analyses this was modified to a three-step scale (good, fair, poor).

The appearance of dyspnoea was categorised into four degrees of difficulty according to a widely used cardiovascular survey method [4]. Persons who suffered from dyspnoea when walking uphill or upstairs comprised the group of mild symptoms. Those out of breath when walking on level ground at normal speed with other people of the same age comprised the intermediate group, those who had to stop walking on level ground due to breathlessness comprised the group of difficult dyspnoea and those who became breathless even while standing still or while washing or dressing themselves, comprised the group of extremely difficult symptoms.

Participants were also asked whether or not they had experienced daily or weekly chest pain during anger or other kind of emotion, palpitation and perspiration without physical exercise, flushing, trembling of hands and voice, and jerking of muscles. Irregular heartbeats daily or weekly were asked, too. Depression was estimated by Beck's [8] depression scale ranging from 0 to 63. The normal score on this scale is below 10. In mild depression the scores are between 10 and 19 [9]. It was also asked how well and how many hours a day the participants had usually slept and how often they had felt day-time sleepiness, which when occurring daily or almost daily has been proved to be associated with depression, insomnia and breath interruptions during sleep [10].

### Analyses

In interpretation of results the coronary heart disease (CHD) patients were divided into two groups. The first comprised coronary patients not having had heart infarction (angina pectoris group) and the second patients having had heart infarction (infarction group). In order to have the best available comparison groups, four randomly selected age and sex matched controls for comparison were selected for every patient. Thus there were altogether 740 controls for the angina pectoris group and 536 con-

**Table 1: The coronary heart disease patients studied according to age and gender**

Age group	Angina pectoris group				Myocardial infarction group			
	Women	Men	Total	%	Women	Men	Total	%
	N	N	N		N	N	N	
30-34	16	14	30	16	4	3	7	5
40-44	20	28	48	26	3	9	12	9
50-54	47	60	107	58	22	93	115	86
Total	83	102	185	100	29	105	134	100

**Table 2: Prevalence of self-reported coronary heart disease according to age and gender**

Age group	Angina pectoris		Myocardial infarction	
	Women	Men	Women	Men
	%	%	%	%
30-34	0.5	0.7	0.1	0.2
40-44	0.7	1.4	0.1	0.4
50-54	1.7	2.8	0.7	4.0

trols for the infarction group. Stroke was ruled out in the control groups, but otherwise there were no differences between the CHD groups and their respective controls.

The associations, odds ratio (OR) with 95% confidence intervals (CI), between symptoms and coronary heart disease with and without heart infarction, were computed by multivariate logistic regression analysis. The analyses were made using the SAS System for Windows, release 8.2/2000.

## Results

The data comprised 319 patients: 185 coronary heart disease patients who had not experienced heart infarction (55.1% were men) and 134 patients who had (78.4% were men) (Table 1). Most of the CHD patients were in the oldest age group, and almost 90% of those who had had a heart infarction were in the age group 50-54. In all age groups the prevalence of self-reported CHD was higher among men than among women (Table 2).

### Perceived state of health

State of health was perceived as good or quite good by 37.3% in the angina group and by 24.6% in the infarction group. The corresponding figures in the control groups were 76.0% and 67.4%, the differences being statistically significant ( $p < 0.001$ ). State of health was perceived as

poor or rather poor by 28.1% in the angina group and by 32.8% in the infarction group.

### Symptoms and complaints

At least mild breathlessness occurred in two thirds of the angina group and three fourths of the infarction group (Table 3). Difficult or extremely difficult breathlessness was reported by 20.2% in the angina group and by 27.8% in the infarction group. The corresponding figures in the control groups were 1.4% and 4.0%, the differences also being statistically significant ( $p < 0.001$ ). Chest pain during anger or any kind of emotion, palpitation and perspiration without physical exercise, irregular heart beats, and jerking of muscles were all both daily and weekly statistically significantly more common among CHD patients than among controls. Almost daily CHD patients reported more daytime sleepiness and trembling of hands and voice than controls. CHD patients also slept more poorly than controls, and sleeping hours  $\leq 6$  in a day was more common among them than among controls. CHD patients scored higher on the depression scale than the controls, the average score being 10.2 (95% CI 9.0-11.4) in the angina group and 5.8 (95% CI 5.4-6.2) in the control group. In the infarction group the average score was 9.7 (95% CI 8.4-11.0). The corresponding figure in the control group was 5.9 (95% CI 5.4-6.5). In both coronary heart disease groups at least mild depression was twice as common as among controls. Flushing was the only

complaint, which was not statistically significantly more common among CHD patients than among controls.

### ORs of reported symptoms

In the multivariate logistic regression analysis chest pain during anger or other kind of emotion and dyspnoea were the symptoms most associated with CHD (Table 4). Irregular heart beats and perspiration without physical exercise were also strongly associated with heart infarction, but not with CHD without heart infarction. On the other hand, jerking of muscles was strongly associated with CHD without heart infarction, but not with heart infarction.

### Discussion

The principal finding in this study was that many working-aged coronary heart disease patients experience unpleasant symptoms such as dyspnoea, chest pain during anger or emotion, irregular heart beats, perspiration without physical exercise, and jerking of muscles. In addition, the frequency of most of the self-reported symptoms among the study population is higher also in respect of

those symptoms, which would not be expected at least among those CHD patients whose disease is in good balance. Working-aged CHD patients may be regarded as a special group compared with the main part of CHD patients who are already by age entitled to a pension. It is likely that working-aged CHD patients have experienced the most widespread and intense exposure to risk factors, which thus has caused them this disease among the first ones within their age group. As working-aged they are wished, however, to return back to normal life and work as soon as possible. According to our study they still have a lot of symptoms concerning their every day life, which harm their recovery and rehabilitation. It is also noteworthy that many of the working aged CHD patients are still in working life. Although chest pain and dyspnoea do not prevent them to work at customer service, many of the symptoms such as trembling of hands interfere their normal jobs while appearing mostly in rest. Trembling of hands and voice are also very irritating symptoms, and they may be considered shaming. Thus they may interfere social life and reduce the quality of life.

**Table 3: Occurrence (%) of symptoms and complaints in coronary heart disease (CHD) patients and the control population**

	Angina pectoris		Controls		Myocardial infarction		Controls	
	N = 177-185	N = 728-736			N = 129-134	N = 514-525		
	%	%	p		%	%	p	
At least mild dyspnoea (Rose and Blackburn 1968)	66.5	33.2	<0.001		75.4	35.3	<0.001	
Chest pain during anger or emotion								
Almost daily	12.2	1.0	<0.001		16.3	0.8	<0.001	
Weekly	12.8	2.5	<0.001		16.3	3.3	<0.001	
Palpitation without physical exercise								
Almost daily	14.1	3.4	<0.001		20.8	3.5	<0.001	
Weekly	15.3	5.2	<0.001		16.9	5.4	<0.001	
Perspiration without physical exercise								
Almost daily	22.7	9.7	<0.001		26.0	11.0	<0.001	
Weekly	18.2	10.2	0.003		16.8	8.3	0.001	
Irregular heart beats								
Almost daily	15.7	3.5	<0.001		24.0	3.7	<0.001	
Weekly	12.4	6.1	0.004		14.0	4.4	<0.001	
Depression (Beck $\geq$ 10)	41.6	20.4	<0.001		43.3	21.0	<0.001	
Sleeping hours $\leq$ 6 in a day	16.8	10.3	0.015		19.4	10.1	0.003	
Poor sleep usually	25.0	15.4	0.002		32.1	13.0	<0.001	
Daytime sleepiness								
Almost daily	33.2	11.2	<0.001		32.1	14.9	<0.001	
Flushing								
Almost daily	11.1	7.0	0.066		10.9	7.6	0.228	
Weekly	12.8	7.1	0.014		9.3	8.0	0.624	
Trembling of hands								
Almost daily	13.8	2.3	<0.001		9.2	3.3	0.004	
Weekly	8.3	4.7	0.053		10.7	4.0	0.003	
Trembling of voice								
Almost daily	3.3	1.1	0.030		5.3	1.2	0.002	
Weekly	3.3	1.2	0.049		3.8	2.3	0.341	
Jerking of muscles								
Almost daily	13.7	2.7	<0.001		12.3	4.4	0.001	
Weekly	11.5	2.7	<0.001		9.2	3.8	0.011	

The study material may be considered representative of the Finnish working-aged population, although the response rate was only 39%. Careful non-response analysis indicated that respondents and non-respondents were comparable in respect of the most important demographic variables [7]. It is possible that CHD patients respond to the questionnaire more actively than other people. On the other hand, there are certainly those among CHD patients who neglect their disease and are not willing to respond. However, we do not know for sure whether there is an over or under estimation of the associations, but we can presume that these two factors compensate each other. Moreover, it is unlikely that the principal association studied, i.e. the association between CHD and appearance of symptoms would be a substantially different one in non-participants. The findings reflect the respondents' own conception of their symptoms. The own conception of symptoms is important, since according to findings from a 3 years' follow-up of 4,000 men, self-reported coronary heart disease predicts very strongly a new coronary event [11]. In addition, the presence of anginal symptoms may be an important independent correlate of prognosis in patients with CHD [12].

Our method to determine the existence of CHD is based on the patients' report on whether a doctor had told them that they suffered from this particular disease. Thus, we cannot know for sure the accuracy of the information reported. Nowadays, CHD is rare among young people [1,2]. In our data there still is some. It is possible that there is a combination of several risk factors in the background.

There are validated instruments, such as the generic The Short Form 36 Health Survey [13] (SF-36), the Nottingham Health Profile [14], and the Seattle Angina Questionnaire [15], to investigate health-related quality of life. Our method to examine the subject was to ask about complaints and symptoms. Most of these questions have been successfully used in cardiovascular surveys [4] and in the Finnish Twin Cohort Study [16]. In addition, mood was estimated according to the Beck's depression scale [8].

The occurrence of dyspnoea and chest pain even during anger or other kind of emotion may be considered a finding that was expected. In an American study on the care of coronary heart patients at the emergency department the most frequently reported symptom was chest pain (70% among men and 71% among women) and dyspnoea (30% men and 29% among women) [17]. The typical chest pain is also a symptom more predictive of an acute coronary attack in working-aged than in older patients [18].

It was suggested as far back as 1987 that palpitations are not an independent risk factor for increased cardiac morbidity or mortality [19]. However, according to a British study those experiencing palpitations at work and while asleep were more likely to have a cardiac cause for their palpitations [20]. Our finding was that working-aged CHD patients report palpitations more often than the control population.

**Table 4: Age- and sex-matched ORs with 95% CI in the multivariate logistic regression analysis for reported symptoms of coronary heart disease (CHD) without and with heart infarction. All of these symptoms were in the same model. Statistically significant associations are bolded.**

	Angina pectoris	Myocardial infarction
	OR (95% CI)	OR (95% CI)
Dyspnoea	<b>3.81 (2.16–6.72)</b>	<b>2.33 (1.21–4.50)</b>
Chest pain during anger or emotion*	<b>3.61 (1.68–7.77)</b>	<b>4.12 (1.72–9.84)</b>
Palpitation without physical exercise*	1.19 (0.55–2.59)	1.39 (0.57–3.39)
Irregular heart beats*	1.46 (0.69–3.08)	<b>3.12 (1.28–7.60)</b>
Perspiration without physical exercise*	1.44 (0.87–2.38)	<b>2.19 (1.15–4.14)</b>
Flushing*	1.11 (0.62–1.97)	2.02 (0.89–4.59)
Trembling of hands*	1.36 (0.58–2.93)	1.01 (0.37–2.67)
Trembling of voice*	2.58 (0.85–7.87)	1.17 (0.29–4.69)
Jerking of muscles*	<b>2.44 (1.20–4.96)</b>	1.37 (0.55–3.41)
Depression (Beck $\geq$ 10)	1.57 (0.99–2.48)	1.01 (0.52–1.98)
Poor sleep	1.30 (0.74–2.27)	1.39 (0.70–2.77)
Daytime sleepiness	1.24 (0.76–2.01)	1.41 (0.78–2.56)
Sleeping hours $\leq$ 6 in a day	1.21 (0.64–2.26)	1.08 (0.50–2.33)

\* almost daily or weekly

The high occurrence of trembling of hands and voice, and jerking of muscles may be considered an unexpected finding. Most CHD patients use beta-blocking medicines, which in addition to protecting the heart muscle also reduce the adrenergic stimulation and thus relieve the symptom of trembling.

Concerning depression our findings support those of previous studies. One out of four of symptomatic coronary heart disease patients have namely been found to have a probable depressive disorder, but none of them had previously been identified as suffering from depression or been treated for this reason [21]. In primary care it is of vital importance to notice symptoms of this illness, since continuing depression has been found to be associated with increased risk of mortality among CHD patients following hospital discharge [22]. It has also been verified that depression is common after coronary heart disease events such as bypass grafting, coronary angioplasty, myocardial infarction and myocardial ischaemia [23]. In our study about 40% of CHD patients had at least minor depression compared with 20% among controls. The high depression rates are probably due to our method to diagnose depression at  $\geq 10$  points in Beck's inventory scale. Thus we do not think there are any selection bias, since the controls were randomly selected. Furthermore, it is not probable that depressed people respond to our questionnaire more readily than people not suffering from low mood.

It was also of noteworthy that daytime sleepiness was connected with coronary heart disease. A cross-sectional study of 5,419 Finnish adult men found a higher prevalence of diagnosed myocardial infarction among those who slept more than nine hours, whilst those sleeping less than six hours per night had more symptomatic coronary disease [24]. In a Swedish study concerning working-aged women poor sleep was associated with an increase in spasmodic chest pain and irregular heart beat [25], whereas in men an association between difficulties falling asleep and CHD mortality has been found [26].

From previous research we know that despite having survived a life-threatening clinical event, CHD patients appear to have continued adverse behaviours such as smoking, being obese and having frequent hangovers more than the control population [27]. The follow-up of our cohort will show in what extent the symptoms we found are indicators of increased risk of CHD among working-aged people and to what extent the symptoms are result of CHD and its care. In both cases particular attention should be paid to these aspects in primary care.

## Conclusions

According to the present findings many working-aged people with self-reported coronary heart disease perceive their state of health as poor or rather poor. They suffer from a wide range of symptoms limiting their every day life. It is noteworthy that many of these symptoms are not only irritating, but constitute a threat to health. The health related quality of life is poor among working-aged coronary heart disease patients.

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

MPTS drafted the manuscript; SBS participated in drafting of the manuscript; MJK participated in the design of the study and the statistical analyses; LHS participated in the statistical analyses; KJM conceived of the study, and participated in its design and co-ordination. All authors have read and approved the final manuscript.

## References

1. Uemura K, Pisa Z: **Trends in cardiovascular disease mortality in industrialised countries since 1950.** *World Health Stat Q* 1988, **41**:155-178.
2. Levi F, Lucchini F, Negri E, La Vecchia C: **Trends in mortality from cardiovascular and cerebrovascular diseases in Europe and other areas of the world.** *Heart* 2002, **88**:119-124.
3. Aromaa A, Heliövaara M, Impivaara O, Knekt P, Maatela J, Joukamaa M: **Health, functional limitations and need for care in Finland. Basic results from the Mini-Finland Health Survey (in Finnish).** *The Social Insurance Institution AL: 32, Helsinki and Turku* 1989.
4. Rose GA, Blackburn H: **Cardiovascular Survey Methods.** In *WHO Monograph Series Number 56* WHO Geneva; 1968.
5. Denollet J: **Health complaints and outcome assessment in coronary heart disease.** *Psychosom Med* 1994, **56**:463-474.
6. Rudisch B, Nemeroff CB: **Epidemiology of comorbid coronary artery disease and depression.** *Biol Psychiatry* 2003, **54**:227-240.
7. Korkeila K, Suominen S, Ojanlatva A, Ahvenainen J, Rautava P, Helenius H, Koskenvuo M: **Non-response and related factors in a nation-wide health survey.** *Eur J Epidemiol* 2001, **17**:991-999.
8. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J: **An inventory for measuring depression.** *Arch Gen Psychiat* 1961, **4**:54-63.
9. Varjonen J, Romanov K, Kaprio J, Heikkilä K, Koskenvuo M: **Self-rated Depression in 12 063 Middle-aged Adults.** *Nord J Psychiat* 1997, **51**:331-338.
10. Hublin C, Kaprio J, Partinen M, Heikkilä K, Koskenvuo M: **Daytime Sleepiness in an Adult Finnish Population.** *J Intern Med* 1996, **239**:417-423.
11. Koskenvuo M, Kaprio J, Rose RJ, Kesäniemi A, Sarna S, Heikkilä K, Langinvainio H: **Hostility as a risk factor for mortality and ischemic heart disease in men.** *Psychosom Med* 1988, **50**:330-340.
12. Cohn PF, Harris P, Barry WH, Rosati RA, Rosenbaum P, Waternaux C: **Prognostic importance of anginal symptoms in angiographically defined coronary artery disease.** *Am J Cardiol* 1981, **47**:233-237.
13. Ware JE, Donald Sherbourne C: **The MOS 36-item Short-Form Health survey (SF-36).** *Medical Care* 1992, **30**:473-483.
14. Hunt SM, McEwan T: **The development of a subjective health indicator.** *Soc of Health and Illness* 1980, **2**:231-246.
15. Spertus JA, Winder JA, Dewhurst TA, Deyo RA, Prodzinski J, McDonnell M, Fihn SD: **Development and evaluation of the Seattle Angina questionnaire: A new functional status measure for coronary artery disease.** *J Am Coll Cardiol* 1995, **25**:333-341.
16. Hublin C, Kaprio J, Partinen M, Koskenvuo M: **Insufficient sleep – a population-based study in adults.** *Sleep* 2001, **24**:392-400.

17. Milner KA, Funk M, Richards S, Wilmes RM, Vaccarino V, Krumholz HM: **Gender differences in symptom presentation associated with coronary heart disease.** *Am J Cardiol* 1999, **84**:396-399.
18. Milner KA, Funk M, Richards S, Vaccarino V, Krumholz HM: **Symptom predictor of acute coronary syndromes in younger and older patients.** *Nurs Res* 2001, **50**:233-241.
19. Knudson MP: **The natural history of palpitations in a family practice.** *J Fam Pract* 1987, **24**:357-360.
20. Summerton N, Mann S, Rigby A, Petkar S, Dhawan J: **New-onset palpitations in general practice: assessing the discriminate value of items within the clinical history.** *Fam Pract* 2001, **18**:383-392.
21. Valkamo M, Hintikka J, Niskanen L, Koivumaa-Honkanen H, Honkalampi K, Viinamäki H: **Depression and associated factors in coronary heart disease.** *Scand Cardiovasc J* 2001, **35**:259-263.
22. Lesperance F, Frasure-Smith N: **Depression in patients with cardiac disease: a practical review.** *J Psychosom Res* 2000, **48**:317-320.
23. Lehto S, Koukkunen H, Hintikka J, Viinamäki H, Laakso M, Pyörälä K: **Depression after coronary heart disease events.** *Scand Cardiovasc J* 2000, **4**:580-583.
24. Partinen M, Putkonen PT, Kaprio J, Koskenvuo M, Hilakivi I: **Sleep disorders in relation to coronary heart disease.** *Acta Med Scand Suppl* 1982, **660**:69-83.
25. Asplund R, Åberg H: **Sleep and cardiac symptoms amongst women aged 40-64 years.** *J Intern Med* 1998, **243**:209-213.
26. Mallon L, Broman JE, Hetta J: **Sleep complaints predict coronary artery disease mortality in males: a 12-year follow-up study of a middle-aged Swedish population.** *J Intern Med* 2002, **251**:207-216.
27. Sumanen M, Koskenvuo M, Immonen-Räihä P, Suominen S, Sundell J, Mattila K: **Secondary prevention of coronary heart disease is disappointing among patients of working-age.** *Fam Pract* 2004, **21**:304-306.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

